

**Diet and trophic relationships of top predators in the oceanic region of the  
archipelago of Madeira**

*“Documento Definitivo”*

**Doutoramento em Biologia e Ecologia das Alterações Globais**  
Especialidade em Biologia e Ecologia Marinha

Joana Chalbert Manso Romero

Tese orientada por:  
José Pedro Granadeiro  
Paulo Catry

Documento especialmente elaborado para a obtenção do grau de doutor



UNIVERSIDADE DE LISBOA

FACULDADE DE CIÊNCIAS



UNIVERSIDADE  
DE LISBOA

Faculdade de Ciências



universidade  
de aveiro

Departamento de Biologia

**Diet and trophic relationships of top predators in the oceanic region of the  
archipelago of Madeira**

**Doutoramento em Biologia e Ecologia das Alterações Globais**

Especialidade em Biologia e Ecologia Marinha

Joana Chalbert Manso Romero

Tese orientada por:

José Pedro Granadeiro

Paulo Catry

Júri:

Presidente:

- Doutora Sólveig Thorsteinsdóttir, Professora Associada com Agregação e Presidente do Departamento de Biologia Animal, da Faculdade de Ciências da Universidade de Lisboa

Vogais:

- Doutor Joan Navarro, Investigador, Instituto de Ciencias del Mar (ICM- CSIC), Espanha
- Doutor Frederico José Oliveira de Almada, Investigador, MARE - Centro de Ciências do Mar e do Ambiente do ISPA - Instituto Universitário de Ciências Psicológicas, Sociais e da Vida;
- Doutor Amadeu Mortágua Velho da Maia Soares, Professor Catedrático, Departamento de Biologia da Universidade de Aveiro;
- Doutora Maria da Luz da Costa Pereira Mathias, Professora Catedrática, Faculdade de Ciências da Universidade de Lisboa;
- Doutor José Pedro Oliveira Neves Granadeiro, Professor Auxiliar com Agregação, Faculdade de Ciências da Universidade de Lisboa, Orientador;
- Doutor Carlos António da Silva Assis, Professor Auxiliar, Faculdade de Ciências da Universidade de Lisboa.

Documento especialmente elaborado para a obtenção do grau de doutor

Fundação para a Ciência e a Tecnologia (FCT) – Bolsa de Doutoramento (PD/BD/127810/2016)

Fundação para a Ciência e a Tecnologia (FCT) – Projecto Oceanwebs (PTDC/MAR-PRO/0929/2014)

Fundação para a Ciência e a Tecnologia (FCT) – Projecto Seamigrant (PTDC/BIA-ANM/3743/2014)

Fundação para a Ciência e a Tecnologia (FCT) – Financiamento estratégico atribuído ao CESAM  
(UIDB/50017/2020, UIDP/50017/2020)

Fundação para a Ciência e a Tecnologia (FCT) – Financiamento estratégico atribuído ao MARE  
(UIDB/04292/2020, UIDP/04292/2020)

Comissão Europeia – Projecto LIFE RECOVER NATURA (LIFE12 NAT/PT/000195)

Agência Regional para o Desenvolvimento da Investigação, Tecnologia e Inovação (ARDITI) – Projecto  
(M1420-01-0145-FEDER-000001-Observatório Oceânico da Madeira-OOM)



**Nota prévia:** A presente tese apresenta artigos científicos já publicados (capítulos 2-5) ou por submeter para publicação (capítulo 6), de acordo com o previsto no nº 2 do artigo 25º do Regulamento de Estudos de Pós-Graduação da Universidade de Lisboa, publicado no Diário da República II série - Nº 60 - 26 de março de 2018. Uma vez que estes trabalhos foram realizados em colaboração o candidato esclarece que participou integralmente na conceção dos trabalhos, obtenção dos dados, análise e discussão dos resultados, bem como na redação dos manuscritos.

Lisboa, dezembro de 2020  
Joana C. M. Romero



*Para os meus pais,  
Por sempre me apoiarem e acreditarem em mim.*

*“O rapaz, porém, pensava no seu tesouro. Quanto mais perto ficava do seu sonho, mais difíceis se tornavam as coisas. Já não funcionava aquilo que o velho rei tinha chamado “sorte de principiante”. O que funcionava, sabia ele, era o teste da persistência e da coragem de quem procura a sua Lenda Pessoal. Por isso ele não podia apressar-se, nem ficar impaciente. Se agisse assim, ia acabar por não ver os sinais que Deus tinha posto no seu caminho”*

O Alquimista, Paulo Coelho





## **Agradecimentos/ Acknowledgements**

Em primeiro lugar, gostaria de agradecer aos meus orientadores, José Pedro Granadeiro e Paulo Catry, por todo o apoio e ajuda prestados durante estes quatro anos de doutoramento. Agradeço a confiança que depositaram em mim e todo o tempo que disponibilizaram para me ensinar e guiar. Estes quatro anos convosco foram uma constante aprendizagem, tanto no campo e laboratório, como a nível académico. Só posso esperar ter conseguido absorver todo o conhecimento e ensinamentos que tinham para me dar. Também vos quero agradecer pelos momentos de amizade e pelos “dar na cabeça” que tão importantes foram para o meu crescimento enquanto investigadora e pessoa. Por isto tudo, um muito obrigada!

Também gostaria de agradecer a todas as entidades que estiveram envolvidas nos trabalhos desta tese ou que de alguma forma possibilitaram a realização da mesma. À Fundação para a Ciência e Tecnologia e ao Centro de Estudos do Ambiente e do Mar, pela atribuição da bolsa que me permitiu ingressar nesta aventura. À Faculdade de Ciências da Universidade de Lisboa por me albergar. Ao Instituto das Florestas e Conservação da Natureza, nomeadamente ao Doutor Paulo Oliveira, Dra. Dília Menezes, Dra. Carolina Santos, Dra. Nádia Coelho e Dr. Pedro Sepulveda, pela ajuda financeira e logística prestada nos diferentes trabalhos realizados nas Desertas e Selvagens. À Direcção Regional das Pescas da Madeira, nomeadamente ao Dr. João Delgado e Dra. Graça Faria, pela disponibilização das instalações e apoio logístico e financeiro providenciados ao projecto. Gostaria também de agradecer a ajuda e apoio indispensáveis da Doutora Margarida Hermida e das bolseiras Bárbara Cavaleiro e Raquel Segur, na recolha de amostras e dados de atuns e ruama na Madeira.

Todo o trabalho que realizei ao longo destes quatro anos não teria tido o mesmo sabor sem a presença, companheirismo e amizade do pessoal da FCUL. Filipe Moniz, muito obrigada pela ajuda dada na Selvagem e no laboratório, pelo passar de conhecimento sobre técnicas de captura de gaivotas e manuseamento de cagaras, “Bebe chico!”, mas acima de tudo pela amizade. À minha PhD-sister, Marie Claire Gatt, muito obrigada pelos momentos memoráveis na Selvagem e nas viagens de navio entre a Madeira e Selvagem (talvez nos queiramos esquecer desta parte), pelas conversas e trocas de ideias sobre as nossas teses, pela ajuda com o R sempre que ele não colaborava comigo. Também gostava de agradecer à Letizia Campioni, a minha companhia na sala. Muito obrigada pelos conselhos e “brainstormings” e, acima de tudo, pela ajuda e companhia na

Selvagem. Ao meu companheiro de laboratório, Fernando Madeira. Antes de tudo quero pedir-te desculpa pelos maus cheiros que tiveste de suportar, cenas de atuns! Muito obrigada pelas horas infindáveis de conversa e parvoíce, sem dúvida foram a melhor companhia que podia ter tido para suportar os longos dias e meses a identificar vértebras e coisas afins. Um grande obrigada à Carolina Vieira pela sua dedicação e ajuda na identificação de fito- e zooplâncton, pela sua vontade de aprender e fazer melhor. Também gostaria de agradecer a todos os outros membros do Wings over water, ao Francesco Ventura, Ana Cordeiro, Edna Correia, Jorge Gutiérrez, Patrícia Pedro, Mohamed Henriques, Maria Alho, Carolina Vieira, João Belo, João Paulino, Martin Beal, Anne-Sophie, Mónica Silva, Vera Nunes, Rita Patrício, Ricardo Furtado, Amadú Insambé, Tumbulo Garcia, Daniel Lopes, Teresa Catry e Rui Rebelo, pelas trocas de ideias, dicas e conhecimento trocados ao longo destes anos.

Não poderia deixar de agradecer à Maria Alho por estar sempre presente em todo este processo, às vezes, um pouco doloroso. Obrigada pelas conversas sobre tudo e sobre nada, à partilha das alegrias e das coisas menos boas, por aturares os meus desesperos e inseguranças. Por estares sempre lá para ouvir-me e por seres a pessoa fantástica que és. Sem dúvida, ajudaste-me a aligeirar as “dores” do doutoramento.

A todos os que me ajudaram no campo e laboratório ao longo destes anos, muito obrigada. Ao João Morgado, Estefania, Paulo Catry, Marie Claire, Maria Alho, Fernando Madeira, Filipe Moniz, Letizia Campioni, Diana Barão, José Pedro Granadeiro, Teresa Catry, Carolina Vieira, Mafalda Pádua, Gabriel Mendes, Rafael Luís e Washington Gouveia. Um grande obrigada também a todos os vigilantes da natureza que ajudaram em toda a logística relacionada com as estadias na Selvagem e Desertas: Jacques da Mata, Fernando Vieira, Carlos Clemente, Carlos Santos, Manuel José, Isamberto Silva, João Paulo Mendes, Maurício Baptista e Marco Camacho.

Um agradecimento muito especial à Dona Augusta e Pierre por todo o carinho e atenção. Por me receberem sempre tão bem na Guiné mas, especialmente, nesta última visita, por me terem proporcionado o melhor ambiente que poderia ter para “atacar” os últimos capítulos da tese. Obrigada por me mostrarem sítios maravilhosos, pelas ostradas de sábado e todas as iguarias preparadas pela Dona Augusta. Um obrigada muito grande também para a Fanceni, Lu e Lara. A vossa companhia foi sempre uma lufada de ar fresco para começar mais uma semana de trabalho. Pelos domingos preguiçosos no terraço e conversas sobre a vida e o mundo, aos nossos jogos da Disney e, sobretudo, ao “Mateus”. Que venham muitos mais momentos como estes.

À minha família, por todo o apoio, amor e carinho que sempre me deram. Por acreditarem em mim e darem-me a força que preciso para continuar. Vocês são e serão sempre a minha “casa” e para onde eu vou querer sempre voltar. Um especial obrigada ao meu irmão por me espicaçar a ser melhor, por me incentivar a sair da minha zona de conforto. Obrigada por estares sempre lá para mim. Obrigada ao meu sobrinho do coração, Du (sim, para mim serás sempre Du, por muito que cresças), por seres o único que realmente me conseguia tirar a cabeça do trabalho e problemas com a tua energia, vivacidade e alegria. Nunca deixes de ser o meu “louquinho”. Aos meus pais... Não tenho palavras para vos descrever o quão agradecida estou por tudo o que fizeram por mim. A pessoa que sou hoje deve-se aos vossos esforços, dedicação e amor incondicional e, por isso, devo-vos tudo. Obrigada por terem acreditado e puxado por mim e, acima de tudo, por me terem proporcionado a vida e experiências que tive até hoje.

Por fim, mas não menos importante, ao Mo. Obrigada pela pessoa que és e que tens sido para mim, pela tua amizade e amor. Obrigada por me teres “aturado” nestes quatro anos de ausências, muito trabalho e stress, mas também de alegria e novas conquistas. Obrigada pela tua compreensão e paciência mesmo nos dias mais difíceis, pela força e apoio que me deste e por me ajudares a levantar mesmo quando eu já não tinha forças. Obrigada por estares lá todos os dias mesmo quando a distância era grande. Obrigada pelos “brainstormings”, comentários aos manuscritos e por tomares conta de mim ao longo de todo este percurso. Obrigada ao meu companheiro, colega e “role-model”, não teria sido capaz sem ti a meu lado. Mais do que ontem e menos do que amanhã.



# Contents

<b>RESUMO.....</b>	<b>1</b>
<b>PALAVRAS-CHAVE .....</b>	<b>2</b>
<b>ABSTRACT .....</b>	<b>3</b>
<b>KEYWORDS .....</b>	<b>4</b>
<b>CHAPTER 1 - GENERAL INTRODUCTION .....</b>	<b>6</b>
<b>DIET AND FORAGING ECOLOGY IN MARINE ECOSYSTEMS .....</b>	<b>6</b>
<b>MARINE TROPHIC WEBS.....</b>	<b>8</b>
<b>THE OCEANIC REGION OF THE ARCHIPELAGO OF MADEIRA .....</b>	<b>10</b>
<b>THESIS OUTLINE .....</b>	<b>12</b>
<b>REFERENCES.....</b>	<b>13</b>
<b>CHAPTER 2 - DIET AND TROPHIC POSITION OF TWO MACKEREL SPECIES IN THE ARCHIPELAGO OF MADEIRA, PORTUGAL.....</b>	<b>20</b>
<b>ABSTRACT.....</b>	<b>20</b>
<b>KEYWORDS .....</b>	<b>20</b>
<b>INTRODUCTION .....</b>	<b>21</b>
<b>METHODS.....</b>	<b>22</b>
<b>RESULTS .....</b>	<b>25</b>
<b>DISCUSSION.....</b>	<b>33</b>
<b>ACKNOWLEDGEMENTS.....</b>	<b>35</b>
<b>REFERENCES.....</b>	<b>35</b>
<b>SUPPORTING INFORMATION.....</b>	<b>40</b>
<b>REFERENCES.....</b>	<b>43</b>
<b>CHAPTER 3 - TUNAS OFF NORTHWEST AFRICA: THE EPIPELAGIC DIET OF THE BIGEYE AND SKIPJACK TUNAS .....</b>	<b>46</b>
<b>ABSTRACT.....</b>	<b>46</b>
<b>KEY WORDS .....</b>	<b>46</b>
<b>INTRODUCTION.....</b>	<b>47</b>
<b>METHODS.....</b>	<b>48</b>
<b>RESULTS .....</b>	<b>52</b>
<b>DISCUSSION.....</b>	<b>59</b>
<b>ACKNOWLEDGEMENTS.....</b>	<b>62</b>
<b>REFERENCES.....</b>	<b>63</b>
<b>SUPPLEMENTARY MATERIAL.....</b>	<b>72</b>
<b>REFERENCES.....</b>	<b>76</b>
<b>CHAPTER 4 - A GULL THAT SCARCELY VENTURES ON THE OCEAN: YELLOW-LEGGED GULLS LARUS MICHAHELLIS ATLANTIS ON THE OCEANIC ISLAND OF MADEIRA .....</b>	<b>78</b>

<b>ABSTRACT.....</b>	<b>78</b>
<b>KEYWORDS .....</b>	<b>78</b>
<b>INTRODUCTION.....</b>	<b>79</b>
<b>METHODS.....</b>	<b>80</b>
<b>RESULTS .....</b>	<b>82</b>
<b>DISCUSSION.....</b>	<b>86</b>
<b>ACKNOWLEDGEMENTS.....</b>	<b>88</b>
<b>REFERENCES.....</b>	<b>89</b>
<b>SUPPLEMENTARY MATERIAL 1.....</b>	<b>94</b>
<b>REFERENCES.....</b>	<b>96</b>
<b>SUPPLEMENTARY MATERIAL 2.....</b>	<b>99</b>
 <b>CHAPTER 5 - SEABIRD DIET ANALYSIS SUGGESTS SUDDEN SHIFT IN THE PELAGIC COMMUNITIES OF THE SUBTROPICAL NORTHEAST ATLANTIC .....</b>	 <b>102</b>
<b>ABSTRACT.....</b>	<b>102</b>
<b>KEYWORDS .....</b>	<b>102</b>
<b>INTRODUCTION.....</b>	<b>103</b>
<b>METHODS.....</b>	<b>104</b>
<b>RESULTS .....</b>	<b>109</b>
<b>DISCUSSION.....</b>	<b>113</b>
<b>ACKNOWLEDGEMENTS.....</b>	<b>118</b>
<b>REFERENCES.....</b>	<b>118</b>
<b>SUPPLEMENTARY MATERIALS.....</b>	<b>126</b>
 <b>CHAPTER 6 - FOOD WEB STRUCTURE OF THE SUBTROPICAL OCEANIC ARCHIPELAGO OF MADEIRA, EASTERN NORTH ATLANTIC .....</b>	 <b>138</b>
<b>ABSTRACT.....</b>	<b>138</b>
<b>KEYWORDS .....</b>	<b>138</b>
<b>INTRODUCTION.....</b>	<b>139</b>
<b>METHODS.....</b>	<b>141</b>
<b>RESULTS .....</b>	<b>146</b>
<b>DISCUSSION.....</b>	<b>151</b>
<b>CONCLUSION .....</b>	<b>154</b>
<b>REFERENCES.....</b>	<b>155</b>
<b>SUPPLEMENTARY MATERIALS.....</b>	<b>161</b>
<b>REFERENCES.....</b>	<b>182</b>
 <b>CHAPTER 7 - GENERAL DISCUSSION .....</b>	 <b>216</b>
<b>DIET, PREY ABUNDANCE AND COMMUNITY ASSESSMENT .....</b>	<b>216</b>
<b>FOOD WEB STRUCTURE AND FISHERIES .....</b>	<b>218</b>
<b>FUTURE RESEARCH.....</b>	<b>220</b>
<b>REFERENCES.....</b>	<b>221</b>

## Resumo

Os estudos de dieta são um elemento importante na compreensão da ecologia de uma espécie, mas são muitas vezes negligenciados e a sua importância subestimada. A dieta de um organismo permite conhecer o habitat em que o animal vive e as suas adaptações ao meio ambiente, mas também permite saber sobre interações com outras espécies. É a combinação de estudos de dieta de várias espécies que permite construir cadeias alimentares e compreender a estrutura e o funcionamento dos ecossistemas. A aplicação deste tipo de estudos no meio marinho tem importância acrescida, uma vez que estudos baseados em observação directa não são suficientes. Esta importância é ainda mais acentuada nas regiões oceânicas, onde as densidades de organismos são baixas e o seu estudo é um desafio.

Esta tese tem como objetivo estudar a dieta e as relações tróficas entre os predadores de topo e as suas presas na região oceânica do arquipélago da Madeira. A dieta e as áreas de alimentação de espécies de importância ecológica e económica, como atuns, aves marinhas e pequenos peixes pelágicos, foram estudadas na tentativa de compreender a sua posição e o seu papel na cadeia alimentar deste ambiente oceânico.

A dieta de duas espécies de peixes pelágicos, a cavala *Scomber colias* e o carapau-negrão *Trachurus picturatus*, foi analisada com base na identificação do conteúdo estomacal de indivíduos capturados ao longo de um ano, nas proximidades da ilha da Madeira. Ambas as espécies são planctívoras e piscívoras, alimentando-se sobretudo de copépodes calanóides e ciclopóides, e de pequenos agulhões *Scomberesox saurus*, clupeídeos, apara-lápis *Macroramphosus scolopax* e mictofídeos.

A dieta do patudo *Thunnus obesus* e do gaiado *Katsuwonus pelamis*, foi estudada identificando o conteúdo estomacal de indivíduos apanhados na Madeira e confirmada através da análise de mercúrio nos tecidos dos atuns e de outros dois peixes epipelágicos, a bicuda *Sphyrna viridensis* e o charuteiro *Seriola rivoliana*. A dieta de ambas as espécies de atum é composta principalmente por espécies epipelágicas, como a cavala e o peixe-rei *Atherina* sp., o que contrasta com o comportamento e dieta mesopelágicos do patudo em outras regiões do mundo.

Também investigámos a utilização de recursos marinhos pela gaivota-de-patas-amarelas *Larus michahellis atlantis* no arquipélago da Madeira. Esta ave costeira mostrou uma grande dependência antropogénica, com elevado uso de ambientes terrestres e associados ao homem. As poucas vezes que utilizou o meio marinho foi, principalmente,

à noite e em associação com embarcações de cerco. A sua dieta refletiu este comportamento, sendo maioritariamente composta por resíduos e alguns peixes.

O estudo da variação na composição de espécies epipelágicas na vasta região da corrente das Canárias, foi realizado utilizando as aves marinhas como indicadores. Com recurso a viagens de alimentação e regurgitos de cagaras *Calonectris borealis* a nidificar nas Ilhas Selvagens, recolhidos por um período de sete anos, este estudo conseguiu detetar uma alteração na comunidade de peixes pelágicos, com um aumento acentuado da população de apara-lápis nos arredores das ilhas Selvagens, em 2017/2018.

Por fim, desenvolvemos um modelo baseado no equilíbrio de biomassas de espécies encontradas na Zona Económica Exclusiva do arquipélago da Madeira, utilizando o software Ecopath with Ecosim. Este capítulo reuniu as informações recolhidas nos outros capítulos e juntou-as aos dados disponíveis na literatura. O nosso modelo estimou uma elevada biomassa de produtores primários, zooplâncton, outros crustáceos e invertebrados, mas também de pequenos peixes pelágicos e mesopelágicos, que foram considerados o principal alimento de predadores de topo. O ecossistema caracterizou-se por um baixo número de ligações entre os níveis tróficos, que está relacionado com uma dieta mais especializada por parte de organismos como golfinhos, aves marinhas e grandes peixes pelágicos. Apesar de o ecossistema ter apresentado um nível trófico médio baixo, o nível trófico das pescas foi bastante alto por, na Madeira, predadores de topo como atuns e peixe-espada, serem os mais pescados. Os predadores de topo também foram considerados importantes modeladores do ecossistema, sendo designados por espécies-chave.

**Palavras-chave:** Dieta, Áreas de alimentação, Relações tróficas, Predadores de topo, Cadeia alimentar.



## Abstract

Diet studies are important for the understanding of the ecology of species, but commonly overlooked and their importance underestimated. The diet of an organism reveals its prey composition, but also its habitat and foraging behaviour. It also allows to have an insight on the interactions with other species and its surroundings. The combined information of diet studies of several species allows to build food webs and understand the structure and functioning of ecosystems. The application of these studies in the marine environment is particularly important, as pure observational studies are not enough. Such importance is even greater in oceanic regions where densities of organisms are low, and their study is challenging.

This thesis aims to study the diet and trophic relationships of top predators and their prey in the oceanic region of the archipelago of Madeira. The diet and foraging areas of ecologically and economically important species, like tunas, seabirds and small pelagic fish, were studied in the attempt to understand their position and role in the food web of this oceanic environment.

The diet of two species of mackerels, the Atlantic chub mackerel *Scomber colias* and the Blue jack mackerel *Trachurus picturatus*, was assessed using stomach contents of individuals caught throughout a year in the vicinities of Madeira island. These are planktivorous and piscivorous species, feeding mostly on calanoid and cyclopoid copepods, and on very small Atlantic saury *Scomberesox saurus*, clupeids, Longspine snipefish *Macroramphosus scolopax*, and myctophids.

The diet of the Bigeye tuna *Thunnus obesus* and Skipjack tuna *Katsuwonus pelamis*, was assessed using stomach contents and confirmed using mercury analysis in the tissues of tunas and two other epipelagic fish, the Yellowmouth barracuda *Sphyræna viridensis* and the Longfin yellowtail *Seriola rivoliana*. The diet of both species was mainly composed by epipelagic prey, like Atlantic chub mackerel and Sand smelts *Atherina* sp., which contrasts with the mesopelagic behaviour and diet of Bigeye tunas elsewhere.

We also investigated the use of marine resources by the Yellow-legged gull *Larus michahellis atlantis* in the archipelago of Madeira. This coastal bird showed a great anthropogenic dependence, with high use of terrestrial and human-associated environments. The few times it used the marine environment, it was mostly at night and

in association with purse-seine vessels. Its diet reflected this behaviour, being mainly composed by human residues and few fish.

The variation in the composition of epipelagic species through a period of seven years in the broad Canary current region was studied using foraging trips and regurgitations of Cory's shearwaters *Calonectris borealis* nesting on Selvagens Islands. This study was able to depict a shift in the community, with a steep increase of the population of Longspine snipefish in the surroundings of Selvagens islands in 2017/2018.

Finally, we developed a mass-balanced model of the Exclusive Economic Zone of the archipelago of Madeira, using the software Ecopath with Ecosim. This chapter brought together the information collected in the other chapters with data available in the literature. The Madeira system was characterized by high biomass of primary producers, zooplankton and other crustaceans and invertebrates, together with small epipelagic and mesopelagic fish, which were also the main prey of top predators. The food web was characterized by a more linear-like food chain, in opposition to a more web-like food chain, with a large proportion of more specialized organisms, like dolphins, shearwaters and large pelagic fish. Despite the low mean trophic level of the system, the mean trophic level of fisheries was very high, targeting mainly top predators, like tunas and Black scabbardfish, which were also the components with most impact in the ecosystem.

**Keywords:** Diet, Foraging areas, Trophic relationships, Top predators, Food web.

# CHAPTER 1

---



**Credits: Joana Romero**

## **General Introduction**

The study of oceanic environments has always represented a challenge for researchers due to the difficulties in sampling this ecosystem, but this is changing with the development of technology and increasing investment in the exploration of the ocean. Many advancements have been made in the understanding of the functioning of oceans, in terms of physical, chemical, and biological processes, but there is still a lot we do not understand. For one, there are still many questions on the impacts of global changes, pollution, and fisheries on the biodiversity and balance of marine ecosystems. One way to address these questions is by assessing the diet of predators, which are wide ocean wanderers, and their trophic relationships with the remaining species of the oceanic food web.

## **Diet and Foraging Ecology in Marine Ecosystems**

Diet studies are essential for the understanding of organisms and their adaptation to the environment. In addition of allowing to study the timing and frequency of feeding (e.g. Granadeiro et al. 1998), energy budgets (e.g. Doidge & Croxall 1985) and competition and interaction among species (e.g. Spitz et al. 2006), diet studies also provide information on prey communities, such as composition, distribution, age and reproduction (e.g. Di Benedetto & Siciliano 2007). Moreover, it can provide insights on the interactions of different species with the fisheries industry (e.g. Wallace et al. 2009), and on the impacts of global changes and human activities on marine species (e.g. Miller & Sydeman 2004).

There are several ways to study the diet of a species, one being the use of stomach contents. For most species the collection of stomach contents requires the animal to be dead, as it is the case of fish and cetaceans, but for others, like seabirds, it is possible to sample their contents in spontaneous regurgitations, or by stomach-flushing (Barrett et al. 2007). Diet analysis is also possible by simply collecting pellets (an agglomerate of hard structures) that are naturally regurgitated after feeding events. The use of stomach contents is usually the most complete method and with higher resolution to describe the diet composition of a species (Karnovsky et al. 2012), as it is often possible to count the number of ingested prey and to identify each organism using morphological characteristics. Even if more digested, prey can still be identified using hard parts, like bones, otoliths, cephalopod beaks, shells, etc. (Granadeiro & Silva 2000, Xavier & Cherel 2009). In more recent years, the development of polymerase chain reaction-based

techniques (PCR) and DNA barcoding have become a powerful ally of this method, making possible the identification of the prey consumed using only a small portion of tissue (Alonso et al. 2014, Correia et al. 2019). Metabarcoding analysis has also proved most useful in studies using faeces, for instance, as it allows to depict more than one prey taxa/species (Ford et al. 2016, Correia et al. 2019). The use of these techniques to identify prey is not always possible due to the high monetary costs they entail and the complexity of the methods. The size and weights of prey are also possible to take in most cases, either through direct observation or using pre-existing equations relating these variables with the size of hard structures (such as otoliths, vertebrae, beaks) of each prey. Still, the analysis of stomach contents present some drawbacks, like the underestimation of easily digested items and short-temporal representation of the diet, as contents only represent the last meal of the individual (Barrett et al. 2007). Another drawback is that it will only represent the prey of the last feeding event. Thus, a good sample size is required to describe the diet more faithfully.

As an attempt to avoid bias and obtain a more detailed and accurate interpretation of the diet, conventional sampling has been allied to the use of biochemical methods, like stable isotope analysis, fatty acid signature, or mercury analysis. Such assays provide tools to assess trophic interactions, diet data over space and time, and can even narrow down the foraging areas of the consumer (Barrett et al. 2007).

Stable isotope analysis has now been widely used as a tool in diet studies (Cherel et al. 2017, Denda et al. 2017, Morera-Pujol et al. 2018). This methodology requires the collection of tissues, like muscle, blood, feathers, among others, to estimate the nitrogen, carbon or sulfur ratios of the consumers. Isotopes accumulate in tissues in predictable ways, allowing to define the trophic level of the consumer using nitrogen, and to identify the feeding habitat (neritic vs. oceanic and epipelagic vs. demersal) of the prey through the carbon isotope ratio (Layman et al. 2012). Also, depending on the tissue used, it is possible to infer different time frames, from a specific period according to its turnover rate, like it happens when using feathers that were grown during a specific period, to months back in time, when using muscle, or to a few weeks or days ago, using different components of the blood of the study species (Layman et al. 2012). Because stable isotopes reflect the food assimilated by studied individuals, it does not carry the bias of different digestibility of prey. However, there is always the need for background information on the potential preys, their isotopic compositions, and foraging areas for a more truthful interpretation of the isotopic signature of the consumer.

Mercury analysis of tissues, per se, cannot infer the diet of individuals, but when associated with stomach contents or stable isotope analysis, they can be a valuable asset to distinguish between prey groups with different mercury concentrations (Di Benedetto et al. 2011). Mercury can be found in the environment in different ways, and once ingested, it has the capacity of magnifying and accumulating in the tissues of its consumers. Its concentration increases from consumer to consumer up to the top predator, which theoretically possess the highest concentrations of mercury of the food chain. Mercury concentrations are not only related to the level of pollution of a habitat, but also with depth. Mercury concentrations can help distinguish between a mesopelagic/bathypelagic and an epipelagic diet, as deep-ocean species have significantly higher concentrations of mercury than epipelagic species due to the higher rate of microbial-mediated methylation of mercury in sub-thermocline low oxygen waters (Choy et al. 2009). In conclusion, mercury is useful to distinguish prey of similar trophic levels and different depths.

Since the advent of very high frequency tracking (VHF), several technologies have been developed, among them those based on the global positioning system (GPS), which has allowed scientists to collect fine-scale location data for far-ranging species. This technology has made possible the study of behaviour, migration, home ranges, human-wildlife conflict and climate change (Hebblewhite & Haydon 2010), but also the definition of foraging areas (Dragon et al. 2012). Moreover, GPS has made possible to track elusive marine species, like seabirds (e.g. Frederiksen et al. 2012), cetaceans (e.g. Sveegaard et al. 2015), sea turtles (e.g. Godley et al. 2008), and even fish (e.g. Horton et al. 2020). Alone, tracking can specify the location or areas used by an organism, but in synchronization with diet studies, it can also define the distribution area of prey (e.g. Alonso et al. 2018) and describe the interannual fluctuations of prey communities (e.g. Hoskins & Arnould 2014), allowing in its turn to delimit important areas, not only for the consumer but also of its prey.

## **Marine Trophic Webs**

A food web is a representation of the flows of energy and matter created by the interaction among organisms (Cohen et al. 1993). It depicts a complex structure that changes through space and time and which influences population dynamics, and the adaptability and evolution of species (Cohen et al. 1993).

The study of food webs has had the main goal of describing the feeding relationships among species in a community. It is a complex subject that started with the study of simple linear food chains and evolved into the study of more reticulated food webs. Food chain is the representation of links between species, starting in the producer and ending at the top predator. There are two types of food chains, grazing and detritus. The first has autotrophs as the origin of energy and nutrients whereas particulate organic matter is the basal energy source in the detritus food chain. The same food web can be constituted by both types of food chains, and depending on the ecosystem, one or the other or even both, can dominate the food web (Odum 1980, Asmus & Asmus 1985, Han et al. 2017). Different types of food webs have been suggested by Robert Paine based on species of a rocky intertidal zone, the connectedness web, the energy flow web, and the functional web (Paine 1980). The connectedness web is based on observation of feeding links, while the energy flow web, as its name suggests, measures the rate at which energy flows through nodes in the web. Lastly, the functional web reflects only the influence of consumers on the consumed species (Paine 1980). Yet, such webs were still too simple or inaccurate to describe the complexity of the natural food webs, which led to the development of present food web models. The connectedness web was brought together with two fundamental species traits, the biomass and abundance of both consumers and sources (Cohen et al. 2003), allowing for the exploration of more complex interactions between trophic structures.

Food webs represent the union points between all the species of an ecosystem. There are variables and processes that might affect a specific group species, but if the food web remains stable, the ecosystem is more likely to remain in equilibrium. Stable food webs are typically characterised by few strong interactions integrated among a group of weak links (McCann et al. 1998, Neutel et al. 2002). Weak links prevent complex food webs with long loops from being unstable (Neutel et al. 2002) or having chaotic dynamics by dampening the potential destabilizing effects of strong consumer-resource interactions (McCann et al. 1998). A higher number of weak links lowers the mean and variance of interaction strengths and increases local stability in communities (Kokkoris et al. 1999). For these reasons, the disappearance of weak links can be damaging for natural communities by decreasing population abundances, ultimately leading to their extinction (McCann et al. 1998). The existence of smaller food chains with fewer trophic levels and a more web-like look, in opposition to long and unstable food chains with a reduced number of interactions between species with higher strengths, has proven to provide more

stability to food webs (Borrelli & Ginzburg 2014). Therefore, it is expectable that generalist-dominated food webs are prone to exhibiting less fluctuations than specialist-dominated food webs, since generalist species create more links with a lower average interaction strength than specialists (McCann et al. 1998). In the same line, a less productive system will house less species and will tend to have fewer links with larger average interaction strengths. More productive systems house higher biodiversity, which has been linked to stability and a higher resistance and resilience of an ecosystem to perturbations (Kondoh 2003, Ives & Carpenter 2007).

There are different perturbations influencing the oceanic environment (Lu et al. 2018, O’Leary et al. 2020), some of natural origin and some triggered by human activities, but most eventually end up being reflected in the ecology and behavior of its species and, consequently, in the functioning of the food web and the ecosystem (Lavoie et al. 2010, Christianen et al. 2017, Michel et al. 2019). Yet, knowledge on the impacts and implications of global changes and other human activities in marine ecosystem are scarce. For that reason, it is important to study the food web as a whole, so we can better understand the ecosystem and implement holistic management and conservation measures, instead of less effective single-species measures.

Food web models can provide an overview of the food web and of the ecosystem. Among the several existing software and packages is the Ecopath with Ecosim (Christensen & Walters 2004), a mass-balanced trophic model that allows to address ecological questions, like the evaluation of the impacts of fishing activities in ecosystems (e.g. Colléter et al. 2015), placement and effects of marine protected areas, and effects of environmental changes (e.g. Heymans et al. 2014). It further allows to study management policy options (Christensen et al. 2008). However, in order to build the model, it is necessary to gather information on the species of the ecosystem, including knowledge on their biomass and production, and diet composition of each of the functional groups/species. Thus, diet studies are important in an ecosystem perspective and not only for knowledge on the ecology of single species.

## **The oceanic region of the archipelago of Madeira**

The archipelago of Madeira is located off Northwest Africa, in the subtropical front where the cold waters from the north and the warm waters from the south meet. It is composed by three groups of oceanic islands, Madeira, Desertas and Porto Santo, which are surrounded by the Canary and the Azores currents. As most oceanic islands, they have



narrow continental shelves, prolonged by vast abyssal plains that can go as deep as ca. 5,000 m. Three hundred kilometers south of Madeira is the archipelago of Selvagem, and part of the Exclusive Economic Zone (EEZ) of Madeira. The Madeira region is relatively poor when compared to the upwelling systems found in African neritic areas (Cropper et al. 2014). Even so, small and localized upwelling phenomena occur near the coast of the island and in between Madeira and Desertas islands. An underwater ridge resulting from the collapsed volcanic crater is probably causing this upwelling and manifestation of cold and highly productive water at the surface (Caldeira et al. 2002). Furthermore, cold core eddies located in the western side of Madeira and Desertas also showed to increase greatly the productivity in this area (Caldeira et al. 2002). Due to these processes, it is possible for the region to harbour a relevant variety of taxa.

The archipelago of Madeira is an important feeding ground for many species during the warmer months (spring to autumn). Many go there to breed, like seabirds, arriving in the beginning of the spring and leaving only when autumn has started. The Cory's shearwater *Calonectris borealis* is the most abundant seabird species breeding in these islands (Granadeiro et al. 2006), although other common species, like the Bulwer's petrel *Bulweria bulwerii* and the Band-rumped storm-petrel *Hydrobates castro*, also breed in Madeira (Meirinho et al. 2014). Underwater, tunas are among the most iconic figures of "summer" in Madeira, as they are considered a delicacy and, therefore, are highly targeted by fisheries (Gouveia et al. 2019). The Bigeye tuna *Thunnus obesus* and the Skipjack tuna *Katsuwonus pelamis* are the most abundant, but the Albacore tuna *Thunnus alalunga* has known to exceed the abundance of Skipjack tuna in some years (Gouveia et al. 2019). The Atlantic bluefin tuna *Thunnus thynnus* and the Yellowfin tuna *Thunnus albacares* can also be found in these waters (Gouveia et al. 2019). Several species of cetaceans, like Fin whales *Balaenoptera physalus*, also visit these waters during their migrations (Freitas et al. 2012). Many others are resident, like the short-finned pilot whale *Globicephala macrorhynchus*, or are sighted all-year-around, like the Sperm whale *Physeter macrocephalus* (Freitas et al. 2012, 2014). Madeira is also known as a good sport-fishing spot, which reflects the abundance of large predatory pelagic fish, like the Atlantic blue marlin *Makaira nigricans*. Going lower in the trophic web, we will find mackerels (Blue jack *Trachurus picturatus* and Atlantic chub mackerels *Scomber colias*) dominating the intermediate trophic levels (Hermida & Delgado 2016). Other small pelagic fishes are present but in lower abundances, like European pilchard *Sardina pilchardus* and Bogue *Boops boops*. Another iconic species of Madeira is the Scabbard

fish *Aphanopus carbo*, the most captured species by the Madeiran fishing fleet (DREM data), and another touristic attraction of the region.

There are presently unmeasured threats that are impacting oceanic ecosystems, such as human exploitation and climate change, and the assessment of their real impact in these ecosystems directly depends on the existence of good quality baseline information on the ecology of species and interactions among food web components. The oceanic ecosystem around the archipelago of Madeira, despite the recognised importance of its biodiversity and services provided to humans, is still largely understudied and several of the most important components of the food web and their interactions are still unknown.

## **Thesis outline**

The general objective of this work was to assess the trophic web of the oceanic region of the archipelago of Madeira. To tackle this, I first described the diet of key element species from this ecosystem, namely of small pelagic fishes, tunas, and seabirds, of which very few information was available. I also characterized the community of the most common and available epipelagic species of this ecosystem and produced a model which could explain the food web system and interactions of this oceanic region.

This thesis comprises a total of seven chapters: a first chapter which introduces the topics discussed in the following chapters: five chapters (Chapter 2-6) corresponding to a series of five articles (one published, one accepted, two submitted and in revision, and one manuscript in progress); and a last chapter where I provide a general discussion about the chapters presented and the main conclusions of the thesis.

### **Chapter 2 - Diet and trophic position of two mackerel species in the archipelago of Madeira, Portugal**

In this chapter we describe for the first time the diet of the Atlantic chub mackerel *Scomber colias* and Blue jack mackerel *Trachurus picturatus* in the oceanic region of the archipelago of Madeira. These are two of the most abundant and important species in this ecosystem, as prey for many top predators that use this area as a migratory corridor and as feeding grounds. Despite several studies performed in neritic areas, oceanic studies are scarce and information on the feeding ecology of these mackerels is limited in these regions.

Chapter 3 - The diet of the Bigeye tuna *Thunnus obesus* and the Skipjack tuna *Katsuwonus pelamis* in the subtropical pelagic region of the northeast Atlantic

In this chapter we describe for the first time the diet of the Bigeye tuna *Thunnus obesus* and Skipjack tuna *Katsuwonus pelamis* in the archipelago of Madeira. This is the first diet study for the Bigeye tuna in the Northeast Atlantic and only the second for the Skipjack tuna, the first being in the Canary Islands.

Chapter 4 - A gull that scarcely ventures on the ocean: Yellow-legged gulls *Larus michahellis atlantis* on the oceanic island of Madeira

In this chapter we describe the distribution and foraging behaviour of the Yellow-legged gull *Larus michahellis atlantis* in the archipelago of Madeira, using GPS-GSM devices and stable isotope analysis.

Chapter 5 - Seabird diet analysis suggests sudden shift in the pelagic communities of the subtropical Northeast Atlantic

In this chapter we studied the variation in the composition of epipelagic species through time in the broad Canary Current region, by analysing foraging trips and regurgitations of Cory's shearwaters *Calonectris borealis* nesting on Selvagem Grande island in 2008-2011 and 2016-2018. Furthermore, we used fisheries, oceanographic data, and the North Atlantic Oscillation as possible explanatory variables for trends in behaviour and diet of Cory's shearwaters.

Chapter 6 – Food web structure of the subtropical oceanic archipelago of Madeira, eastern North Atlantic

In this chapter we aim to develop an ecosystem-based model of the Exclusive Economic Zone of the archipelago of Madeira, using the Ecopath with Ecosim software. As little-known ecosystems, oceanic environments, like the archipelago of Madeira, are still in need of base line information before we can understand the extent of global changes impacts. This chapter is still a work in progress.

## References

Alonso H, Granadeiro JP, Dias MP, Catry T, Catry P (2018) Fine-scale tracking and diet information of a marine predator reveals the origin and contrasting spatial distribution of prey. Prog Oceanogr 162:1–12

- Alonso H, Granadeiro JP, Waap S, Xavier J, Symondson WOC, Ramos JA, Catry P (2014) An holistic ecological analysis of the diet of Cory's shearwaters using prey morphological characters and DNA barcoding. *Mol Ecol* 23:3719–3733
- Asmus H, Asmus R (1985) The importance of grazing food chain for energy flow and production in three intertidal sand bottom communities of the northern Wadden Sea. *Helgoländer Meeresuntersuchungen* 39:273–301
- Barrett RT, Camphuysen K, Anker-Nilssen T, Chardine JW, Furness RW, Garthe S, Hüppop O, Leopold MF, Montevecchi WA, Veit RR (2007) Diet studies of seabirds: A review and recommendations. *ICES J Mar Sci* 64:1675–1691
- Beneditto APM Di, Siciliano S (2007) Stomach contents of the marine tucuxi dolphin (*Sotalia guianensis*) from Rio de Janeiro, south-eastern Brazil. *J Mar Biol Assoc United Kingdom* 87:253–254
- Beneditto APM Di, Souza CMM de, Kehrig HA, Rezende CE (2011) Use of multiple tools to assess the feeding preference of coastal dolphins. *Mar Biol* 158:2209–2217
- Borrelli JJ, Ginzburg LR (2014) Why there are so few trophic levels: Selection against instability explains the pattern. *Food Webs* 1:10–17
- Caldeira RMA, Groom S, Miller P, Pilgrim D, Nezlin NP (2002) Sea-surface signatures of the island mass effect phenomena around Madeira Island, Northeast Atlantic. *Remote Sens Environ* 80:336–360
- Cherel Y, Xavier JC, Grissac S De, Trouvé C, Weimerskirch H (2017) Feeding ecology, isotopic niche, and ingestion of fishery-related items of the wandering albatross *Diomedea exulans* at Kerguelen and Crozet Islands. *Mar Ecol Prog Ser* 565:197–215
- Choy CA, Popp BN, Kaneko JJ, Drazen JC (2009) The influence of depth on mercury levels in pelagic fishes and their prey. *Proc Natl Acad Sci* 106:13865–13869
- Christensen V, Walters CJ (2004) Ecopath with Ecosim: Methods, capabilities and limitations. *Ecol Modell* 172:109–139
- Christensen V, Walters C, Pauly D, Forrest R (2008) Ecopath with Ecosim version 6: User Guide.
- Christianen MJA, Heide T Van Der, Holthuijsen SJ, Reijden KJ Van Der, Borst ACW, Olff H (2017) Biodiversity and food web indicators of community recovery in intertidal shellfish reefs. *Biol Conserv* 213:317–324
- Cohen AJE, Beaver R a, Cousins SH, Deangelis DL, Goldwasser L, Heong KL, Holt RD, Kohn AJ, Lawton JH, Martinez N, O'Malley R, Page LM, Patten BC, Pimm SL,

- Polis GA, Rejmánek M, Schoener TW, Schoenly K, Sprules WG, Teal JM, Ulanowicz RE, Warren PH, Wilbur HM, Yodzis P (1993) Improving Food Webs. *Ecology* 74:252–258
- Cohen JE, Jonsson T, Carpenter SR (2003) Ecological community description using the food web, species abundance, and body size. *Proc Natl Acad Sci U S A* 100:1781–1786
- Colléter M, Valls A, Guitton J, Gascuel D, Pauly D, Christensen V (2015) Global overview of the applications of the Ecopath with Ecosim modeling approach using the EcoBase models repository. *Ecol Modell* 302:42–53
- Correia E, Granadeiro JP, Mata VA, Regalla A, Catry P (2019) Trophic interactions between migratory seabirds, predatory fishes and small pelagics in coastal West Africa. *Mar Ecol Prog Ser* 622:177–189
- Cropper TE, Hanna E, Bigg GR (2014) Spatial and temporal seasonal trends in coastal upwelling off Northwest Africa, 1981-2012. *Deep Res Part I Oceanogr Res Pap* 86:94–111
- Denda A, Stefanowitsch B, Christiansen B (2017) From the epipelagic zone to the abyss: Trophic structure at two seamounts in the subtropical and tropical Eastern Atlantic - Part I zooplankton and micronekton. *Deep Res Part I Oceanogr Res Pap* 130:63–77
- Doidge DW, Croxall JP (1985) Diet and energy budget of the Antarctic fur seal, *Arctocephalus gazella* at South Georgia. *Antarct Nutr cycles food webs*:543–550
- Dragon AC, Bar-Hen A, Monestiez P, Guinet C (2012) Comparative analysis of methods for inferring successful foraging areas from Argos and GPS tracking data. *Mar Ecol Prog Ser* 452:253–267
- Ford MJ, Hempelmann J, Hanson MB, Ayres KL, Baird RW, Emmons CK, Lundin JI, Schorr GS, Wasser SK, Park LK (2016) Estimation of a killer whale (*Orcinus orca*) population's diet using sequencing analysis of DNA from feces. *PLoS One* 11:1–14
- Frederiksen M, Moe B, Daunt F, Phillips RA, Barrett RT, Bogdanova MI, Boulinier T, Chardine JW, Chastel O, Chivers LS, Christensen-Dalsgaard S, Clément-Chastel C, Colhoun K, Freeman R, Gaston AJ, González-Solís J, Goutte A, Grémillet D, Guilford T, Jensen GH, Krasnov Y, Lorentsen SH, Mallory ML, Newell M, Olsen B, Shaw D, Steen H, Strøm H, Systad GH, Thórarinnsson TL, Anker-Nilssen T

- (2012) Multicolony tracking reveals the winter distribution of a pelagic seabird on an ocean basin scale. *Divers Distrib* 18:530–542
- Freitas L, Alves F, Ribeiro C, Dinis A, Nicolau C, Carvalho A (2014) Estudo técnico-científico de suporte à proposta de criação de áreas de operação para a actividade de whale watching e a respectiva capacidade de carga (Deliverable A.7\_IIA).
- Freitas L, Dinis A, Nicolau C, Ribeiro C, Alves F (2012) New records of cetacean species for Madeira archipelago with an updated checklist. *Bol do Mus Munic do Funchal (História Nat)* 62:25–43
- Godley BJ, Blumenthal JM, Broderick AC, Coyne MS, Godfrey MH, Hawkes LA, Witt MJ (2008) Satellite tracking of sea turtles: Where have we been and where do we go next? *Endanger Species Res* 4:3–22
- Gouveia L, Amorim A, Alves A, Hermida M (2019) Updated fishery statistics of tuna species caught off Madeira archipelago. *Collect Vol Sci Pap ICCAT* 75:2081–2097
- Granadeiro JP, Dias MP, Rebelo R, Santos CD, Catry P (2006) Numbers and Population Trends of Cory's Shearwater *Calonectris diomedea* at Selvagem Grande, Northeast Atlantic. *Waterbirds* 29:56–60
- Granadeiro JP, Nunes M, Silva M, Furness RW (1998) Flexible foraging strategy of Cory's shearwater, *Calonectris diomedea*, during the chick-rearing period. *Anim Behav* 56:1169–1176
- Granadeiro JP, Silva MA (2000) The use of otoliths and vertebrae in the identification and size-estimation of fish in predator-prey studies. *Cybium* 24:383–393
- Han D, Xue Y, Zhang C, Ren Y (2017) A mass balanced model of trophic structure and energy flows of a semi-closed marine ecosystem. *Acta Oceanol Sin* 36:60–69
- Hebblewhite M, Haydon DT (2010) Distinguishing technology from biology: A critical review of the use of GPS telemetry data in ecology. *Philos Trans R Soc B Biol Sci* 365:2303–2312
- Hermida M, Delgado J (2016) High trophic level and low diversity: Would Madeira benefit from fishing down? *Mar Policy* 73:130–137
- Heymans JJ, Coll M, Libralato S, Morissette L, Christensen V (2014) Global patterns in ecological indicators of marine food webs: A modelling approach. *PLoS One* 9
- Horton TW, Block BA, Drumm A, Hawkes LA, O'Cuaig M, Maoiléidigh NÓ, O'Neill R, Schallert RJ, Stokesbury MJW, Witt MJ (2020) Tracking Atlantic bluefin tuna from foraging grounds off the west coast of Ireland. *ICES J Mar Sci*

- Hoskins AJ, Arnould JPY (2014) Relationship between long-term environmental fluctuations and diving effort of female Australian fur seals. *511*:285–295
- Ives AR, Carpenter SR (2007) Stability and diversity of ecosystems. *Science* (80- ) *317*:58–62
- Karnovsky NJ, Hobson KA, Iverson SJ (2012) From lavage to lipids: Estimating diets of seabirds. *Mar Ecol Prog Ser* *451*:263–284
- Kokkoris GD, Troumbis AY, Lawton JH (1999) Patterns of species interaction strength in assembled theoretical competition communities. *Ecol Lett* *2*:70–74
- Kondoh M (2003) Foraging adaptation and the relationship between food-web complexity and stability. *Science* (80- ) *299*:1388–1391
- Lavoie RA, Hebert CE, Rail JF, Braune BM, Yumvihoze E, Hill LG, Lean DRS (2010) Trophic structure and mercury distribution in a Gulf of St. Lawrence (Canada) food web using stable isotope analysis. *Sci Total Environ* *408*:5529–5539
- Layman CA, Araujo MS, Boucek R, Hammerschlag-Peyer CM, Harrison E, Jud ZR, Matich P, Rosenblatt AE, Vaudo JJ, Yeager LA, Post DM, Bearhop S (2012) Applying stable isotopes to examine food-web structure: An overview of analytical tools. *Biol Rev* *87*:545–562
- Lu Y, Yuan J, Lu X, Su C, Zhang Y, Wang C, Cao X, Li Q, Su J, Ittekkot V, Garbutt RA, Bush S, Fletcher S, Wagey T, Kachur A, Sweijid N (2018) Major threats of pollution and climate change to global coastal ecosystems and enhanced management for sustainability. *Environ Pollut* *239*:670–680
- McCann K, Hastings A, Huxel GR (1998) Weak trophic interactions and the balance of nature. *Nature* *395*:794–798
- Meirinho A, Barros N, Oliveira N, Catry P, Lecoq M, Paiva V, Geraldés P, Granadeiro JP, Ramírez I, Andrade J (2014) Atlas das aves marinhas de Portugal. Sociedade Portuguesa para o Estudo das Aves. Lisboa
- Michel LN, Danis B, Dubois P, Eleaume M, Fournier J, Gallut C, Jane P, Lepoint G (2019) Increased sea ice cover alters food web structure in East Antarctica. *Sci Rep* *9*:1–11
- Miller AK, Sydeman WJ (2004) Rockfish response to low-frequency ocean climate change as revealed by the diet of a marine bird over multiple time scales. *Mar Ecol Prog Ser* *281*:207–216
- Morera-Pujol V, Ramos R, Pérez-Méndez N, Cerdà-Cuellar M (2018) Multi-isotopic assessments of spatio-temporal variability of diet: the case of two sympatric gulls


in the Western Mediterranean Institut de Recerca de la Biodiversitat (IRBio) and Departament de Biologia Evolutiva, Ecologia i Ciències Ambient. Mar Ecol Prog Ser 606:4–7

- Neutel AM, Heesterbeek JAP, Ruiter PC De (2002) Stability in real food webs: Weak links in long loops. *Science* (80- ) 296:1120–1123
- O’Leary BC, Hoppit G, Townley A, Allen HL, McIntyre CJ, Roberts CM (2020) Options for managing human threats to high seas biodiversity. *Ocean Coast Manag* 187:105110
- Odum EP (1980) the Status of Three Ecosystem-Level Hypotheses Regarding Salt Marsh Estuaries: Tidal Subsidy, Outwelling, and Detritus-Based Food Chains. ACADEMIC PRESS, INC.
- Paine RT (1980) Food Webs: Linkage, Interaction Strength and Community Infrastructure. *Journal of An*
- Spitz J, Rousseau Y, Ridoux V (2006) Diet overlap between harbour porpoise and bottlenose dolphin: An argument in favour of interference competition for food? *Estuar Coast Shelf Sci* 70:259–270
- Sveegaard S, Galatius A, Dietz R, Kyhn L, Koblitz JC, Amundin M, Nabe-Nielsen J, Sinding MHS, Andersen LW, Teilmann J (2015) Defining management units for cetaceans by combining genetics, morphology, acoustics and satellite tracking. *Glob Ecol Conserv* 3:839–850
- Wallace BP, Avens L, Braun-McNeill J, McClellan CM (2009) The diet composition of immature loggerheads: Insights on trophic niche, growth rates, and fisheries interactions. *J Exp Mar Bio Ecol* 373:50–57
- Xavier JC, Cherel Y (2009) Cephalopod beak guide for the Southern Ocean.



## CHAPTER 2

---



**Credits: Joana Romero**

*This chapter is published as:*

*Joana Romero\*, Carolina Vieira\*, Susana Garrido, Margarida Hermida, Paulo Catry, Graça Faria, José Pedro Granadeiro (2021). Diet and trophic position of two mackerel species in the archipelago of Madeira, Portugal. Journal of Fish Biology, 202:1-13.*

*\*contributed equally*

# Diet and trophic position of two mackerel species in the archipelago of Madeira, Portugal

## Abstract

The Atlantic chub mackerel *Scomber colias* and the Blue jack mackerel *Trachurus picturatus* are two abundant species in the Macaronesia region which include the archipelago of Madeira, Portugal. Both are key species in the trophic web, being important prey for several local top predators, such as seabirds and marine mammals. However, little is known about their feeding ecology in oceanic environments. In this study, we describe the diets of the *S. colias* and the *T. picturatus* in the oceanic region of Madeira throughout a year and by seasons. *Scomber colias* fed on a broader range of prey groups than *T. picturatus*, but for both species, zooplankton (particularly calanoid copepods) and fish were the most important food items, according to stomach contents. The diet of *S. colias* included a higher proportion of fish, namely Atlantic saury *Scomberesox saurus* and *S. colias*, than that of *T. picturatus*, that included mostly Longspine snipefish *Macroramphosus scolopax*. *Trachurus picturatus* also consumed higher proportions of decapods and other copepods. Seasonal variation was found in the diet of both species, with zooplanktonic species being more important in colder months (February to April) for *S. colias* and during warm months (May to October) for *T. picturatus*. Their diet in other seasons were dominated by fish. Despite consuming similar prey, Carbon and Nitrogen stable isotope analysis of muscle of *S. colias* and *T. picturatus* showed little overlap in their diets, and *T. picturatus* showed higher  $\delta^{15}\text{N}$  and a narrower isotopic niche.

**Keywords:** Copepods, Diet, Mackerels, Stable isotopes, Stomach contents, Trophic position.

## Introduction

The Atlantic chub mackerel *Scomber colias* Gmelin 1789 and the Blue jack mackerel *Trachurus picturatus* Bowdich 1825 are two abundant species in the temperate and subtropical coastal waters of the Atlantic. *Scomber colias* are distributed from the Bay of Biscay to the Azores and down to South Africa (Collette, 1983; Hernández and Ortega, 2000). While *T. picturatus* share the northern limits of *S. colias*, these are restricted to Mauritania in the southern limit (Shaboneyev and Kotlyar, 1979; Shaboneyev and Ryazantseva, 1977). Both *S. colias* and *T. picturatus* occupy the surface layers of the water column but can also go as deep as 300 and 500 m, respectively (Hernández and Ortega, 2000; Menezes *et al.*, 2009). They can also be found in more offshore waters, in seamounts and close to islands, such as those of the archipelago of Madeira, Portugal (Vasconcelos *et al.*, 2012, 2018). These species occupy intermediate trophic levels and are key prey species in the pelagic trophic structure for many predators, such as cephalopods (e.g. Martins, 1982), large pelagic fish, like tunas (e.g. Logan *et al.*, 2011), seabirds (e.g. Alonso *et al.*, 2014; Romero *et al.* 2019) and cetaceans (Giménez *et al.*, 2017; Marçalo *et al.*, 2018). Both fish species have significant economic value worldwide, including in the oceanic region of Madeira, where an average of 400 and 200 tonnes (2008-2018, DRPM, Unpublished data) of *T. picturatus* and *S. colias*, respectively, are fished every year. These species also have important roles in the tuna fishing activities as bait (Hermida and Delgado, 2016; Tejerina *et al.*, 2019).

Because of the importance of both mackerel species in fisheries, several studies have focused on several aspects of their ecology, like stock assessment, life history parameters, reproduction and trace elements (Garcia *et al.*, 2015; Lozano-Bilbao *et al.*, 2019; Raimundo *et al.*, 2013; Vasconcelos *et al.*, 2006, 2011, 2012). However, there is still limited knowledge on their feeding ecology (e.g. Bachiller and Irigoien, 2015; Garrido *et al.* 2015; Wahbi *et al.*, 2015) and few studies were carried out in oceanic environments (Castro, 1993; Denda *et al.*, 2017; Hirsch and Christiansen, 2010). To date there are no detailed studies on the diet of these two species in the vast oceanic area that surrounds the archipelago of Madeira. Such information may provide an insight on the planktonic communities of this poorly known ecosystem and allows to better understand what sustains the pelagic trophic web of this region.

The main goal of this study is to describe the diet of the *S. colias* and the *T. picturatus* in the oceanic region of Madeira, and its seasonal variability throughout the

year. Based on the analysis of stomach contents and the use of stable isotope analysis of Carbon and Nitrogen, we (1) identify the most important prey for each species in overall and in each season, (2) describe their trophic position, and (3) compare the diet of the two mackerel species.

## Methods

### *Ethical Statement*

The fish analysed in this study were acquired dead by the Madeira Regional Fisheries Directorate (DRP), under the framework of the Programa Nacional de Recolha de Dados da Pesca (multiannual European Union programme for the collection, management and use of data in the fisheries and aquaculture sectors).

### *Sampling*

Between February 2017 and January 2018, *S. colias* and *T. picturatus* individuals were obtained from commercial landings of local purse-seine fishing vessels (mostly using an 18mm mesh size), operating south of Madeira Island. A total of 20 individuals of each species were collected every month, except in July, August and December (for both species), and November (for the *T. picturatus*) (see Supporting Information, Table S1). For each individual, we recorded the total length (see Supporting Information, Table S1), total mass, mass of the stomach and of its content, and also of the gonads (IPIMAR, 2009).

### *Stomach contents analysis*

To avoid variability in dietary data due to ontogenetic differences, we selected individuals of each species belonging to the modal size classes of 20-24 cm and 19-22 cm for *S. colias* and *T. picturatus*, respectively, for each month. In June (for *S. colias*) and September (for *T. picturatus*) there were not enough individuals within the selected size range. Therefore, we sampled nine *S. colias* of 22-27 cm, and four *T. picturatus* of 15-20 cm.

Sampling followed the procedures described in Garrido *et al.* (2015). We carefully collected the entire content of the cardiac and pyloric regions, to avoid food items in more advanced stages of digestion in the intestine. For both species, groups of stomach contents from 3 individuals collected in the same haul were combined to form a single sample, and an aliquot was used to identify and count prey items. Stomach samples were filtered using a 200  $\mu\text{m}$  sieve (see Supporting Information, Table S2,). Fractions  $<200 \mu\text{m}$  were homogenized, placed in a Utermohol chamber and identified using an inverted microscope

(Leica DM IL LED) at a magnification of 200x. Fractions >200 µm were homogenized in a known volume of water and analysed under a stereomicroscope at a magnification of 160x (Olympus SZX12). Since *T. picturatus* is not a consumer of phytoplankton (e.g. Hirsch and Christiansen, 2010; Kompowski, 1976), this group was only quantified in the stomachs of *S. colias*. All food items were identified to the lowest possible taxonomic level. Fish scales found in the stomachs of both mackerel species were disregarded, as they were most probably swallowed during the fishing event (Hernández and Ortega, 2000; Hirsch and Christiansen, 2010; Kompowski, 1976).

#### *Diet characterization*

To assess the percentage of empty stomachs of each species, we calculated the vacuity index as the number of empty samples in relation to the total number of samples of each species. The feeding intensity (FI%) of both mackerel species, used to assess the stomach fullness, was calculated as:

$$FI\% = \frac{\text{stomach content mass}}{\text{total mass} - \text{gonads mass}} \times 100$$

To describe the diet of the *S. colias* and the *T. picturatus*, we calculated the frequency of occurrence (FO%), as the number of samples in which a prey group occurred in relation to the total number of samples. To account for the importance of prey for dietary energy we estimated the contribution of prey to dietary carbon by estimating the percentage of carbon content (CC%) as the mass of carbon of each prey group in the sample in relation to total dietary carbon in that sample. The carbon content of each prey was calculated using published relationships of volume and weight with their carbon content of phytoplankton, zooplankton and ichthyoplankton organisms (see Supporting Information, Table S3). To assess the importance of each prey in the diet of the two mackerels, the modified index of relative importance (mIRI) was calculated according to the equation (Hayse, 1990):

$$mIRI = FO\% \text{ of group prey} \times CC\% \text{ of group prey}$$

The diversity of the diet of the two mackerel species was compared using the Shannon-Wiener index, based on the mean percentage of carbon content of each prey (lowest taxonomic level identified). To avoid bias due to the different number of samples collected for each species (n=22 and n=27, for *T. picturatus* and *S. colias*, respectively), and to estimate the variability within each species, we calculated the means and standard deviations of 1000 random samples (with repetition) of n=22 from each species. Dietary

overlap between *S. colias* and *T. picturatus* was calculated using the Schoener's index, based on the carbon contribution of each prey (S):

$$S = 1 - 0.5 \left( \sum_{i=1}^n |C_{ij} - C_{ik}| \right)$$

$C_{ij}$  and  $C_{ik}$  are the proportion of carbon content of prey  $i$  (lowest taxonomic level identified) in the diet of species  $j$  and species  $k$ . The index varies between zero (diets are completely different) and one (identical proportions of all prey) and values  $>0.6$  are considered to represent a significant diet overlap (Wallace and Ramsey, 1983).

The temporal variability of diet content was assessed by dividing the year in four seasons: Season 1 (February-April), Season 2 (May-July), Season 3 (August-October) and Season 4 (November-January). For each season, the relative importance of each prey based on the carbon content contribution was calculated, such as the frequency of occurrence and mIRI. Differences in prey groups (%CC) among seasons were described using multivariate ordination (Principal Component Analysis, PCA).

#### *Stable isotope analysis*

White muscle of *S. colias* (n=12) and *T. picturatus* (n=12) individuals was collected for nitrogen and carbon stable isotope analysis. Before analysis, samples were lipid-extracted using a chloroform-methanol (2:1) solution. The analysis was performed using a continuous-flow isotope-ratio mass spectrometer (CF-IRMS) on a Sercon Hydra 20-22 (Sercon, UK) coupled to a EuronEA (EuroVector, Italy) elemental analyser, which automatizes the samples preparation (Preston and Owens, 1983), in the Stable Isotopes and Instrumental Analysis Facility of the University of Lisbon. The results are expressed relative to atmospheric nitrogen for  $\delta^{15}\text{N}$ , and the Vienna Pee Dee Belemnite for  $\delta^{13}\text{C}$ . The precision of the analysis was calculated using values from six to nine replicates of laboratory standard material (casein),  $\leq 0.08\text{‰}$  for  $\delta^{15}\text{N}$  and  $\leq 0.06\text{‰}$  for  $\delta^{13}\text{C}$ .

To compare the isotopic niches of *S. colias* and *T. picturatus*, we estimated the corrected standard ellipse area (SEAc) of each species corresponding to 95% of the data and the overlap of these ellipses using the function *maxLikOverlap* of the *SIBER* R package (Jackson *et al.*, 2011). The proportion of the overlapping area was calculated as the ratio of the overlap area with the total area of the two polygons minus the overlap area (or with the area of the polygon of each mackerel). The trophic position (TP) of the consumers was estimated from the isotopic ratios according to Vander Zanden and Rasmussen (2001):

$$TP = TP_b + (\delta^{15}N_c - \delta^{15}N_{base}) / \Delta N$$

where  $TP_b$  is the baseline of the trophic web (in this case, Particulate Organic Matter (POM)=1.5),  $\delta^{15}N_c$  is the  $\delta^{15}N$  of the consumer, and  $\delta^{15}N_{base}$  is the mean  $\delta^{15}N$  of POM.  $\Delta N$  is the enrichment in  $^{15}N$  per trophic level and was assumed to be 3.4% (Post, 2002). All analyses were performed using R version 3.5.2 (R Core Team, 2018).

## Results

### Overall diet

A total of 27 samples, pooled from 81 stomachs of *S. colias* were analysed, none of which was empty. From these, 66 prey types, belonging to 41 orders, were identified (Table 1). For *T. picturatus*, 22 samples pooled from 66 stomachs were analysed, with a vacuity index of 4.6%. Twenty-eight prey types, belonging to 15 orders, were described for this species (Table 1). In general, *S. colias* presented a more diverse diet (Shannon-Wiener index: *S. colias* =  $2.6 \pm 0.2$  (SE) and *T. picturatus* =  $2.3 \pm 0.1$ ) and higher feeding intensities (FI%=1.1, range=[0.6-2.0]) than the *T. picturatus* (FI%=0.3 [0.2-0.4]).

In this study, *S. colias* fed on phytoplankton, zooplankton, fish, cephalopods, and tunicates, and *T. picturatus* fed on zooplankton and fish. Overall, fish represented the highest mean contribution to *S. colias* dietary carbon ( $51.5 \pm 48.2\%$  (SD), n=22), slightly higher than zooplankton ( $36.3 \pm 45.6\%$ ), while the pattern was inversed in terms of frequency of occurrence (100% for zooplankton vs. 66% for fish). Contrarily, for *T. picturatus*, zooplankton presented the highest mean contribution to dietary carbon ( $63.6 \pm 49.2\%$ , n=27) and was present in more stomachs (FO%=95%), followed by fish (%CC= $31.8 \pm 47.7\%$  and FO%=31%). The mIRI was higher for zooplankton than for fish in the diet of both species due to the lower frequency of occurrence of fish in the stomachs (Table 1). *Scomber colias* occasionally included small cephalopods in their diet (FO% ca. 15%), a food item which was not detected in the diet of *T. picturatus*.

Both mackerel species preyed mainly on copepods, which was the most important group among planktonic crustaceans, both in frequency of occurrence and carbon content (Table 1). Among these, calanoids (*Candacia* sp. and *Pleuromamma* sp.) and unidentified copepods were the dominant prey and the ones with the highest mIRI (Table 1). Cyclopoid copepods (*Oncaea* sp.) were also consumed, occurring more frequently in the *S. colias* diet than in *T. picturatus*. Crustacean decapods and pteropods (Gastropoda) were also important in the diet of *T. picturatus*. Eggs of crustaceans, fish and gastropods were

frequently found in the stomachs of *S. colias* and represented on average  $1.2 \pm 0.8\%$  of contribution to dietary carbon.

Despite being present in all *S. colias* stomachs analysed, the carbon contribution of phytoplankton ( $1.4 \pm 4.5\%$ ) was very low, mostly comprising dinoflagellates (*Dinophyceae*) and diatoms (*Pennales*, *Achnanthes*).

Atlantic saury *Scomberesox saurus* Walbaum 1792 and Longspine snipefish *Macroramphosus scolopax* Linnaeus 1758 were the most important fish prey identified in the diet of *S. colias* and *T. picturatus*, respectively. Cannibalism was observed in 11% of the samples of *S. colias* analysed and represented the second most important fish contribution to its diet.

#### Seasonal variation

In Season 1, *S. colias* consumed the highest variety of prey, including unidentified fish and crustaceans, among them decapods and copepods (calanoids: *Candacia* sp. and *Euchaeta* sp., and cyclopoids: *Oncaea* sp. and *Corycaeus* sp.) (Table 1). In Season 2 Clupeidae and unidentified fish were the most important for the diet of *S. colias* as well as copepods (*Candacia* sp., *Corycaeus* sp., *Scolecithrix* sp. and *Sapphirina* sp.). In Season 3, fish dominated the diet of *S. colias* (*M. scolopax*, *S. colias*, *S. saurus* and other unidentified species), as in Season 4 (unidentified, European pilchard *Sardina pilchardus* Walbaum 1792 and *S. colias*), in which copepods were also important (*Candacia* sp., *Corycaeus* sp. and *Pleuromamma* sp.). The carbon content of cephalopods in the stomachs of *S. colias* were highest in Season 4 (Table 1).

For *T. picturatus*, Season 1 was characterized by higher consumption of myctophids and other unidentified fish, crustaceans (among them calanoids and other copepods, Brachyura and Anomura), and pteropods (Table 1). In Season 2, unidentified fish, pteropods, copepods, among them *Candacia* sp., and other crustaceans were the most consumed. Season 3 was dominated by decapods, including Anomura, fish, namely *M. scolopax*, and copepods, namely *Pleuromamma* sp.. In season 4, the most important prey of *T. picturatus* were crustaceans, with copepods like *Candacia* sp. and *Pleuromamma* sp. representing the bulk of its diet.

The multivariate ordination (PCA) of prey groups showed higher differentiation in the prey consumed between each season than between each mackerel species (Figure 1). For *S. colias*, Seasons 3 and 1 were the most different amongst each other and Season 2 and 4 were the most similar, being these last two differentiated from the other seasons



by a higher consumption of crustacean eggs, cephalopods and fish. Seasons 1 and 3 are more segregated due to the consumption of a more varied set of prey in the former, and fish in the latter. For *T. picturatus*, Seasons 3 and 4 were the most different, while Season 1 and 2 the most similar. Seasons 1 and 2 are different from other seasons due to a higher consumption of decapods and gastropods. Season 3 was characterized by a higher consumption of decapods and fish, while Season 4 was distinct due to *T. picturatus* consuming more copepods and other crustaceans.




Figure 1 – Principal component analysis (PCA) of the percentage of carbon contribution of prey and their predators, Atlantic chub mackerel *Scomber colias* (ChubMack) and Blue jack mackerel *Trachurus picturatus* (BJackMac), per season (S1: February-April, S2: May-July, S3: August-October, S4: November-January). Prey were grouped in Epipelagic fish (Epipel Fish), Unidentified fish (N. ID. Fish), Crustacean eggs (Crustac Eggs), Cephalopods, Other Prey, Phytoplankton, Fish Eggs, Calanoids, Other Copepods, Other Crustaceans (Other Crustac), Decapods, Gastropods, Mesopelagic Fish (Mesopelagic Fish).

### *Trophic levels and Isotopic niches*

There was a significant difference between the  $\delta^{15}\text{N}$  isotopic ratios of both mackerel species,  $9.6 \pm 0.2$  (SD)‰ and  $8.6 \pm 0.7$ ‰ for *T. picturatus* and *S. colias*, respectively (ANOVA:  $F_{1,22}=24.3$ ,  $P<0.001$ ). Conversely,  $\delta^{13}\text{C}$  ratios were not significantly different,  $-19.9 \pm 0.4$ ‰ and  $-20.0 \pm 0.5$ ‰ for *T. picturatus* and *S. colias*, respectively ( $F_{1,22}=0.364$ ,  $P=0.55$ ). The estimated trophic levels were significantly different ( $F_{1,22}=24.3$ ,  $P<0.001$ ) for the two mackerel species,  $3.09 \pm 0.05$  and  $2.78 \pm 0.21$  for *T. picturatus* and *S. colias*, respectively. The isotopic niche of *T. picturatus* was smaller than that of *S. colias* (*T. picturatus* isotopic area: 1.4 and *S. colias* isotopic area: 5.7;  $P=0.99$ ; Figure 2). The total dietary overlap in the isotopic space was 16%, which agrees with the value of Schoener's index, estimated at 0.51. The dietary overlap for the *S. colias* corresponded to 17% and for the *T. picturatus* 70% (Figure 2).




Figure 2 – Standard ellipse areas (SEAc, 95%) of individual Atlantic chub mackerel *Scomber colias* (in grey) and Blue jack mackerel *Trachurus picturatus* (in black) sampled in the archipelago of Madeira between February 2017 and January 2018.

Table 1 - Diet composition of Atlantic chub mackerel *Scomber colias* and Blue jack mackerel *Trachurus picturatus* captured in the archipelago of Madeira, per season (Season 1: February-April; Season 2: May-July; Season 3: August-October; Season 4: November-January) and annual.

Prey	<i>Scomber colias</i>										<i>Trachurus picturatus</i>									
	Season 1		Season 2		Season 3		Season 4		Annual		Season 1		Season 2		Season 3		Season 4		Annual	
	%CC†	mIRI‡	%CC	mIRI	%CC	mIRI	%CC	mIRI	%CC	mIRI	%CC	mIRI	%CC	mIRI	%CC	mIRI	%CC	mIRI	%CC	mIRI
<b>Phytoplankton</b>	<b>3.8</b>	<b>382.7</b>	<b>0.3</b>	<b>29.7</b>	<b>&lt;0.1</b>	<b>0.4</b>	<b>0.2</b>	<b>15.9</b>	<b>1.4</b>	<b>137.8</b>										
<b>Diatoms</b>	<b>0.3</b>	<b>27.2</b>	<b>&lt;0.1</b>	<b>0.2</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>0.7</b>	<b>&lt;0.1</b>	<b>9.3</b>										
Bacillariophyceae	0.2	18.0	<0.1	0.2	<0.1	<0.1	<0.1	0.2	<0.1	5.4										
Achnanthales																				
<i>Achnanthes</i> sp.	<0.1	1.6							<0.1	0.2										
<i>Cocconeis</i> sp.							<0.1	<0.1	<0.1	<0.1										
Pennales	0.1	7.4	<0.1	<0.1	<0.1	<0.1	<0.1	0.4	<0.1	2.5										
<b>Chlorophyta</b>																				
Pyramimonadales																				
<i>Pyramimonas</i> sp.	<0.1	3.7	<0.1	0.3					<0.1	0.8										
<b>Coccolithophores</b>	<b>0.2</b>	<b>13.5</b>			<b>&lt;0.1</b>	<b>0.1</b>	<b>&lt;0.1</b>	<b>1.1</b>	<b>&lt;0.1</b>	<b>3.8</b>										
<b>Dinoflagellates</b>	<b>2.7</b>	<b>274.9</b>	<b>0.3</b>	<b>28.5</b>	<b>&lt;0.1</b>	<b>0.4</b>	<b>0.1</b>	<b>13.9</b>	<b>1.0</b>	<b>101.1</b>										
Dinophyceae																				
Dinophysiales																				
<i>Dinophysis</i> sp.			<0.1	0.3					<0.1	<0.1										
Gonyaulacales																				
<i>Gonyaulax</i> sp.					<0.1	<0.1			<0.1	<0.1										
<i>Lingulodinium polyedra</i>	<0.1	4.1							<0.1	0.5										
Gymnodiniales																				
<i>Gymnodinium</i> sp.			<0.1	0.1					<0.1	<0.1										
Peridinales																				
<i>Protoperidinium</i> sp.			<0.1	1.4					<0.1	<0.1										
<i>Scrippsiella</i> sp.	<0.1	3.6							<0.1	0.4										
Prorocentrales																				
<i>Prorocentrum</i> sp.									<0.1	<0.1										
Cyst dinoflagellates	0.5	48.5	<0.1	2.6	<0.1	<0.1			0.2	12.4										
Unidentified	2.2	217.7	0.2	24.0	<0.1	0.3	0.1	13.9	0.8	81.1										

Unidentified phytoplankton	0.6	61.3	<0.1	0.7					0.2	11.4										
Zooplankton	73.4	7341.5	19.7	1966.9	0.3	31.2	33.3	3330.6	36.3	3631.4	55.6	4940.7	66.7	6667.5	66.7	6666.8	99.9	9999.9	63.6	6075.6
Crustacea																				
Hexanauplia																				
Copepoda (subclass)	44.0	4400.8	14.6	1463.2	<0.1	8.5	30.6	3059.5	24.7	2473.9	18.3	101.6	32.2	2147.4	19.4	1296.1	84.6	8460.6	29.6	1881.1
Calanoida									14.1	1253.6									13.5	612.6
Acartia sp.	0.1	1.0					<0.1	<0.1	<0.1	0.2										
Calanus helgolandicus	2.6	28.8	<0.1	0.2			1.0	16.4	1.1	12.1										
Paracalanus sp.			<0.1	<0.1					<0.1	<0.1										
Candacia sp.	14.2	948.2	3.9	391.4	<0.1	1.8	7.3	729.6	7.2	616.5	<0.1	<0.1	5.8	96.0	1.8	89.2	15.9	1061.3	3.9	88.7
Chirundina sp.															0.7	33.6			<0.1	0.3
Eucalanus sp.	0.6	6.8							0.2	0.8										
Euchaeta sp.	3.4	112.8							1.1	12.5			3.9	65.1					1.1	4.8
Heterorhabdus papilliger			0.3	21.1					<0.1	1.0										
Mesocalanus tenuicornis							<0.1	<0.1	<0.1	<0.1			<0.1	0.1					<0.1	<0.1
Metridia sp.	1.0	11.0							0.3	1.2										
Pleuromamma sp.			<0.1	0.1	<0.1	0.4	7.2	598.2	1.6	53.3					13.4	671.3	25.8	1719.7	4.7	64.6
Scolecithrix sp.			3.4	283.5	<0.1	<0.1	0.4	17.8	0.8	27.8					0.6	38.1	4.4	145.9	0.7	8.8
Temora longicornis							<0.10	0.1	<0.1	<0.1										
Temora stylifera	0.4	8.2					1.1	52.9	0.4	6.6	0.3	3.4							0.1	0.5
Unidentified	3.4	113.0	<0.1	0.1	<0.1	0.1	0.4	6.8	1.2	31.7	7.2	159.4							2.9	26.7
Cyclopoida									4.1	414.8									1.9	71.5
Corycaeus sp.	1.9	129.3	1.4	120.4	<0.1	2.3	3.9	327.9	1.8	150.5			3.2	53.2					0.8	3.9
Oithona sp.	0.1	0.7	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	0.6							0.1	4.5	<0.1	<0.1
Oncaea sp.	4.2	422.2	0.9	90.1	<0.1	0.5	0.5	48.3	1.7	171.6	0.5	17.5	2.9	95.7			<0.1	0.6	0.9	27.3
Sapphirina sp.			1.6	137.0			0.9	29.9	0.6	14.6					0.9	42.8			<0.1	0.4
Harpacticoida	<0.1	<0.1	0.1	1.9	<0.1	0.7	1.2	58.5	0.3	11.2										
Unidentified Copepoda	12.1	1212.2	2.9	293.6	<0.1	1.0	6.8	682.6	6.2	621.2	10.3	571.7	16.5	1098.3	2.1	106.5	38.4	3836.2	14.1	834.8
Cirripedia (Infraclass)			<0.1	<0.1					<0.1	<0.1										
Malacostraca	11.6	513.5	1.0	54.5	0.2	20.7	<0.1	2.0	4.1	245.8	15.9	528.4	<0.1	0.1	45.0	3753.3	3.8	127.0	14.1	513.8
Amphipoda	0.8	18.1	<0.1	<0.1	<0.1	1.1	<0.1	0.2	0.3	10.3	<0.1	<0.1			2.9	145.2			0.3	2.4
Decapoda	8.7	291.1	<0.1	0.4	0.1	5.1			2.9	86.9			<0.1	0.1	27.1	2259.1	3.8	127.0	6.0	136.7
Brachyura (Infraorder)	2.0	66.4	1.1	35.6	<0.1	0.7			0.9	30.1	9.2	102.6			1.1	54.8			3.9	35.2

Anomura (Infraorder)					0.1	6.5	<0.1	0.6	<0.1	0.6	6.6	73.4			13.9	696.5			3.9	36.1
Isopoda					<0.1	0.1			<0.1	<0.1										
Mysida	<0.1	0.3							<0.1	<0.1										
<b>Ostracoda</b>			<b>0.1</b>	<b>2.0</b>					<b>&lt;0.1</b>	<b>0.1</b>										
<b>Crustacea eggs</b>	<b>5.4</b>	<b>120.2</b>	<b>&lt;0.1</b>	<b>0.1</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>0.6</b>	<b>30.0</b>	<b>0.8</b>	<b>78.7</b>	<b>&lt;0.1</b>	<b>0.6</b>	<b>&lt;0.1</b>	<b>2.2</b>	<b>&lt;0.1</b>	<b>0.6</b>	<b>&lt;0.1</b>	<b>1.2</b>		
Copepoda eggs	<0.1	<0.1	<0.1	<0.1			<0.1	<0.1	<0.1	<0.1										
Unidentified eggs	<0.1	1.1	3.1	307.9	<0.1	<0.1	0.4	44.7	0.8	78.7	<0.1	0.6	<0.1	2.2	<0.1	0.6	<0.1	1.2	<0.1	1.5
<b>Unidentified Crustacea</b>	<b>3.3</b>	<b>334.4</b>	<b>0.5</b>	<b>47.6</b>	<b>&lt;0.1</b>	<b>0.8</b>	<b>0.7</b>	<b>67.9</b>	<b>1.4</b>	<b>137.3</b>	<b>15.0</b>	<b>998.4</b>	<b>3.1</b>	<b>204.3</b>	<b>1.0</b>	<b>80.1</b>	<b>10.9</b>	<b>1093.5</b>	<b>8.5</b>	<b>621.2</b>
<b>Bivalvia</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>0.2</b>	<b>7.6</b>	<b>&lt;0.1</b>	<b>0.1</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>1.0</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>							<b>&lt;0.1</b>	<b>&lt;0.1</b>
<b>Gastropoda</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>0.6</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>6.3</b>	<b>347.4</b>	<b>31.3</b>	<b>3133.0</b>	<b>1.2</b>	<b>81.2</b>	<b>0.6</b>	<b>20.2</b>	<b>0.6</b>	<b>20.2</b>		
Pteropoda			<0.1	<0.1	<0.1	0.6	<0.1	<0.1	<0.1	<0.1	5.8	324.7	28.2	2823.5	1.2	81.2	0.6	20.2	10.3	654.5
<b>Gastropoda eggs</b>	<b>0.4</b>	<b>19.6</b>	<b>0.1</b>	<b>5.2</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>0.2</b>	<b>8.9</b>	<b>0.2</b>	<b>4.3</b>					<b>&lt;0.1</b>	<b>0.8</b>	<b>&lt;0.1</b>	<b>1.1</b>
<b>Unidentified Gastropoda</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>							<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>0.4</b>	<b>4.5</b>	<b>3.1</b>	<b>51.6</b>					<b>1.0</b>	<b>9.1</b>
<b>Fish eggs</b>	<b>5.4</b>	<b>120.2</b>	<b>&lt;0.1</b>	<b>0.1</b>	<b>&lt;0.1</b>	<b>&lt;0.1</b>	<b>0.6</b>	<b>30.0</b>	<b>1.9</b>	<b>57.4</b>										
<b>Microzooplankton</b>	<b>8.6</b>	<b>864.5</b>	<b>0.1</b>	<b>11.2</b>	<b>&lt;0.1</b>	<b>0.2</b>	<b>0.9</b>	<b>94.4</b>	<b>3.1</b>	<b>300.2</b>										
<b>Echinodermata (phylum)</b>	<b>4.5</b>	<b>400.1</b>					<b>0.5</b>	<b>44.6</b>	<b>1.6</b>	<b>77.9</b>										
<b>Foraminifera (phylum)</b>	<b>0.2</b>	<b>14.3</b>							<b>&lt;0.1</b>	<b>1.6</b>										
<b>Radiozoa (phylum)</b>			<b>&lt;0.1</b>	<b>&lt;0.1</b>					<b>&lt;0.1</b>	<b>&lt;0.1</b>										
<b>Ciliophora (phylum)</b>																				
Tintinnina (suborder)	4.0	354.0	0.1	11.2	<0.1	0.2	0.4	34.1	1.4	128.4										
<b>Tunicata</b>	<b>0.1</b>	<b>3.0</b>							<b>&lt;0.1</b>	<b>0.3</b>										
<b>Appendicularia</b>																				
Copelata																				
<i>Oikopleura</i> sp.	0.1	1.5							<0.1	0.2										
<b>Thaliacea</b>																				
Salpida	<0.1	<0.1							<0.1	<0.1										
<b>Teleostei</b>	<b>18.9</b>	<b>421.9</b>	<b>58.3</b>	<b>3884.3</b>	<b>99.7</b>	<b>9967.2</b>	<b>45.2</b>	<b>2258.5</b>	<b>51.5</b>	<b>2859.1</b>	<b>33.3</b>	<b>1110.2</b>	<b>33.3</b>	<b>1110.8</b>	<b>33.3</b>	<b>1111.1</b>			<b>31.8</b>	<b>1011.9</b>
<b>Epipelagic fish</b>			<b>16.6</b>	<b>276.7</b>	<b>66.4</b>	<b>4429.3</b>	<b>33.3</b>	<b>1108.7</b>	<b>25.8</b>	<b>670.0</b>					<b>28.6</b>	<b>954.2</b>			<b>7.8</b>	<b>70.9</b>
Clupeidae																				
<i>Sardina pilchardus</i>							16.7	277.6	3.7	13.7										
<i>Unidentified</i>			16.6	276.7					3.7	13.7					8.5	141.3			2.3	10.5
Centriscidae																				



## Discussion

This work is the first to describe the diet of *S. colias* and *T. picturatus* in the archipelago of Madeira. It also provides novel data on the zooplanktonic communities in this poorly known oceanic region of the Atlantic. The size classes analysed in this study correspond to a narrow size class range (modal size class in the fishery), which excludes ontogenetic variations of the diet. Our data showed that both species fed on planktonic and micronektonic prey from the epipelagic environment, with a modest contribution of mesopelagic prey. A seasonal variation was observed in both mackerel's diet which seems to be related to the availability of fish and cephalopods, temporarily replacing the smaller sized planktonic species. *S. colias* fed on a wider variety of prey than *T. picturatus*, which overall resulted in moderate overlap in their diet compositions and isotopic niches.

Around Madeira, the diet of *S. colias* suggests that this species is an opportunistic forager, while *T. picturatus* showed to have a more selective diet that incorporates larger prey and a higher degree of piscivory. These results agree with other studies in the Mediterranean (Battaglia *et al.*, 2019; Sever *et al.*, 2006), subtropical and tropical eastern Atlantic (Castro and Santana del Pino, 1995; Denda *et al.*, 2017; Gushchin and Corten, 2017; Kompowski, 1976) and Iberian coast (Bachiller and Irigoien, 2015; Garrido *et al.*, 2015). The wider prey variety presented by the *S. colias* has been previously described (Bachiller and Irigoien, 2015) and is most probably related to a higher ability of the feeding apparatus to retain smaller prey and an eventual filter-feeding behaviour (Costalago *et al.*, 2015). This opportunistic feeding behaviour may explain the more diverse diet found in this study than in studies conducted in other regions (e.g. Castro and Santana del Pino, 1995; Garrido *et al.*, 2015; Wahbi *et al.*, 2015).

Among the planktonic prey, calanoid copepods were the most consumed by both mackerel species in this study, but this is also true in other studies performed in the Portuguese coast (Garrido *et al.*, 2015) and the Àmpere seamount (Denda *et al.*, 2017), for example. Mysids and euphausiids, which had important roles in the diet of these mackerels in other regions (Bachiller and Irigoien, 2015; Battaglia *et al.*, 2019; Castro, 1993; Kompowski, 1976; Sever *et al.*, 2006;), and in a previous study conducted in Madeira (Costa *et al.*, 2013), were infrequent in our study. The differences found between Costa *et al.* (2013) and our study suggests long-term changes in the abundance of those species in the Madeira oceanic region, suggesting the existence of a dynamic planktonic community.

Fish was an important prey in the diet of *S. colias* and *T. picturatus* around the Madeira waters, alongside zooplanktonic species. To the best of our knowledge, this was only described for *S. colias* in two other studies off Mauritania and Turkey coasts (Gushchin and Corten, 2017; Sever *et al.*, 2006), as fish are often reported having little importance in the Atlantic and in the Mediterranean. Species such as *M. scolopax*, *S. pilchardus*, myctophids (Castro, 1993; Denda *et al.*, 2017; Gushchin and Corten, 2017; Sever *et al.*, 2006; Wahbi *et al.*, 2015), and *S. saurus* (this study) can be found in mackerel's stomachs in these regions. Our results also contrast with data obtained in the Portuguese coast, where both mackerel species have a strictly planktonic diet (Garrido *et al.*, 2015). Although the slightly lower average size of the *T. picturatus* analysed in Garrido *et al.* (2015) (ca.18 cm, total length) might contribute to partly explain the difference in diet, as younger individuals tend to consume smaller crustaceans, like copepods, euphausiids and decapod larvae (Kompowski, 1976), the same does not hold for *S. colias*. It is possible that the disparity found in the diet of the *S. colias* in different regions is linked to its opportunistic behaviour (Hernández and Ortega, 2000) and to different prey communities in different areas.

Cannibalism has been previously reported in *S. colias* and other pelagic forage fish, such as *S. pilchardus* (Castro, 1993; Garrido *et al.*, 2008), supporting the same conclusions reached for *S. colias* in this study. On the other hand, this was not observed for *T. picturatus* here, which consumed mainly *M. scolopax* and myctophids, following the results found in other studies (Battaglia *et al.*, 2019; Hirsch and Christiansen, 2010). Unlike what has been reported in other studies of individuals of similar sizes (e.g. Battaglia *et al.*, 2019; Kompowski, 1976), cephalopods were not consumed by *T. picturatus* in our study. The reasons for this behaviour remain unclear, as cephalopods are diverse and abundant in this region (Hastie *et al.*, 2009).

This study found seasonal variations in the diets of the *S. colias* and *T. picturatus*. Zooplanktonic (and also phytoplanktonic for *S. colias*) species were constantly present in the stomachs of both mackerels, which indicates that they comprise the base of the diet for these species, being available all year round. Although planktonic species are easier to capture, their caloric content is lower than that of bigger more nutritious prey, which, on its turn, also require more energy to be preyed. Here, we hypothesise that the seasonal differences observed in the diet of these two mackerel species are directly linked to the abundance of planktonic species and the abundance of fish and cephalopods. Mackerels may choose their prey to fit the best energetic balance.



While the two mackerel species fed on similar prey types, there was a wider range of different prey types in the diet of *S. colias* than in the diet of *T. picturatus*, resulting in differences in the index of dietary diversity. These differences in the relative importance of different prey also resulted in a low dietary overlap from the point of view of the *S. colias*, and in the smaller isotopic niche of the *T. picturatus*. This difference in the diet is further noticeable in their trophic levels, which is significantly higher in the *T. picturatus* than in the *S. colias*. In general, *S. colias* seem to have consumed more fish but also ingested phytoplankton, which can reduce their trophic level. This opportunistic behaviour likely confers an advantage for *S. colias*, which are able to prey on a higher diversity of prey sizes and prey types in this oligotrophic environment. Such species, like these mackerels, are excellent indicators of the existing planktonic communities (Reid *et al.*, 2001). Further studies on their seasonal variations may shed more light on the communities comprising the base of oceanic food webs like the pelagic ecosystem of the archipelago of Madeira.

## Acknowledgements

This work was supported by Fundação para a Ciência e a Tecnologia (FCT Portugal) through the project Oceanwebs (PTDC/MAR-PRO/0929/2014), the strategic projects UIDB/04292/2020 and UIDP/04292/2020 granted to MARE, and UIDB/50017/2020 and UIDP/50017/2020 granted to CESAM. This study was partially supported by the Oceanic Observatory of Madeira Project (M1420-01-0145-FEDER-000001-Observatório Oceânico da Madeira-OOM). PD/BD/127810/2016 was granted to JR by FCT. MH was supported by a postdoctoral grant from ARDITI (Agência Regional para o Desenvolvimento da Investigação, Tecnologia e Inovação), Project M1420-09-5369-FSE-000001. Thanks are due to Fátima Quintela and Alexandra D. Silva for helping with zooplankton and phytoplankton identification, respectively.

## References

- Alonso, H., Granadeiro, J. P., Waap, S., Xavier, J., Symondson, W. O. C., Ramos, J. A., & Catry, P. (2014). An holistic ecological analysis of the diet of Cory's shearwaters using prey morphological characters and DNA barcoding. *Molecular Ecology*, **23**, 3719-3733.
- Bachiller, E., & Irigoien, X. (2015). Trophodynamics and diet overlap of small pelagic fish species in the Bay of Biscay. *Marine Ecology Progress Series*, **534**, 179-198.

- Battaglia, P., Pagano, L., Consoli, P., Esposito, V., Granata, A., Guglielmo, L., Pedá, C., Romeo, T., Zagami, G., Vicchio, T. M., Guglielmo, R., & Andaloro, F. (2019). Consumption of mesopelagic prey in the Strait of Messina, an upwelling area of the central Mediterranean Sea: Feeding behaviour of the blue jack mackerel *Trachurus picturatus* (Bowdich, 1825). *Deep-Sea Research Part I: Oceanographic Research Papers*, **155**, 1-13.
- Castro, J. J. (1993). Feeding ecology of chub mackerel *Scomber japonicus* in the Canary Islands area. *South African Journal of Marine Science*, **13**, 323-328.
- Castro, J. J., & Santana del Pino, A. (1995). Feeding preferences of *Scomber japonicus* in the Canary Islands area. *Scientia Marina*, **59**, 325-333.
- Collete, B. B., & Nauen, C. E. (1983). FAO species catalogue, Vol. 2. Scombrids of the world. An annotated and illustrated catalogue of tunas, mackerels, bonitos and related species known to date. *FAO Fisheries Synopsis*, **125**, 137.
- Costa, G., Santamaria, M. T. G., Vasconcelos, J., Perera, C. B., & Melo-Moreira, E. (2013). Endoparasites of *Trachurus picturatus* (Pisces: Carangidae) from the Madeira and Canary Islands: Selecting parasites for use as tags. *Scientia Marina*, **77**, 61-68.
- Costalago, D., Garrido, S., & Palomera, I. (2015). Comparison of the feeding apparatus and diet of European sardines *Sardina pilchardus* of Atlantic and Mediterranean waters: ecological implications. *Journal of Fish Biology*, **86**, 1348-1362.
- Denda, A., Stefanowitsch, B., & Christiansen, B. (2017). From the epipelagic zone to the abyss: Trophic structure at two seamounts in the subtropical and tropical Eastern Atlantic - Part II Benthopelagic fishes. *Deep-Sea Research Part I: Oceanographic Research Papers*, **130**, 78-92.
- Garcia, A., Pereira, J. G., Canha, Â., Reis, D., & Diogo, H. (2015). Life history parameters of blue jack mackerel *Trachurus picturatus* (Teleostei: Carangidae) from north-east Atlantic. *Journal of the Marine Biological Association of the United Kingdom*, **95**, 401-410.
- Garrido, S., Ben-Hamadou, R., Oliveira, P. B., Cunha, M. E., Chícharo, M. A., & Van Der Lingen, C. D. (2008). Diet and feeding intensity of sardine *Sardina pilchardus*: Correlation with satellite-derived chlorophyll data. *Marine Ecology Progress Series*, **354**, 245-256.
- Garrido, S., Silva, A., Pastor, J., Dominguez, R., Silva, A. V., & Santos, A. M. (2015). Trophic ecology of pelagic fish species off the Iberian coast: diet overlap,

- cannibalism and intraguild predation. *Marine Ecology Progress Series*, **539**, 271-286.
- Giménez, J., Marçalo, A., Ramírez, F., Verborgh, P., Gauffier, P., Esteban, R., Nicolau, L., González-Ortegón, E., Baldó, F., Vilas, C., Vingada, J., Forero, M. G., & De Stephanis, R. (2017). Diet of bottlenose dolphins (*Tursiops truncatus*) from the Gulf of Cadiz: Insights from stomach content and stable isotope analyses. *PLoSOne*, **12**, e0184673. <https://doi.org/10.1371/journal.pone.0184673>
- Gushchin, A. V., & Corten, A. (2017). Feeding of pelagic fish in waters of Mauritania: 3. Atlantic Chub mackerel *Scomber colias*, Atlantic horse mackerel *Trachurus trachurus*, Cunene horse mackerel *Trachurus trecae*. *Journal of Ichthyology*, **57**, 410-423.
- Hastie, L., Pierce, G., Wang, J., Bruno, I., Moreno, A., Piatkowski, U., & Robin, J. (2009). Cephalopods in the North-eastern Atlantic: Species, biogeography, ecology, exploitation and conservation. In R. N. Gibson, R. J. A. Atkinson & J. D. M. Gordon (Eds.), *Oceanography and Marine Biology: An Annual Review* (pp. 111-190). Taylor & Francis.
- Hayse, J. W. (1990). Feeding habits, age, growth, and reproduction of Atlantic spadefish *Chaetodipterus faber* (Pisces: Ephippidae) in South Carolina. *Fishery Bulletin*, **88**, 67-83.
- Hermida, M., & Delgado, J. (2016). High trophic level and low diversity: Would Madeira benefit from fishing down? *Marine Policy*, **73**, 130-137.
- Hernández, J. J. C., & Ortega, A. T. S. (2000). Synopsis of Biological Data on The Chub Mackerel (*Scomber japonicus* Houttuyn, 1782). FAO Fishery Synopsis. Rome, FAO.
- Hirsch, S., & Christiansen, B. (2010). The trophic blockage hypothesis is not supported by the diets of fishes on Seine Seamount. *Marine Ecology*, **31**, 107-120.
- IPIMAR (2009). Manual de Amostragem Biológica de Sardinha, pp. 45.
- Jackson, A. L., Inger, R., Parnell, A. C., & Bearhop, S. (2011). Comparing isotopic niche widths among and within communities: SIBER – Stable Isotope Bayesian Ellipses in R. *Journal of Animal Ecology*, **80**, 595-602.
- Kompowski, A. (1976). A study on the food and feeding habits of *Trachurus trachurus*, *Tr. trecae*, *Tr. picturatus* and *Caranx rhonchus* in the region of Cape Blanc. *Acta Ichthyologica et Piscatoria*, **VI**, 35-57.
- Logan, J. M., Rodríguez-Marín, E., Goñi, N., Barreiro, S., Arrizabalaga, H., Golet, W.,

- & Lutcavage, M. (2011). Diet of young Atlantic bluefin tuna (*Thunnus thynnus*) in eastern and western Atlantic foraging grounds. *Marine Biology*, **158**, 73-85.
- Lozano-Bilbao, E., Díaz, Y., Lozano, G., Jurado-Ruzafa, A., Hardisson, A., Rubio, C., Jiménez, S., González-Weller, D., & Gutiérrez, Á. J. (2019). Metal Content in Small Pelagic Fish in the North-West Africa. *Thalassas: An International Journal of Marine Sciences*, **35**, 643-653.
- Marçalo, A., Nicolau, L., Giménez, J., Ferreira, M., Santos, J., Araújo, H., Silva, A., Vingada, J., & Pierce, G. J. (2018). Feeding ecology of the common dolphin (*Delphinus delphis*) in Western Iberian waters: has the decline in sardine (*Sardina pilchardus*) affected dolphin diet? *Marine Biology*, **165**, 44.
- Martins, H. R. (1982). Biological studies of the exploited stock of *Loligo forbesi* (Mollusca: Cephalopoda) in the Azores. *Journal of the Marine Biological Association of the United Kingdom*, **62**, 799-808.
- Menezes, G. M., Rosa, A., Melo, O., & Pinho, M. R. (2009). Demersal fish assemblages off the Seine and Sedlo seamounts (northeast Atlantic). *Deep-sea Research Part II: Topical Studies in Oceanography*, **56**, 2683-2704.
- Post, D. M. (2002). Using stable isotopes to estimate trophic position: models, methods, and assumptions. *Ecology*, **83**, 703-718.
- Preston, T., & Owens, N. J. P. (1983). Interfacing an Automatic Elemental Analyser with an Isotope Ratio Mass Spectrometer: The Potential for Fully Automated Total Nitrogen and Nitrogen-15 Analysis. *Analyst*, **108**, 971-977.
- R Core Team (2018). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. Available at: <http://www.R-project.org/>.
- Raimundo, J., Vale, C., Caetano, M., Giacomello, E., Anes, B., & Menezes, G. M. (2013). Natural trace element enrichment in fishes from a volcanic and tectonically active region (Azores archipelago). *Deep-sea Research Part II: Topical Studies in Oceanography*, **98**, 137-147.
- Reid, P., De Fatima Borges, M., & Svendsen, E. (2001). A regime shift in the North Sea circa 1988 linked to changes in the North Sea horse mackerel fishery. *Fisheries Research*, **50**, 163-171.
- Romero, J., Catry, P., Menezes, D., Coelho, N., Silva, J. P., & Granadeiro, J. P. (2019). A gull that scarcely ventures on the ocean: yellow-legged gulls *Larus michahellis atlantis* on the oceanic island of Madeira. *Ardeola*, **66**, 101-112.

- Sever, T. M., Bayhan, B., Bilecenoglu, M., & Mavili, S. (2006). Diet composition of the juvenile chub mackerel (*Scomber japonicus*) in the Aegean Sea (Izmir Bay, Turkey). *Journal of Applied Ichthyology*, **22**, 145-148.
- Shabonev, I. Y., & Kotlyar, A. N. (1979). A comparative morpho-ecological analysis of the eastern Pacific forms of *Trachurus symmetricus* and the Atlantic Ocean horse mackerel, *Trachurus picturatus picturatus*. *Journal of Ichthyology*, **19**, 24-29.
- Shabonev, I. Y., & Ryazantseva, Y. I. (1977). Population structure of the oceanic horse mackerel (*Trachurus picturatus picturatus*). *Journal of Ichthyology*, **17**, 954-958.
- Tejerina, R., Hermida, M., Faria, G., & Delgado, J. (2019). The purse-seine fishery for small pelagic fishes off the Madeira Archipelago. *African Journal of Marine Science*, **41**, 373-383.
- Vander Zanden, M., & Rasmussen, J. (2001). Variation in  $\delta^{15}\text{N}$  and  $\delta^{13}\text{C}$  trophic fractionation: Implications for aquatic food web studies. *Limnology and Oceanography*, **46**, 2061-2066.
- Vasconcelos, J., Afonso-Dias, M., & Faria, G. (2012). Atlantic chub mackerel (*Scomber colias*) spawning season, size and age at first maturity in Madeira waters. *Arquipélago. Life and Marine Sciences*, **29**, 43-51.
- Vasconcelos, J., Alves, A., Gouveia, E., & Faria, G. (2006). Age and growth of the Blue jack mackerel, *Trachurus picturatus* Bowdich, 1825 (Pisces: Teleostei) off the Madeira Arquipélago. *Life and Marine Sciences*, **23A**, 47-57.
- Vasconcelos, J., Dias, M. A., & Faria, G. (2011). Age and growth of the Atlantic chub mackerel *Scomber colias* Gmelin, 1789 off Madeira Island. *Arquipélago. Life and Marine Sciences*, **28**, 57-70.
- Vasconcelos, J., Sousa, R., Riera, R., Delgado, J., Faria, G., & Gordo, L. S. (2018). Stock assessment of the blue jack mackerel, *Trachurus picturatus*, in the North-eastern Atlantic. *Fisheries Management and Ecology*, **25**, 233-239.
- Wahbi, F., Tojo, N., Ramzi, A., Somoue, L., Manchih, K., & Errhif, A. (2015). Seasonal and size-dependent variability in diet of *Scomber colias* (Gmelin, 1789) of the Atlantic Coast of the Northwest Africa. *International Journal of Advanced Research*, **3**, 485-497.
- Wallace, R., & Ramsey, J. (1983). Reliability in Measuring Diet Overlap. *Canadian Journal of Fisheries and Aquatic Sciences*, **40**, 347-351.

## Supporting Information

Table S1 - Number of individuals collected for each month and species (Blue jack mackerel *Trachurus picturatus* and Atlantic chub mackerel *Scomber colias*) in the south of Madeira island, between February 2017 and January 2018, and respective range of total lengths.

Date	<i>Scomber colias</i>		<i>Trachurus picturatus</i>	
	N	TL <sup>†</sup>	N	TL
February 2017	20	24.3±1.8 (20.5-27.9)	20	22.2±1.3 (19.8-24.6)
March 2017	20	21.3±1.2 (19.6-23.6)	20	20.3±1.2 (18.7-24.5)
April 2017	20	22.4±0.9 (20.9-24.2)	20	19.5±0.6 (18.6-20.6)
May 2017	21	21.0±1.9 (18.3-25.2)	19	19.8±0.7 (18.4-21.0)
June 2017	20	25.3±1.3 (22.6-27.1)	20	20.2±0.8 (18.7-21.6)
September 2017	23	20.0±2.2 (17.2-24.6)	4	18.1±2.0 (15.3-19.6)
October 2017	20	22.6±1.7 (20.8-28.1)	20	19.9±1.7 (17.5-24.8)
November 2017	20	21.5±0.7 (20.4-22.5)	-	-
January 2018	20	23.6±0.9 (21.8-25.7)	20	19.6±1.0 (17.7-21.0)

<sup>†</sup>Total length

Table S2 - Number of stomachs analysed for each month and species (Blue jack mackerel *Trachurus picturatus* and Atlantic chub mackerel *Scomber colias*).

Date	<i>Trachurus picturatus</i>	<i>Scomber colias</i>	
	≥200µm	≥200µm	<200µm
February 2017	9	9	-
March 2017	9	9	3
April 2017	9	9	3
May 2017	9	9	3
June 2017	9	9	-
September 2017	3	9	3
October 2017	9	9	6
November 2017	-	9	-
January 2018	9	9	6
Total	66	81	24

Table S3 - Carbon content value of each prey item identified in the stomachs of the Atlantic chub mackerel *Scomber colias* and the Blue jack mackerel *Trachurus picturatus*, and references from which the data were obtained.

Prey	Final Carbon content value (µg)	Mean length and mass reference	Reference
Phytoplankton n.i.†	0,00145	-	(Garrido <i>et al.</i> , 2008)
Dinophyceae n.i.	0,00327	-	(Garrido <i>et al.</i> , 2008)
Dinophyceae cyst	0,00327	-	(Garrido <i>et al.</i> , 2008)
<i>Dinophysis</i> sp.	0,00495	-	(Garrido <i>et al.</i> , 2008)
<i>Gonyaulax</i> sp.	0,00536	(Espinoza & Bertrand, 2008)	(Espinoza and Bertrand, 2008)
<i>Lingulodinium polyedra</i>	0,00369	-	(Lewis and Hallett 1997)
<i>Gymnodinium</i> sp.	0,00225	-	(Menden-Deuer & Lessard, 2000)
<i>Scrippsiella</i> sp.	0,00123	-	(Garrido <i>et al.</i> , 2008)
<i>Protoperidinium</i> sp.	0,02257	-	(Garrido <i>et al.</i> , 2008)
<i>Prorocentrum</i> sp.	0,00205	-	(Garrido <i>et al.</i> , 2008)
Bacillariophyceae n.i.	0,00088	-	Mean values calculated from different prey from this study This study mean values
Pennales (Bacillariophyceae)	0,00072	-	(Garrido <i>et al.</i> , 2008)
<i>Achnanthes</i> sp.	0,00145	-	= Phytoplankton n.i.
<i>Cocconeis</i> sp.	0,00006	(Espinoza & Bertrand, 2008)	(Espinoza & Bertrand, 2008)
<i>Pyramimonas</i> sp. (Chlorophyta)	0,00145	-	= Phytoplankton n.i.
<i>Coccolithophora</i> sp.	0,00145	-	= Phytoplankton n.i.
Tintinnina n.i.	0,18062	-	(Garrido <i>et al.</i> , 2008)
Equinodermata n.i.	1,85000	Larvae 1mm; (Espinoza & Bertrand, 2008)	(Espinoza & Bertrand, 2008)
Foraminifera n.i.	0,01460	-	(Garrido <i>et al.</i> , 2008)
Radiozoa n.i.	0,00045	-	(Espinoza & Bertrand, 2008)
Crustacea n.i.	1,86038	-	(Garrido <i>et al.</i> , 2008)
Crustaceaeggs n.i.	0,01469	-	(Garrido <i>et al.</i> , 2008)
Crustaceaegg laying n.i.	0,86795	-	= Copepoda eggs
Copepoda n.i.	4,07082	-	(Garrido <i>et al.</i> , 2008)
Copepoda eggs	0,86795	-	(Garrido <i>et al.</i> , 2008)
Copepoda eggsack	0,01469	-	(Garrido <i>et al.</i> , 2008)
Calanoida n.i.	26,47974	-	Mean values obtained from Calanoid species detected in this study
<i>Acartia</i> sp.	4,02296	-	(Garrido <i>et al.</i> , 2008)
<i>Chirundina</i> sp.	26,47974	-	= Calanoida n.i.
<i>Calanus helgolandicus</i>	63,53851	-	(Garrido <i>et al.</i> , 2008)
<i>Mesocalanus tenuicornis</i>	26,47974	-	= Calanoida n.i.
<i>Candacia</i> sp.	23,42223	-	(Garrido <i>et al.</i> , 2008)
<i>Eucalanus</i> sp.	30,03926	-	(Garrido <i>et al.</i> , 2008)
<i>Euchaeta</i> sp.	37,00245	-	(Garrido <i>et al.</i> , 2008)

<i>Heterorhabdus papilliger</i>	26,47974	-	= Calanoida n.i.
<i>Metridia</i> sp.	48,66667	-	(Hopkins <i>et al.</i> , 1984)
<i>Paracalanus</i> sp.	14,85697	-	(Garrido <i>et al.</i> , 2008)
<i>Pleuromamma</i> sp.	22,02708	-	(Garrido <i>et al.</i> , 2008)
<i>Scolecithrix</i> sp.	22,49178	-	(Espinoza & Bertrand, 2008)
<i>Temora longicornis</i>	38,84597	-	(Garrido <i>et al.</i> , 2008)
<i>Temora stylifera</i>	6,27166	-	(Garrido <i>et al.</i> , 2008)
<i>Corycaeus</i> sp.	12,98689	-	(Garrido <i>et al.</i> , 2008)
<i>Oithona</i> sp.	0,93131	-	(Garrido <i>et al.</i> , 2008)
<i>Oncaea</i> sp.	0,63272	-	(Garrido <i>et al.</i> , 2008)
<i>Sapphirina</i> sp.	16,86000	TL <sup>‡</sup> (Lopes <i>et al.</i> , 2007)	(Lopes <i>et al.</i> , 2007)
Harpacticoida n.i.	4,84607	-	(Garrido <i>et al.</i> , 2008)
Parasite Copepoda	4,07082	-	= Copepoda n.i.
Amphipoda n.i.	7,62509	-	(Garrido <i>et al.</i> , 2008)
Decapoda n.i.	137,116	-	(Garrido <i>et al.</i> , 2008)
Brachyura n.i.	43,14121	-	(Garrido <i>et al.</i> , 2008)
Anomura n.i. (Megalops larvae)	137,116	CL <sup>§</sup> = 1333µm; This study	(Nikolioudakis <i>et al.</i> , 2012)
Isopoda n.i.	80,000	-	(Elizalde <i>et al.</i> , 1999)
Mysida n.i.	60,15672	-	(Garrido <i>et al.</i> , 2008)
Ostracoda n.i.	2,55884	(ostracodjuvenile) = 6.035µg dm <sup>¶</sup> (Borme <i>et al.</i> , 2009)	(Van Der Lingen, 2002)
Cirripedia n.i.	26,16838	-	(Garrido <i>et al.</i> , 2008)
Bivalvia	45,49852	TL= 1.25mm; (Espinoza & Bertrand, 2008) dm (James, 1987)	(Van Der Lingen, 2002)
Lamelibranchia veligera (Bivalvia)	6,13745	-	(Garrido <i>et al.</i> , 2008)
Cephalopoda	249260	TL = 8cm	(Espinoza & Bertrand, 2008)
Gastropoda n.i.	8,47578	-	(Garrido <i>et al.</i> , 2008)
Gastropoda egg-laying	0,86795	-	= Copepoda eggs
Pteropoda	4,36044	-	(Garrido <i>et al.</i> , 2008)
Appendicularia	5,80192	-	(Garrido <i>et al.</i> , 2008)
Salpidae	10,04400	L <sup>¥</sup> = 6.2mm; Mean value of the maximum and minimum values (Heron <i>et al.</i> , 1988)	(Heron <i>et al.</i> , 1988)
Pisces n.i.	625610	-	Mean values of Pisces in this study
Pisces eggs	13,41405	-	(Garrido <i>et al.</i> , 2008)
<i>Scomberesox saurus</i>	480262	TL = 5cm	(Santos <i>et al.</i> , 2002)
Clupeidae n.i.	1126508	-	= <i>S. pilchardus</i>
<i>Sardina pilchardus</i>	1126508	TL = 8cm	(Coelho, 2009)
<i>Scomber colias</i>	1051304	TL = 8cm	(Vasconcelos <i>et al.</i> , 2011)
<i>Macroramphosus scolopax</i>	462907	TL = 5cm	(Espinoza & Bertrand, 2008)
Myctophidae n.i.	7066,5	L = 2cm; (Espinoza & Bertrand, 2008)	(Froese, 1998)

† n.i.: not identified

‡ TL: Total length

§ CL: Carapace length

¶ dm: dry mass

¥ L: Length



## References

- Borme, D., Tirelli, V., Brandt, S. B., Fonda Umani, S., & Arneri, E. (2009). Diet of *Engraulis encrasicolus* in the northern Adriatic Sea (Mediterranean): ontogenetic changes and feeding selectivity. *Marine Ecology Progress Series*, **392**, 193-209.
- Coelho, P. (2009). Modelação do crescimento de *Sardina pilchardus* (Walbaum, 1792) em duas zonas (oceanograficamente distintas) da costa portuguesa (Masters thesis). Available from Sapientia – Universidade do Algarve: <http://hdl.handle.net/10400.1/513>.
- Elizalde, M., Weber, O., Pascual, A., Sorbe, J. C., & Etcheber, H. (1999). Benthic response of *Munnopsurus atlanticus* (Crustacea Isopoda) to the carbon content of the near-bottom sedimentary environment on the southern margin of the Cap-Ferret Canyon (Bay of Biscay, northeastern Atlantic Ocean). *Deep-sea Research Part II: Topical Studies in Oceanography*, **46**, 2331-2344.
- Espinoza, P., & Bertrand, A. (2008). Revisiting Peruvian anchovy (*Engraulis ringens*) trophodynamics provides a new vision of the Humboldt Current system. *Progress in Oceanography*, **79**, 215–227.
- Froese, R. (1998). Length-weight relationships for 18 less-studied fish species. *Journal of Applied Ichthyology*, **14**, 117-118.
- Garrido, S., Ben-Hamadou, R., Oliveira, P. B., Cunha, M. E., Chícharo, M. A., & Van Der Lingen, C. D. (2008). Diet and feeding intensity of sardine *Sardina pilchardus*: Correlation with satellite-derived chlorophyll data. *Marine Ecology Progress Series*, **354**, 245-256.
- Heron, A. C., McWilliam, P., & Dal Pont, G. 1988. Length-weight relation in the salp *Thalia democratica* and potential of salps as a source of food. *Marine Ecology Progress Series*, **42**, 125-132.
- Hopkins, C. C. E., Tande, K. S., & Grønvik, S. (1984). Ecological investigations of the zooplankton community of Balsfjorden, Northern Norway: an analysis of growth environment in *Metridia longa* (Lubbock), *Calanus finmarchicus* (Gunnerus), *Thysanoessa inermis* (Krøyer) and *T. raschi* (M. Sars). *Journal of Experimental Marine Biology and Ecology*, **82**, 77-99.
- James, A. G. (1987). Feeding ecology, diet and field-based studies on feeding selectivity of the cape anchovy *Engraulis capensis* Gilchrist. *South African Journal of Marine Science*, **5**, 673-692.

- Lewis J., & Hallett, R. (1997). *Lingulodinium polyedrum* (Gonyaulax polyedra) a blooming dinoflagellate. An annual review. In R. N. Gibson & M. Barnes (Eds.). *Oceanography and Marine Biology – volume 35* (pp. 89-93). Applied Ecology Research Group, School of Biological and Health Sciences, University of Westminster, 115 New Cavendish Street, London.
- Lopes, R. M., Dam, H. G., Aquino, N. A., Monteiro-Ribas, W., & Rull, L. (2007). Massive egg production by a salp symbiont, the poecilostomatoid copepod *Sapphirina angusta* Dana, 1849. *Journal of Experimental Marine Biology and Ecology*, **348**, 145-153.
- Menden-Deuer, S., & Lessard, E. J. (2000). Carbon to volume relationships for dinoflagellates, diatoms, and other protist plankton. *Limnology and Oceanography*, **45**, 569-579.
- Nikolioudakis, N., Isari, S., Pitta, P., & Somarakis, S. (2012). Diet of sardine *Sardina pilchardus*: An ‘end-to-end’ field study. *Marine Ecology Progress Series*, **453**, 173-188.
- Santos, M. N., Gaspar, M. B., Vasconcelos, P., & Monteiro, C. C. (2002). Weight-length relationships for 50 selected fish species of the Algarve coast (southern Portugal). *Fisheries Research*, **59**, 289–295.
- Van Der Lingen, C. D. (2002). Diet of Sardine *Sardinops sangax* in the Southern Benguela upwelling ecosystem. *South African Journal of Marine Science*, **24**, 301-316.
- Vasconcelos, J., Afonso-Dias, M., & Faria, G. (2012). Atlantic chub mackerel (*Scomber colias*) spawning season, size and age at first maturity in Madeira waters. *Arquipélago. Life and Marine Sciences*, **29**, 43-51.
- Vasconcelos, J., Dias, M. A., & Faria, G. (2011). Age and growth of the Atlantic chub mackerel *Scomber colias* Gmelin, 1789 off Madeira Island. *Arquipélago. Life and Marine Sciences*, **28**, 57-70.

## CHAPTER 3

---



Credits: Raquel Segur

*This chapter is published as:*

*Joana Romero, Paulo Catry, Margarida Hermida, Verónica Neves, Bárbara Cavaleiro, Lúcia Gouveia, José Pedro Granadeiro (2021). Tunas off northwest Africa: the epipelagic diet of the Bigeye and Skipjack tunas. Fisheries Research 238: 105914.*

# Tunas off northwest Africa: the epipelagic diet of the Bigeye and Skipjack tunas

## Abstract

Tunas are among the most fished top predators worldwide, which has led to the depletion of many of their stocks. The decline of their populations has potentially great impacts in the food-web and the ecosystems. To better understand the impacts in the most important tuna areas, basic knowledge on the diet of these species in each region is required. Here, we describe the diet of the two most fished tuna species in the archipelago of Madeira, the Bigeye tuna *Thunnus obesus* and the Skipjack tuna *Katsuwonus pelamis*, based on stomach contents. To gain further insights into the diet, and also better assess the possible bias caused by the occurrence of live bait in stomachs, we compared tuna mercury values with those of two other predators with similar diets that are not directly targeted by fisheries. Bigeye tunas fed mostly on Atlantic chub mackerel *Scomber colias*, mackerel *Trachurus* sp. and Longspine snipefish *Macroramphosus scolopax*, also consuming some mesopelagic prey, including myctophids and cephalopods. Skipjack tunas had an epipelagic diet constituted mainly of Longspine snipefish and Sand smelt *Atherina* sp.. There were interannual variations in diet likely linked to interannual pelagic community shifts. Bait did not bias the results of the stomach analysis of these tunas and bait species were proved to be part of the natural diet of both tuna species. Baseline data provided by this study should allow for more informed decisions for an efficient ecosystem-based fisheries management.

**Key words:** *Thunnus obesus*, *Katsuwonus pelamis*, Diet, Mercury, Northeast Atlantic Ocean

## Introduction

Marine top predators are among the most threatened functional groups in the wide ocean, with an estimated loss of biomass of large predatory fishes at around 90% in relation to pre-industrial numbers (Myers and Worm 2003). Among this group, tunas are one of the most targeted by the fishery industry. Every year, around half a million tonnes of tunas are captured worldwide (2000-2018; ICCAT 2020). Such pressure has led to the depletion of many stocks and, consequently, to the inclusion of several tuna species in the IUCN Red list of threatened species (IUCN 2020). Furthermore, the decline of top predators impacts the structure of communities and the diversity of food webs, as well as the productivity and connectivity of the ecosystems (Doney et al. 2012). Impacts are not only observed directly on their prey but can produce cascading effects on other marine species and communities. For example, the increase in tuna fishing activity in the 1960's in the eastern tropical Atlantic Ocean and the consequent decline of tuna populations in that area is thought to have caused a shift in the diet and a dramatic decline of the once massive Sooty tern *Onychoprion fuscatus* population of Ascension island which depends on interactions with sub-surface predators to locate and catch prey (Reynolds et al. 2019).

The Bigeye tuna *Thunnus obesus* (Vulnerable) and the Skipjack tuna *Katsuwonus pelamis* (Least concern) are the main targeted tuna species around the archipelago of Madeira (Gouveia et al. 2019), located in the subtropical NE Atlantic Ocean. In this archipelago there is a strong tuna fishing tradition taking place mostly from March to October (Gouveia and Mejuto 2003). Here tunas are caught mostly using pole-and-line, which uses small pelagic fish as live bait. In Madeira, an average of 1667 ( $\pm 481$  SD) and 549 ( $\pm 461$  SD) tonnes of Bigeye and Skipjack tuna, respectively, are landed every year (2007-2017; Gouveia et al. 2019), comprising an average of 35% of local fisheries landings and even reaching 50% in some years (Hermida and Delgado 2016).

Despite the considerable economic importance of tunas in the subtropical and temperate NE Atlantic, more specifically in Madeira, Azores and the Canary Islands, few ecological studies on these species have been conducted in this region. This contrasts with the various studies conducted in the Pacific on their distribution (eg. Houssard et al. 2019; Lehodey et al. 1997), movements (eg. Schaefer et al. 2009), reproduction (eg. Hunter et al. 1986), and contamination (eg. Chen et al. 2014), and also in the South and NW Atlantic (Matthews et al. 1977; Matsumoto and Miyabe 2002; da Silva et al. 2019). Large knowledge gaps on the ecology of these species remain in the Atlantic, such as migratory

routes, breeding periods and diet. To the best of our knowledge, a single study on the Skipjack tuna diet in the Canary Islands was published to this date (Ramos et al. 1995), and none on Bigeye tunas.

Tunas are opportunistic predators which means that their main prey shift according to region, reflecting prey availability in the different ocean compartments (Ménard et al. 2006; Gorni 2016; Ohshimo et al. 2018). Therefore, knowledge on their diet will contribute to our understanding of food web dynamics and allow to infer broad community-scale changes in the abundance, availability, and diversity of poorly known mid-trophic prey. Such knowledge is required for an ecosystem-based fisheries management and the conservation of large pelagic predators, including tunas.

Analysis of stomach contents is a direct and reliable way to thoroughly assess the diet of tunas (eg. Glaser et al. 2015; Varela et al. 2019). However, the use of live bait during fishing events may raise concerns about what proportion of stomach content is bait. Indirect techniques, like determination of mercury concentration in tissues have allowed to infer the diet of several predators for which sampling stomach contents is difficult (Layman et al. 2012; Teffer et al. 2014). Mercury concentrations can help distinguishing between a mesopelagic/bathypelagic diet and an epipelagic one, as deep-ocean species have significantly higher concentrations of mercury than epipelagic species due to the higher rate of microbial mediated methylation of mercury in sub-thermocline low oxygen waters (Choy et al. 2009).

The aim of this study is to describe the diet of two important top predators, the Bigeye and the Skipjack tunas, in the pelagic region around the Madeira archipelago, using stomach content analysis. To control for potential biases due to the use of live bait in this fishery, we also determined the mercury concentration in tissues of tunas and of two other abundant pelagic predators from the region with epipelagic diets, the Yellowmouth barracuda *Sphyraena viridensis* and the Longfin yellowtail *Seriola rivoliana*.

## Methods

### *Tuna sampling*

The stomachs of a total of 71 Bigeye tuna and 61 Skipjack tuna from a total of 16 and 10 fishing events, respectively, were obtained directly from local fishermen. Fish were captured using pole-and-line fishing vessels in Madeira island in 2016/2017 and 2016-2018 for Bigeye and Skipjack tuna, respectively. The individuals were measured

( $\pm 0.5\text{cm}$ ), weighted ( $\pm 1\text{g}$ ), and the stomachs removed and kept frozen until sorting in the laboratory. A sample of blood was also collected for analysis of mercury concentration, and frozen until further processing in the laboratory. We obtained the coordinates of 22 tuna fishing events from local fishermen that took place in September 2017 and May-October 2018 which allowed us to produce an approximate map of the areas used for this fishing activity (Figure 1).




Figure 1 - Main areas (in grey) used by the tuna fishing vessels from Madeira island (95% Kernel Utilization Distribution). Areas were defined using coordinates of tuna fishing events in September 2017 and May-October 2018, as provided by fishermen (n=22). Isobathic lines of 1000, 2000, 3000, 4000m

### *Prey identification*

Once in the laboratory, stomachs were thawed, and the contents carefully removed and classified according to digestion levels and type of prey (fish, cephalopods, crustaceans). Digestion levels for fish and cephalopods were attributed according to Aloncle and Delaporte (1974) and Alonso et al. (2018), respectively (Supplementary Material, Table A.1). Whenever the digestion level corresponded to I or II, fish prey were weighted ( $\pm 0.1\text{g}$ ), measured ( $\pm 0.1\text{cm}$ ) and identified using morphologic characteristics (Whitehead et al. 1989). Individuals with digestion levels of III or higher were identified using our reference collection of fish skeletons and otoliths from the NE Atlantic

(currently comprising over 700 specimens of ca. 100 species). Specific vertebrae were measured in order to obtain size estimations and weights of each individual (Granadeiro and Silva 2000). When identification using hard structures was not possible, a small piece of muscle was collected for DNA barcoding analysis. Cephalopods were identified using the lower beak (Clarke 1986; Lu and Ickeringill 2002) and DNA-barcoding analysis. Beaks with no tissue attached were excluded from the quantitative analysis due to the impossibility of knowing for how long they had been in the stomach. This way we avoid over-representation of cephalopod prey in the tuna's diet. The number of individuals in a sample was estimated by counting all identifiable structures. The Blue jack mackerel is the most common species of the genus *Trachurus* to be found in this marine region. However, due to the similarities of vertebrae between *Trachurus picturatus* and *Trachurus trachurus* and inability to confirm the exact species, we refer to this genus as *Trachurus* sp..

Tissues of 21 fish, 23 cephalopods and 6 crustaceans were sampled from stomach contents of Bigeye tuna and Skipjack tuna from which DNA was extracted using the E.Z.N.A. Tissue DNA kit (Omega Bio-tek). Optimized PCR conditions were used to amplify the 3' end region of the 16S rRNA gene of each prey DNA using the universal primers 16Sar and 16SSbr of Palumbi (1996) (Alonso et al. 2014). Both directions of the PCR products were sequenced in outsourcing (Macrogen Inc.). The resulting sequences were queried using BLAST (NCBI) and searches with similarity values higher than 98% were considered as positive identifications.

#### *Diet characterization*

The diets of Bigeye and the Skipjack tuna were characterized using numeric frequency (NF, number of individuals of a species in relation to the total number of individuals found in the stomachs) and frequency of occurrence (FO, number of stomachs in which a specific prey was found in relation to the total number of stomachs analysed). The Shannon-Wiener diversity index was used to calculate the prey diversity found in the stomachs of the two tuna species. Ontogenetic differences (length classes: 60-80, 80-100 and >100cm) in the diet of Bigeye tunas were assessed using Permutational multivariate analysis of variance (PERMANOVA).

#### *Bait*

In Madeira, Bigeye tuna's fishery involves the use of live bait to attract fish to the surface, particularly Atlantic chub mackerel *Scomber colias* and Blue jack mackerel



*Trachurus picturatus*. On the other hand, Skipjack tuna catches use European pilchard *Sardina pilchardus*, Sand smelt *Atherina* sp. or Bogue *Boops boops*, and when abundant, Longspine snipefish *Macroramphosus scolopax*.

Live bait has been considered a nuisance in diet studies of tuna due to the difficulty to distinguish it from the natural diet (Ankenbrandt 1985; Ramos et al. 1995). To address this issue, we (1) compared mercury concentrations in the tissues of the tunas studied with two other local predatory fishes known to have an epipelagic diet: Yellowmouth barracuda *Sphyræna viridensis* and Longfin yellowtail *Seriola rivoliana*; we also (2) excluded the bait (species known to be used as bait and with digestion levels I and II) from the original stomach content dataset and compared it with the diet inferred from the complete stomach content dataset (NF and FO) (Ankenbrandt 1985; Ramos et al. 1995).

Muscle samples of Yellowmouth barracuda and Longfin yellowtail were obtained from commercial fisheries in Madeira island and from a seizure of illegal fishing in the Selvagens islands, respectively. Each individual was measured ( $\pm 1$ cm) and weighted ( $\pm 1$ g) upon capture. White muscle was collected for measurement of mercury concentration.

Mercury concentrations were analysed using blood for the tuna species and muscle for the two other predators. By using the tuna's blood, we attempted to minimize the contribution of assimilated diet that would have been obtained in more distant waters used by tunas before arriving in Madeira. As it was not possible to collect blood for the two other predator species, white muscle was used instead. These predators are not known to perform long migrations like tunas (McClellan and Cummings 1997; Fontes and Afonso 2017), therefore, both tissues should represent the mercury values accumulated in the marine environment around the archipelago of Madeira. Because mercury accumulates differently among tissues (Storelli et al. 2005; Voegborlo et al. 2007), we also measured the concentration of mercury in the muscle of the same individuals of Bigeye tuna ( $n=20$ ) that we analysed for blood (Supplementary Material, Figure A.1).

The blood and muscle tissues were first lyophilized and reduced into a homogenous powder that was then used to determine the total concentration of mercury. Samples (weighing 3-17mg) were analysed using thermal decomposition atomic absorption spectrometry alongside gold amalgamation in LECO AMA-254 equipment. A maximum coefficient of variation of 10% was defined for a minimum of 2 mercury readings (Furtado et al. 2019). Tort-2 and Tort-3 were used as the certified reference material to support the precision of the method (muscle tissue, certified value of

0.27±0.06 mg·kg<sup>-1</sup> and 0.292±0.22 mg·kg<sup>-1</sup>, respectively). The recovery efficiency for Tort 2 was 82.2±0.8% (n=6), and for Tort 3 was 84.4±2.4% (n=8). Mercury concentrations are reported on a dry weight basis (dw).

All statistical analyses were performed using R version 4.0.2 (R Core Team 2020).

## Results

### *Bigeye tuna diet*

A total of 762 prey items were found in 69 stomachs of Bigeye tuna (2 stomachs were empty), of which 95% were identified to family, genus or species level, resulting in 24 species, 23 genus and 20 different families. Fish was the most consumed prey, followed by cephalopods (Table 1). The Bigeye tuna presented an epipelagic diet (NF=84%) with only 9% of the prey (comprising the genus *Hygophum* and *Diaphus*) belonging to mesopelagic layers. The Atlantic chub mackerel *Scomber colias* was the most consumed species overall (NF=41%), followed by mackerel *Trachurus* sp. (NF=25%) and Longspine snipefish (NF=10%). Cephalopods comprised ca. 3.5% of the prey items. Other cephalopods, represented by older beaks with no remaining flesh, were also found in the stomachs of this tuna species (Supplementary Material, Table A.2). The Shannon-Wiener diversity for the Bigeye tuna was 1.79±0.14 (n=69).

The importance of the main prey of the Bigeye tuna changed between 2016 and 2017, with the number of Atlantic chub mackerel decreasing from 2016 to 2017 (NF=47% (n=642) to 2% (n=120)), and the number of Longspine snipefish increasing from 2% to 54% (NF) (Figure 2). However, Longspine snipefish were only found in 27% of the stomachs (n=11) in 2017 against the 12% of 2016 (n=58), while the Atlantic chub mackerel was found in 72% of the stomachs in 2016 and 45% in 2017. Finally, no variation in the consumption of mesopelagic and epipelagic were found between the different tuna length classes (PERMANOVA:  $F_{2,65} = 1.03$ ,  $r^2 = 0.03$ ,  $p=0.39$ ).

### *Skipjack tuna diet*

A total of 550 prey individuals were found in 57 stomachs of Skipjack tuna (4 empty stomachs), and 98% of the prey were identified resulting in 8 species, 10 genus and 8 families (Table 1). Fish was the most consumed prey by the Skipjack tuna, which also showed a preference for epipelagic species (NF=97%). The Longspine snipefish was the most consumed species (NF=35%), followed by Sand smelt *Atherina* sp. (NF=27%). The Atlantic chub mackerel and the European pilchard also had an important role in the

Skipjack tuna diet (NF=14% and 10%, respectively). The Shannon-Wiener Diversity Index for the Bigeye tuna was  $1.59 \pm 0.12$ .

We found differences among years with the most consumed prey shifting from Sand smelt (NF=55%, n=138) and Atlantic chub mackerel (NF=33%, n=138) in 2016, to Longspine snipefish (NF=38%, n=292; FO=40%, n=30), European pilchard (NF= 19%, n=292) and Sand smelt (NF=16%, n=292) in 2017 (Figure 2). The consumption of Longspine snipefish increased even more in 2018 (NF=84%, n=120; FO=61%, n=13). No myctophids were found in Skipjack stomachs.

Table 1 - Diet of the Bigeye tuna *Thunnus obesus* and the Skipjack tuna *Katsuwonus pelamis* inferred through numeric frequency (NF) and frequency of occurrence (FO). Numbers between brackets represent the sample size

Taxa	Family	Species	Bigeye		Skipjack	
			NF (762)	FO (69)	NF (550)	FO (57)
Teleostei			92.7	98.6	97.8	100
	Scombridae	<i>Scomber colias</i>	41.1	69.6	14.4	47.4
	Carangidae	<i>Trachurus</i> sp.	25.7	69.6	4.5	21.1
	Centriscidae	<i>Macroramphosus scolopax</i>	10.6	14.5	35.8	35.0
	Atherinidae	<i>Atherina</i> sp.			27.1	36.8
	Sparidae	<i>Boops boops</i>	1.2	10.1	4.5	14.0
	Scomberesocidae	<i>Scomberesox saurus</i>	0.8	2.9		
	Clupeidae	<i>Sardina pilchardus</i>			10.4	22.8
		<i>Sardinella</i> sp.			0.2	1.8
		Unidentified			0.7	7.0
	Bramidae	<i>Brama brama</i>	0.3	2.9		
	Molidae	<i>Ranzania laevis</i>	0.3	1.4		
	Alepisauridae	<i>Alepisaurus ferox</i>	0.4	2.9		
	Paralepididae	<i>Arctozenus risso</i>	1.0	7.2		
		<i>Sudis hyalina</i>	0.1	1.4		
	Trichiuridae	Unidentified			0.2	1.8
	Nomeidae	<i>Cubiceps gracilis</i>	2.9	7.2		
	Scopelarchidae	<i>Scopelarchus</i> sp.	0.3	1.4		
	Myctophidae	<i>Diaphus</i> sp.	0.8	5.8		
		<i>Diogenichthys atlanticus</i>	0.1	1.4		
		<i>Gonichtys cocco</i>	0.1	1.4		
		<i>Hygophum hygomii</i>	0.1	1.4		
		<i>Hygophum reinhardtii</i>	0.3	1.4		
		<i>Hygophum taaningi</i>	0.4	1.4		
		<i>Hygophum</i> sp.	4.9	1.4		
		Unidentified fish		1.3	8.7	
Cephalopoda			3.5	15.9	0.2	1.8
	Mastigoteuthidae	<i>Mastigoteuthis</i> sp.	0.1	1.4		

Ommastrephidae	<i>Todarodes sagittatus</i>	0.5	2.9		
	<i>Ommastrephes bartramii</i>			0.2	1.8
	Unidentified	0.5	5.8		
Onychoteuthidae	<i>Onychoteuthis</i> sp.	0.5	2.9		
Decapodiformes	<i>Unidentified</i>	0.3	2.9		
Argonautidae	<i>Argonauta</i> sp.	0.3	2.9		
Octopodidae	Unidentified	0.1	1.4		
Unidentified cephalopod		1.2	4.3		
<b>Arthropoda</b>		<b>3.7</b>	<b>14.5</b>	<b>0.4</b>	<b>1.8</b>
Penaeidae	<i>Funchalia villosa</i>	0.3	2.9		
Decapoda	<i>Unidentified</i>	1.0	1.4		
Platyscelidae	<i>Platyscelus</i> sp.	1.3	1.4		
Unidentified crustacean		1.0	11.6	0.4	1.8
<b>Gastropoda</b>				<b>1.6</b>	<b>1.8</b>
<b>Tunicata</b>		<b>0.1</b>	<b>1.4</b>		
Thaliacea	Unidentified	0.1	1.4		




Figure 2 - Interannual variation of prey found in the stomachs of Bigeye *Thunnus obesus* and Skipjack tuna *Katsuwonus pelamis* captured in Madeira pelagic region in 2016-2018

#### Exclusion of bait from diet

Only trivial differences were found in the relative importance of prey in the diet of these tunas when all bait species with digestion levels I and II were excluded (Supplementary Material, section A1, Figure A.2 and A.3). Furthermore, in the stomach contents of Skipjack tunas captured in 2018 (when we had information on the bait used), we found highly digested remains (digestion state IV) of species used as bait and which should only be found undigested if the tunas only ate them as bait (Supplementary Material, section A.1, Table A.2).

### Mercury concentration comparisons

Differences among predators in mercury concentrations were also found (ANOVA:  $F_{3,82}=22.35$ ,  $p<0.001$ ). Two distinct groups were observed: Bigeye tuna (blood) and Yellowmouth barracuda (muscle), which had the highest values, and Skipjack tuna (blood) and Longfin yellowtail (muscle) which presented the lowest values (Table 2). Only the Bigeye tuna presented a significant positive correlation between mercury concentration in blood and weight ( $r(18)=0.73$ ,  $p=0.0002$  on log transformed data) (Figure 3); individuals  $<100\text{cm}$  had  $1.08\pm0.25 \text{ mg}\cdot\text{kg}^{-1} \text{ dw}$  (0.57-1.58) and individuals  $>100\text{cm}$  had  $2.12\pm0.96 \text{ mg}\cdot\text{kg}^{-1} \text{ dw}$  (1.50-4.04).




Figure 3 - Mercury concentration ( $\text{mg}\cdot\text{kg}^{-1}$  dry weight) per fork length of each individual and species, Bigeye tuna *Thunnus obesus*, Skipjack tuna *Katsuwonus pelamis*, Yellowmouth barracuda *Sphyraena viridensis* and Longfin yellowtail *Seriola rivoliana*

Table 2 - Mercury concentrations (Hg; wet weight (ww); dry weight (dw) when mentioned) of tunas, yellowtails and barracudas worldwide. The wet-to-dry weight ratios were estimated as 1 mg·kg<sup>-1</sup> wet weight to 3.3 mg·kg<sup>-1</sup> dw (Houssard et al. 2019). Values are represented as Mean±SD (Min-Max), when possible

Species	Region	N	Tissue	Length (cm)	Weight (kg)	Hg (mg·kg <sup>-1</sup> )	Reference
<i>Bigeye tuna</i> <i>Thunnus obesus</i>	Madeira	20	Blood	97.6±35.8 (66-170)	25.2±25.6 (7.7-86.0)	1.44±0.76 dw (0.57-4.04)	This study
	Madeira	20	Muscle	97.6±35.8 (66-170)	25.2±25.6 (7.7-86.0)	1.76±1.29 dw (0.74-5.53)	This study
	Atlantic	30	Muscle	(60-167)		0.76 (0.34-1.29)	Besada et al. 2006
	NE Brazil	30	Muscle	(54-139)		0.545±0.339 (0.09-1.75)	Lacerda et al. 2017
	Azores	15	Muscle		10.6±0.8(SE)	0.14±0.08	Torres et al. 2016
	Atlantic	7	Muscle		38.9 (35-43)	0.27±0.01	Yamashita et al. 2005
	Atlantic	121	Muscle	(70-200)	56±33	0.89±0.47 (0.32-3.13)	Chen et al. 2011
	Pacific	30	Muscle		59.4 (41-99)	0.98±0.34	Yamashita et al. 2005
	Hawaii	50	Muscle	92±5 (84-98)		0.43±0.20 (0.15-1.01)	Ferriss and Essington 2011
	Taiwan	75	Muscle	140.8±27.4 (81-200)	53.4±28.1 (9-140)	0.93±0.67 (0.16-3.32)	Chen et al. 2014
	N Pacific	6	Muscle	113.7±33.5 (75-164)		0.54±0.21 (0.20-0.76)	Blum et al. 2013
	Hawaii	50	Muscle		41.2±20.4 (11.3-89.8)	0.6±0.3	Kaneko and Ralston 2007

Skipjack tuna <i>Katsuwonus pelamis</i>	W Pacific	726	Muscle	91±31 (31-175)		0.43±0.48 (0.005-2.865)	Houssard et al. 2019
	Indian	27	Muscle	87±46		0.34±0.29	Bodin et al. 2017
	W Indian	110	Muscle	130±31 (53-198)	41±27 (1-110)	0.68±0.23 (0.22-1.88)	Chen et al. 2011
	Madeira	18	Blood	49.9±3.4 (43-58)	2.6±0.7 (1.8-4.2)	0.23±0.09 <b>dw</b> (0.12-0.49)	This study
	Azores	53	Muscle	49.7 (28-84)		0.19 (0.09-0.34)	Andersen and Depledge 1997
	Azores	15	Muscle		3.3±0.2(SE)	0.04±0.01	Torres et al. 2016
	Canary Islands	132	Muscle			0.34 (0.15, min)	Armas et al. 1993
	Central N Pacific	10	Muscle		8.6±1.3 (6.4-10.4)	0.3±0.1	Kaneko and Ralston 2007
	N Pacific	6	Muscle	65.7±16.03 (35-80)		0.28±0.13 (0.06-0.41)	Blum et al. 2013
	Reunion Island	37	Muscle	68 (41-85)	9 (1-16)	0.67±0.26 <b>dw</b>	Kojadinovic et al. 2007
Longfin yellowtail <i>Seriola rivoliana</i>	W Indian	13	Muscle	57±10		0.21±0.11	Bodin et al. 2017
	Madeira	20	Muscle	43.8±1.6 (41-48)	1.6±0.2 (1.3-2.2)	0.26±0.05 <b>dw</b> (0.17-0.34)	This study
	Florida	17	Muscle	74.3 (43.8-84.0)		0.56 (0.1-1.4)	Adams et al. 2003
<i>Seriola rivoliana</i> + <i>Seriola dumerili</i>	Bermuda	17	Muscle		(ca. 0.6-4.7)	0.17±0.14 (0.03-0.53)	Dewailly et al. 2008
	Hawaii	8	Muscle	77.9±2.9		0.76±0.07 (0.51-1.12)	Sackett et al. 2015

<i>Seriola dorsalis</i>	California	72	Muscle	84.2±16.6 (50.7-120.7)		1.27±0.89 <b>dw</b>	Madigan et al. 2018
Yellowmouth barracuda <i>Sphyraena viridensis</i>	Madeira	8	Muscle	63±2.7 (59-67)	1.2±0.2 (1.1-1.6)	1.09±0.20 <b>dw</b> (0.82-1.4)	This study
<i>Sphyraena barracuda</i>	Florida	13	Muscle	109.5±12.98		1.71±0.88 (3.4, max)	Rumbold et al. 2018
	Florida	62	Muscle	62.8 (11.9-109.6)		0.87 (0.08-3.10)	Adams et al. 2003
<i>Sphyraena sphyraena</i>	Mediterranean	5	Muscle	(24.9-31.0)	(55.6-107.3)	0.21±0.12 (0.15-0.42)	Hornung et al. 1980



## Discussion

The use of stomach contents allowed us to document the diet of the two most fished tuna species, the Bigeye and the Skipjack tuna, in the waters around the archipelago of Madeira. Here, both species presented a diet based on epipelagic organisms, and in the case of the Bigeye tuna complemented by some mesopelagic prey.

### *Diet of the Bigeye and Skipjack tuna*

The diet of the Bigeye tuna captured in the archipelago of Madeira consists mostly of epipelagic fish prey, in contrast with what was found in other regions in the Atlantic and Pacific. The published evidence suggests that in both oceans, its diet consists mainly of mesopelagic species, both fish and cephalopods, such as Bramidae, Alepisauridae, Myctophidae, Gempylidae, Paralepididae (eg. Matthews et al. 1977; Moteki et al. 2001; Allain 2005; Young et al. 2010; Junior et al. 2012) and Ommastrephidae (Logan et al. 2013). These families were very scarcely represented, or entirely absent in our study. Besides our study, where Scombridae and Carangidae were the main prey items, only one other study performed in the equatorial eastern Pacific in a pole-and-line fishery, found the Bigeye tuna to mostly prey on an epipelagic species (the Buccaneer anchovy *Stolephorus buccaneeri*) (Hida 1973). In this region, the bait consisted mostly of 4 epipelagic forage fish which were unimportant as prey items (Hida 1973).

Bigeye tunas are known to perform daily vertical migrations, descending on average up to 500m during the day and ascending to the surface at night, to forage (Matsumoto et al. 2005). However, a study carried out in the Azores showed that the Bigeye tunas in this archipelago do not perform vertical migrations as deep as elsewhere (Arrizabalaga et al. 2008). In that region, tunas remained mostly in the upper 50m layer, rarely venturing beyond the 300m depth. Arrizabalaga et al. (2008) hypothesized that oceanographic features in the Azores would enhance primary production and concentrate tuna food resources in surface layers, allowing Bigeye tuna to forage on shallower waters. This may suggest that, like in Madeira, this tuna concentrates on epipelagic prey in the Azores. Alternatively, it could be that in the Azores mesopelagic prey are more accessible at shallow depths, which is strongly suggested by their prominence in the diet of surface foragers (Monteiro et al. 1996; Granadeiro et al. 2002). To our knowledge, there are no diet studies of Bigeye tuna performed in the Azores which could confirm or contradict any of these hypotheses.

Some studies have suggested that there is an ontogenetic variation in the diet of several species of tunas (eg. Graham et al. 2007; Lacerda et al. 2017), including the Bigeye tuna which feeds on a higher proportion of mesopelagic species when larger due to its higher capacity to reach greater depths (Ohshimo et al. 2018). However, in our study, we could find no differences in the diet of smaller and larger size-classes.

The diet of the Skipjack tuna in the archipelago of Madeira is similar to what has been documented in other regions of the Atlantic, with a predominance of Clupeidae, Scombridae, Carangidae and Serranidae (Postel 1955; Batts 1972; Dragovich 1970; Dragovich and Potthoff 1972). Around the Canary Islands, where pole-and-line technique with live bait (sardine and sand smelt) is also used, the diet of this species is composed mainly of Atlantic chub mackerel (Ramos et al. 1995). In the Pacific, families like Exocoetidae, Engraulidae and reef fish are also part of the skipjack's diet (Alverson 1963; Hida 1973; Allain 2005). In smaller proportions, mesopelagic species, such as Gonostomatidae, Gempylidae and Myctophidae are also part of its diet in the Atlantic and the Pacific oceans (Ahlstrom and Counts 1958; Dragovich 1970; Ankenbrandt 1985). Still, its diet is mostly epipelagic. In our study the only mesopelagic prey found was a Trichiuridae. Crustaceans, such as euphausiids (Ankenbrandt 1985), stomatopods and megalopa larvae, completed the diet of this tuna in several regions of the Atlantic and Pacific (Alverson 1963; Dragovich 1970; Batts 1972; Dragovich and Potthoff 1972; Bernard et al. 1985). However, in Madeira, the contribution of crustaceans was negligible.

As opportunistic species, tunas are known to shift their diet among foraging regions but also when prey community shifts (Olson et al. 2014). The increase of Longspine snipefish in the diet of the Bigeye and Skipjack tuna follows the trend already observed in the diet of the Cory's shearwaters *Calonectris borealis* in the same oceanic region (J. Romero unpubl. data), where the authors suggested that a shift happened in the pelagic community of the subtropical northeast Atlantic with an increase of the Longspine snipefish population from 2016 to 2017/2018.

#### *The importance of bait in the diet of tunas*

In this study, the characterization of the diet of the Bigeye and the Skipjack tunas considered the issues posed by the use of bait during the fishing events. The two predators which were used for comparison, the Yellowmouth barracuda and the Longfin yellowtail, feed on a

variety of schooling epipelagic species, such as engraulids, clupeids, scombrids, carangids and sparids (Allam et al. 1999; Barreiros et al. 2002, 2003; Kalogirou et al. 2012; Manooch and Haimovici 1983). It is worth mentioning that in the Azores, the main diet of barracudas and yellowtails are Blue jack mackerel and Atlantic chub mackerel (Barreiros et al. 2002, 2003). In the archipelago of Madeira, the yellowfin feeds on European pilchards and Blue jack mackerels (Cavaleiro et al. 2018) while the barracuda feeds on Atlantic chub mackerels (M. Hermida unpubl. data). These species corresponded to the main tuna prey we observed in this study.

The exclusion of the least digested prey used as bait in Madeira (see eg. Ankenbrandt 1985; Ramos et al. 1995) had minimal impact in the assessment of the diet. Furthermore, highly digested remains of bait species were found in Skipjack tunas, strongly suggesting they are also caught as natural prey. Therefore, the decision to exclude bait species from the diet analysis (rather than discarding only freshly ingested prey) as done in some studies (eg. Ankenbrandt 1985; Ramos et al. 1995), may be excluding prey that are actually important in the diet of these predators.

The analysis of mercury revealed two different groups of predators, one with higher concentrations (Bigeye tuna and Yellowmouth barracuda) and one with lower concentrations (Skipjack tuna and the Longfin yellowtail). Bigeye tuna showed slightly higher Hg values than the other predators, most likely because the diet of the former includes mesopelagic species, which have high concentrations of mercury (e.g. Monteiro et al. 1996). This is further supported by Choy et al. (2009) which concluded that predators with a mesopelagic diet have higher concentrations of mercury than those with an epipelagic diet. The barracuda also presented higher mercury values despite reportedly having an epipelagic diet. The reasons for this are still unknown, but the fact that Hg levels in Bigeye tuna are only slightly higher than in barracuda further supports the conclusion that mesopelagics are not an important component of the tuna's diet, or else we would expect a larger difference between these species.

Overall, the mercury concentrations in tissues of predators presented in this study are lower than in others from the Atlantic, Pacific, and Indian oceans (Table 2). While intraspecific differences in mercury burdens have also been attributed to the availability of mercury in different foraging regions (Peterson et al. 1973; Ferriss and Essington 2011;

Houssard et al. 2019), the use of conventional diet in our study, indicates that the differences in relation to other areas might be better explained by an epipelagic diet of the Bigeye tuna. Other variables, such as elimination and uptake rate of mercury, may explain the variations observed among species (Peterson et al. 1973). According to Ferriss and Essington (2014), the Skipjack tuna has a higher elimination rate than the Bigeye tuna (0.376 and 0.077  $y^{-1}$ , respectively) which might explain the lower concentrations. Trudel and Rasmussen (1997) stated that the elimination rate responds negatively to the predator's body mass which could explain the differences between these two species. There are no elimination rates described for barracudas and yellowtails, remaining unclear what are the mechanisms driving the differences in mercury concentrations for barracudas and yellowtails.

The mercury values described in this study for the Bigeye, Skipjack, yellowtail and barracuda were obtained using blood for tunas and muscle tissue for the two other predators. When comparing the mercury values for blood and muscle for the Bigeye tuna, the calculated mean blood-muscle ratio was 1:1.17. This suggests that mercury levels in blood and muscle are broadly comparable, but more studies are needed to better assess that, as 3 (out of 20) of the individuals, which were also 3 of the biggest sampled tunas, presented higher mercury ratios.

The baseline information provided in this study raises new questions on the functioning of the ecosystem of this region. The Bigeye and the Skipjack tuna presented an epipelagic diet in the pelagic region of the archipelago of Madeira, which in the case of the Bigeye tuna suggests a different behaviour than the one often presented by this species. Further studies on the variability in oceanographic variables and its effects on the distribution and abundance of prey, as well as on the vertical movements of tuna in this area are required to understand their behaviour and explain why this mesopelagic predator presents an epipelagic diet in Madeira. The results of this study will allow for more informed decisions on an ecosystem-based fisheries management by considering not only the tuna but also the role they may have in the regional food webs and their interactions with other species.

## **Acknowledgements**

This work was supported by Fundação para a Ciência e a Tecnologia (FCT Portugal) through the project Oceanwebs (PTDC/MAR-PRO/0929/2014) and Seamigrant (PTDC/BIA-ANM/3743/2014), the strategic projects UIDB/04292/2020 and

UIDP/04292/2020 granted to MARE, and UIDB/50017/2020 and UIDP/50017/2020 granted to CESAM. PD/BD/127810/2016 was granted to JR by FCT. MH was supported by a postdoctoral grant from ARDITI (Regional Agency for the Development of Research, Technology and Innovation of Madeira), Project M1420-09-5369-FSE-000001. BC was supported by a research grant within the project OceanWebs (PTDC/MAR-PRO/0929/2014). Bigeye tuna and barracudas used in this study were acquired by ARDITI through the Oceanic Observatory of Madeira Project (M1420-01-0145-FEDER-000001-Observatório Oceânico da Madeira-OOM). Skipjack tuna were acquired by the Regional Directorate of Fisheries (DRP) of Madeira, under the framework of the Programa Nacional de Recolha de Dados da Pesca (multiannual Union programme for the collection, management and use of data in the fisheries and aquaculture sectors). The authors are grateful to the Regional Directorate of Fisheries (DRP) of Madeira for their collaboration and support, particularly the Research Service director, João Delgado. Thanks are also due to Filipe Moniz and Carolina Vieira for helping sieving and identifying some of the stomach contents, and Cheila Raposo and Carolina Vieira for performing the mercury analysis.

## References

- Adams DH, McMichael Jr. RH, Henderson GE (2003) Mercury levels in marine and estuarine fishes of Florida 1989–2001. Florida Marine Research Institute Technical Report TR-9. 2nd ed. rev. 57 pp.
- Ahlstrom EH, Counts RC (1958) Development and distributions of *Vinciguerria lucetia* and related species in the eastern Pacific. Fish Bull 58:363–416.
- Allain V (2005) Diet of large pelagic predators of the Western and Central Pacific Ocean. Noumea, New Caledonia. 18 pp.
- Allam SM, Faltas SN, Ragheb E (1999) Food and feeding habits of barracudas in the Egyptian Mediterranean water off Alexandria. Bull Nat Inst Ocean Fish 25:395–410.
- Aloncle H, Delaporte F (1974) Données nouvelles sur le germon *Thunnus alalunga* Bonnaterre 1788 dans le nord-est Atlantique. Rev Trav Inst Pêches marit 37(4):475–578.
- Alonso H, Granadeiro JP, Dias MP, Catty T, Catty P (2018) Fine-scale tracking and diet information of a marine predator reveals the origin and contrasting spatial distribution of prey. Prog Oceanogr 162:1–12. <https://doi.org/10.1016/j.pocean.2018.02.014>.
- Alonso H, Granadeiro JP, Waap S, Xavier J, Symondson WOC, Ramos JA, Catty P (2014)

- An holistic ecological analysis of the diet of Cory's shearwaters using prey morphological characters and DNA barcoding. *Mol Ecol* 23:3719–3733. <https://doi.org/10.1111/mec.12785>
- Alverson FG (1963) The Food of Yellowfin and Skipjack Tunas in the Eastern Tropical Pacific Ocean. *Inter-american Trop tuna Comm* 7:295–396.
- Andersen JL, Depledge MH (1997) A Survey of Total Mercury and Methylmercury in Edible Fish and Invertebrates from Azorean Waters. *Mar Environ Res* 44:331–350.
- Ankenbrandt L (1985) Food habits of bait-caught skipjack tuna, *Katsuwonus pelamis*, from the southwestern Atlantic Ocean. *Fish Bull* 83:379–393.
- Armas JD, Diaz J, Galan JM (1993) Mercury content of the skipjack, *Katsuwonus pelamis*, off the Canary Islands and adjacent waters. *Bol del Inst Esp Oceanogr* 9:9–14.
- Arrizabalaga H, Pereira JG, Royer F, Galuardi B, Goñi N, Artetxe I, Arregi I, Lutcavage M (2008) Bigeye tuna (*Thunnus obesus*) vertical movements in the Azores Islands determined with pop-up satellite archival tags. *Fish Oceanogr* 17:74–83.
- Barreiros JP, Santos RS, Borba AE (2002) Food habits, schooling and predatory behaviour of the yellowmouth barracuda, *Sphyrna viridensis* (Perciformes: Sphyrnaidae) in the Azores. *Cybiu* 26:83–88.
- Barreiros JP, Morato T, Santos RS, Borba AE (2003) Interannual changes in the diet of the almaco jack, *Seriola rivoliana*, (Perciformes: Carangidae) from the Azores. *Cybiu*, 27: 37–40.
- Batts BS (1972) Food habits of the skipjack tuna, *Katsuwonus pelamis*, in North Carolina waters. *Chesap Sci* 13:193–200.
- Bernard HJ, Hedgepeth JB, Reilly SB (1985) Stomach contents of albacore, skipjack, and bonito caught off southern California during summer 1983. *CalCOFI Rep* 26:175–182.
- Besada V, González JJ, Schultze F (2006) Concentraciones de mercurio, cadmio, plomo, arsénico, cobre y zinc en atún blanco, rabil y patudo procedentes del Océano Atlántico. *Ciencias Mar* 32:439–445.
- Blum JD, Popp BN, Drazen JC, Anela Choy C, Johnson MW (2013) Methylmercury production below the mixed layer in the North Pacific Ocean. *Nat Geosci* 6: 879–884. <http://dx.doi.org/10.1038/ngeo1918>.
- Bodin N, Lesperance D, Albert R, Hollanda S, Michaud P, Degroote M, Churlaud C,

- Bustamante P (2017) Chemosphere Trace elements in oceanic pelagic communities in the western Indian Ocean. *Chemosphere* 174:354–362. <http://dx.doi.org/10.1016/j.chemosphere.2017.01.099>.
- Cavaleiro B, Hermida M, Saraiva A (2018) Parasites of amberjacks from the archipelago of Madeira, Eastern Atlantic. *Dis Aquat Organ* 131:133–142.
- Chen CY, Lai CC, Chen KS, Hsu CC, Hung CC, Chen MH (2014) Total and organic mercury concentrations in the muscles of Pacific albacore (*Thunnus alalunga*) and bigeye tuna (*Thunnus obesus*). *Mar Pollut Bull* 85:606–612. <http://dx.doi.org/10.1016/j.marpolbul.2014.01.039>.
- Chen MH, Teng PY, Chen CY, Hsu CC, Teng PY, Chen CY, Organic CCH (2011) Organic and total mercury levels in bigeye tuna, *Thunnus obesus*, harvested by Taiwanese fishing vessels in the Atlantic and Indian Oceans. 4(1): 15-21. <https://doi.org/10.1080/19393210.2010.535908>
- Choy CA, Popp BN, Kaneko JJ, Drazen JC (2009) The influence of depth on mercury levels in pelagic fishes and their prey. *Proc Natl Acad Sci* 106:13865–13869. <https://doi.org/10.1073/pnas.0900711106>.
- Clarke, M. R. 1986. A handbook for the identification of cephalopod beaks. Clarendon Press, Oxford.
- da Silva, G. B., Hazin, H. G., Hazin, F. H. V., and Vaske-Jr, T. 2019. Diet composition of bigeye tuna (*Thunnus obesus*) and yellowfin tuna (*Thunnus albacares*) caught on aggregated schools in the western equatorial Atlantic Ocean. *J Appl Ichthyol* 35: 1111–1118.
- Dewailly É, Rouja P, Dallaire R, Pereg D, Tucker T, Ward J, Weber JP, Maguire JS, Julien P (2008) Balancing the risks and the benefits of local fish consumption in Bermuda. *Food Addit Contam - Part A Chem Anal Control Expo Risk Assess* 25:1328–1338. <https://doi.org/10.1080/02652030802175285>
- Doney SC, Ruckelshaus M, Emmett Duffy J, Barry JP, Chan F, English CA, Galindo HM, Grebmeier JM, Hollowed AB, Knowlton N, Polovina J, Rabalais NN, Sydeman WJ, Talley LD (2012) Climate Change Impacts on Marine Ecosystems. *Ann Rev Mar Sci* 4:11–37. <https://doi.org/10.1146/annurev-marine-041911-111611>.
- Dragovich A (1970) The food of Skipjack and Yellowfin tunas in the Atlantic Ocean. *Fish*

Bull 68:445–460

- Dragovich A, Potthoff T (1972) Comparative study of food of skipjack and yellowfin tunas off the coast of west africa. Fish Bull 70:1087–1110.
- Ferriss BE, Essington TE (2011) Regional patterns in mercury and selenium concentrations of yellowfin tuna (*Thunnus albacares*) and bigeye tuna (*Thunnus obesus*) in the Pacific Ocean. 2056:2046–2056. <http://dx.doi.org/10.1139/F2011-120>.
- Ferriss BE, Essington TE (2014) Does trophic structure dictate mercury concentrations in top predators? A comparative analysis of pelagic food webs in the Pacific Ocean. Ecol Modell 278:18–28. <http://dx.doi.org/10.1016/j.ecolmodel.2014.01.029>.
- Fontes J, Afonso P (2017) Long-term residency and movements of yellowmouth barracuda (*Sphyræna viridensis*) at a shallow seamount. Mar Biol 164:1–10. <http://doi.org/10.1007/s00227-016-3045-1>.
- Furtado R, Pereira ME, Granadeiro JP, Catry P (2019) Body feather mercury and arsenic concentrations in five species of seabirds from the Falkland Islands. Mar Pollut Bull 149:110574. <https://doi.org/10.1016/j.marpolbul.2019.110574>.
- Glaser SM, Waechter KE, Bransome NC (2015) Through the stomach of a predator: Regional patterns of forage in the diet of albacore tuna in the California Current System and metrics needed for ecosystem-based management. J Mar Syst 146:38–49. <http://doi.org/10.1016/j.jmarsys.2014.07.019>.
- Gorni GR (2016) Description of diet of pelagic fish in the southwestern Atlantic, Brazil. Biota Neotrop, 13(1). <https://doi.org/10.1590/S1676-06032013000100006>.
- Gouveia L, Amorim A, Alves A, Hermida M (2019) Updated fishery statistics of tuna species caught off Madeira archipelago. Collect Vol Sci Pap ICCAT 75:2081–2097.
- Gouveia L, Mejuto J (2003) Seasonality and interannual variability in catches of skipjack tuna (*Katsuwonus pelamis*) and bigeye tuna (*Thunnus obesus*) in the area around the Archipelago of Madeira. Col Vol Sci Pap ICCAT 55:1853–1867.
- Graham BS, Grubbs D, Holland K, Popp BN (2007) A rapid ontogenetic shift in the diet of juvenile yellowfin tuna from Hawaii. Mar Biol 150:647–658. <http://doi.org/10.1007/s00227-006-0360-y>
- Granadeiro JP, Monteiro LR, Silva MC, Furness RW (2002) Diet of Common Terns in the Azores, Northeast Atlantic. Waterbirds 25:149–155. [66](http://dx.doi.org/10.1675/1524-</a></p></div><div data-bbox=)



- 4695(2002)025[0149:DOCTIT]2.0.CO;2.
- Granadeiro JP, Silva MA (2000) The use of otoliths and vertebrae in the identification and size-estimation of fish in predator-prey studies. *Cybium* 24:383–393.
- Hermida M, Delgado J (2016) High trophic level and low diversity: Would Madeira benefit from fishing down? *Mar Policy* 73:130–137. <http://dx.doi.org/10.1016/j.marpol.2016.07.013i>.
- Hida T (1973) Food of Tunas and dolphins (Pisces: Scombridae and Coryphaenidae) with emphasis on the distribution and biology of their prey *Stolephorus buccaneeri* (Engraulidae). *Fish Bull US* 71:135–143.
- Hornung H, Zismann L, Oren OH (1980) Mercury in twelve Mediterranean trawl fishes of Israel. *Environ Int* 3:243–248.
- Houssard P, Point D, Tremblay-Boyer L, Allain V, Pethybridge H, Masbou J, Ferriss BE, Baya PA, Lagane C, Menkes CE, Letourneur Y, Lorrain A (2019) A Model of Mercury Distribution in Tuna from the Western and Central Pacific Ocean: Influence of Physiology, Ecology and Environmental Factors. *Environ Sci Technol* 53:1422–1431. <http://doi.org/10.1021/acs.est.8b06058>.
- Hunter J, Macewicz B, Sibert J (1986) the Spawning Frequency of Skipjack Tuna, *Katsuwonus Pelamis*, From the South Pacific. *Fish Bull* 84:895–903.
- ICCAT. 2020. International Commission for the Conservation of Atlantic Tunas. <https://www.iccat.int/en/accesingdb.html>. (accessed 23 March 2020)
- IUCN. 2020. IUCN Red List of Threatened Species. <https://www.iucnredlist.org/>. (accessed 17 February 2020).
- Junior T V., Travassos PE, Hazin FH V., Tolotti MT, Barbosa TM (2012) Forage fauna in the diet of Bigeye tuna (*Thunnus obesus*) in the Western tropical Atlantic Ocean. *Brazilian J Oceanogr* 60:89–97.
- Kalogirou S, Mittermayer F, Pihl L, Wennhage H (2012) Feeding ecology of indigenous and non-indigenous fish species within the family Sphyraenidae. *J Fish Biol* 80:2528–2548. <http://doi.org/10.1111/j.1095-8649.2012.03306.x>.
- Kaneko JJ, Ralston NVC (2007) Selenium and Mercury in Pelagic Fish in the Central North Pacific Near Hawaii. :242–254. <http://doi.org/10.1007/s12011-007-8004-8>
- Kojadinovic J, Potier M, Corre M Le, Cosson RP, Bustamante P (2007) Bioaccumulation of

- trace elements in pelagic fish from the Western Indian Ocean. *Environ Pollut* 146:548–566. <http://doi.org/10.1016/j.envpol.2006.07.015>
- Lacerda LD, Goyanna F, Bezerra MF, Silva GB (2017) Mercury Concentrations in Tuna (*Thunnus albacares* and *Thunnus obesus*) from the Brazilian Equatorial Atlantic Ocean. *Bull Environ Contam Toxicol* 98:149–155. <http://doi.org/10.1007/s00128-016-2007-0>
- Layman CA, Araujo MS, Boucek R, Hammerschlag-Peyer CM, Harrison E, Jud ZR, Matich P, Rosenblatt AE, Vaudo JJ, Yeager LA, Post DM, Bearhop S (2012) Applying stable isotopes to examine food-web structure: An overview of analytical tools. *Biol Rev* 87:545–562. <http://doi.org/10.1111/j.1469-185X.2011.00208.x>
- Lehodey P, Bertignac M, Hampton J, Lewis A, Picaut J (1997) El Niño Southern Oscillation and tuna in the western Pacific. *Nature* 389:715–718.
- Logan JM, Toppin R, Smith S, Galuardi B, Porter J, Lutcavage M (2013) Contribution of cephalopod prey to the diet of large pelagic fish predators in the central North Atlantic Ocean. *Deep Res Part II Top Stud Oceanogr* 95:74–82. <http://dx.doi.org/10.1016/j.dsr2.2012.06.003>.
- Lu CC, Ickeringill R (2002) Cephalopod beak identification and biomass estimation techniques: tools for dietary studies of southern Australian finfishes. *Museum Victoria Sci Reports* 6:1–65. <https://doi.org/10.24199/j.mvsvr.2002.06>
- Madigan DJ, Snodgrass OE, Fisher NS (2018) From migrants to mossbacks: Tracer- and tag-inferred habitat shifts in the California yellowtail *Seriola dorsalis*. *Mar Ecol Prog Ser* 597:221–230. <https://doi.org/10.3354/meps12593>
- Manooch CSI, Haimovici M (1983) Foods of greater amberjack, *Seriola dumerili*, and Almaco Jack, *Seriola rivoliana* (Pisces: Carangidae), from the South Atlantic Bight. *J Elisha Mitchell Sci Soc* 99:1–9.
- Matsumoto T, Miyabe N (2002) Preliminary report on the maturity and spawning of bigeye tuna *Thunnus obesus* in the central Atlantic Ocean. *ICCAT Collect Vol Sci Pap* 54:246–260.
- Matsumoto T, Saito H, Miyabe N (2005) Swimming Behavior of Adult Bigeye Tuna Using Pop-Up Tags. *Col Vol Sci Pap ICCAT* 57:151–170.
- Matthews FD, Damkaer DM, Knapp LW, Collette BB (1977) Food of Western North Atlantic Tunas (*Thunnus*) and Lancetfish (*Alepisaurus*). NOAA Technical Reports

- NMFS SSRF-7061: 19 pp.
- McClellan DB, Cummings NJ (1997) Preliminary Analysis of Tag and Recapture Data of the Greater Amberjack, *Seriola dumerili*, in the Southeastern United States. Proc Gulf Caribb Fish Inst 49.
- Ménard F, Labruno C, Shin YJ, Asine AS, Bard FX (2006) Opportunistic predation in tuna: A size-based approach. Mar Ecol Prog Ser 323:223–231. <https://doi.org/10.3354/meps323223>
- Monteiro LR, Ramos JA, Furness RW, Delnevo AJ (1996) Movements, morphology, breeding, molt, diet and feeding of seabirds in the azores. Colon Waterbirds 19:82–97
- Moteki M, Arai M, Tsuchiya K, Okamoto H (2001) Composition of piscine prey in the diet of large pelagic fish E tropic Pacific Ocean. Fish Sci 67:1063–1074
- Myers RA, Worm B (2003) Rapid worldwide depletion of predatory fish communities. Nature 423:280–283. <http://www.nature.com/doi/10.1038/nature01610>.
- Ohshimo S, Hiraoka Y, Sato T, Nakatsuka S (2018) Feeding habits of bigeye tuna (*Thunnus obesus*) in the North Pacific from 2011 to 2013. Mar Freshw Res 69:585–606. <https://doi.org/10.1071/MF17058>
- Olson RJ, Duffy LM, Kuhnert PM, Galván-Magaña F, Bocanegra-Castillo N, Alatorre-Ramírez V (2014) Decadal diet shift in yellowfin tuna *Thunnus albacares* suggests broad-scale food web changes in the eastern tropical Pacific Ocean. Mar Ecol Prog Ser 497:157–178. <https://doi.org/10.3354/meps10609>
- Palumbi SR (1996) Nucleic acids II: the polymerase chain reaction. In: Molecular Systematics, Hillis DM, p 205–247.
- Peterson CL, Klawe WL, Sharp GD (1973) Mercury in Tunas: A Review. Fish Bull 71:603–613.
- Ramos AG, Lorenzo JM, Pajuelo JG (1995) Food habits of bait-caught skipjack tuna, *Katsuwonus pelamis*, off the Canary Islands. Sci Mar 59:365–369.
- Reynolds SJ, Hughes BJ, Wearn CP, Dickey RC, Brown J, Weber NL, Weber SB, Paiva VH, Ramos JA (2019) Long-term dietary shift and population decline of a pelagic seabird—A health check on the tropical Atlantic? Glob Chang Biol 25:1383–1394. <https://doi.org/10.1111/gcb.14560>
- Rumbold DG, Lienhardt CT, Parsons ML (2018) Mercury Biomagnification Through a Coral

- Reef Ecosystem. Arch Environ Contam Toxicol 75:121–133. <https://doi.org/10.1007/s00244-018-0523-0>.
- Sackett DK, Drazen JC, Choy CA, Popp B, Pitz GL (2015) Mercury sources and trophic ecology for Hawaiian bottomfish. Environ Sci Technol 49:6909–6918. <https://doi.org/10.1021/acs.est.5b01009>
- Schaefer KM, Fuller DW, Block BA (2009) Tagging and Tracking of Marine Animals with Electronic Devices. In: Tagging and Tracking of Marine Animals with Electronic Devices. p 121–144.
- Storelli MM, Giacomini-Stuffler R, Storelli A, Marcotrigiano GO (2005) Accumulation of mercury, cadmium, lead and arsenic in swordfish and bluefin tuna from the Mediterranean Sea: A comparative study. Mar Pollut Bull 50:1004–1007.
- Team RC (2020) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.r-project.org/>.
- Teffer AK, Staudinger MD, Taylor DL, Juanes F (2014) Trophic influences on mercury accumulation in top pelagic predators from offshore New England waters of the northwest atlantic ocean. Mar Environ Res 101:124–134. <http://dx.doi.org/10.1016/j.marenvres.2014.09.008>.
- Torres P, Rodrigues A, Soares L, Garcia P (2016) Metal Concentrations in Two Commercial Tuna Species from an Active Volcanic Region in the Mid-Atlantic Ocean. Arch Environ Contam Toxicol 70:341–347. <http://doi.org/10.3354/meps11707>
- Trudel M, Rasmussen JB (1997) Modeling the elimination of mercury by fish. Environ Sci Technol 31:1716–1722.
- Varela JL, Cañavate JP, Medina A, Mourente G (2019) Inter-regional variation in feeding patterns of skipjack tuna (*Katsuwonus pelamis*) inferred from stomach content, stable isotope and fatty acid analyses. Mar Environ Res 152.
- Voegborlo RB, Adimado AA, Ephraim JH (2007) Total mercury distribution in different tissues of frigate tuna (*Auxis thazard thazard*) from the atlantic coastal waters of Ghana, Gulf of Guinea. Environ Monit Assess 132:503–508. <http://doi.org/10.1007/s10661-006-9552-7>
- Whitehead PJP, Bauchot M-L, Hureau J-C, Nielsen J, Tortonese E (1989) Fishes of the North-eastern Atlantic and the Mediterranean. UNESCO, Paris.

- Yamashita Y, Omura Y, Okazaki E (2005) Total mercury and methylmercury levels in commercially important fishes in Japan. *Fish Sci* 71:1029–1035.
- Young JW, Lansdell MJ, Campbell RA, Cooper SP, Juanes F, Guest MA (2010) Feeding ecology and niche segregation in oceanic top predators off eastern Australia. *Mar Biol* 157:2347–2368. <http://doi.org/10.1007/s00227-010-1500-y>

## Supplementary Material

Table A.1 - Classification of the digestion level of tuna's prey for fishes and cephalopods according to Aloncle and Delaporte (1974) and Alonso et al. (2018), respectively.

Type of prey	Digestion Level	Description
Fish	I	The fish that was just swallowed, still presents all scales and original colour.
	II	The fish is whole but partly or entirely stripped of its scales and skin. Muscles are visible.
	III	The fish is split but the flesh is still attached to the skeleton.
	IV	The fish is only represented by skeleton debris.
Cephalopods	I	Cephalopod complete
	II	Mantle complete
	III	More than half of mantle present
	IV	Only a piece of mantle or isolated tentacles present
	V	Loose fresh beak (with flesh attached and/or not eroded)




Figure A.1 - Regression between mercury concentration (mg.kg<sup>-1</sup> dw) in blood and muscle of Bigeye tunas *Thunnus obesus* captured in the pelagic waters of the archipelago of Madeira.

Table A.2 - Cephalopod species present in the stomach contents of the Bigeye tuna *Thunnus obesus* and identified using their beaks. These beaks did not have any flesh attached which means that they were not ingested recently, and, therefore, were excluded from the diet characterization to avoid overestimation. NF – numeric frequency (sample size); FO – Frequency of occurrence (sample size).

Prey	Order/Family	Species	NF (140)	FO (29)
<b>Decapodiformes</b>				
	Mastigoteuthidae	<i>Mastigoteuthis</i> sp.	0.7	3.4
	Ommastrephidae	<i>Todarodes sagittatus</i>	4.3	10.3
		Unidentified	11.4	20.7
	Onychoteuthidae	<i>Onychoteuthis</i> sp.	10.0	10.3
	Brachioteuthidae	<i>Brachioteuthis risei</i>	1.4	6.9
	Chroteuthidae	<i>Planchoteuthis danae</i>	8.6	10.3
	Oegopsida	Unidentified	8.6	13.8
	Unidentified		17.9	24.1
<b>Octopodiformes</b>				
	Octopodidae	Unidentified	2.9	6.9
	Amphitretidae	<i>Japetella diaphana</i>	1.4	3.4
	Argonautidae	<i>Argonauta</i> sp.	4.3	10.3
	Argonautoidae (Superfamily)	Unidentified	0.7	3.4
<b>Unidentified</b>			<b>27.9</b>	<b>44.8</b>

#### A.1 - Bait

Removing prey with low digestion levels (I and II) from the analysis increased the percentage of empty stomachs from 3% to 32% (n=71) for the Bigeye tuna, and from 6% to 11% (n=61) for the Skipjack tuna. Still, no large shifts in the relative importance of prey were observed in the diet of these tunas, whether bait was included or not in the analysis (Fig. S2 and S3).




Figure A.2 - Diet of the Bigeye tuna *Thunnus obesus* (Numeric frequency - NF and Frequency of occurrence - FO) including all individuals found in the stomachs (n=69, Bait) and excluding bait species (Atlantic chub mackerel *Scomber colias*, mackerel *Trachurus sp.* and Bogue *Boops boops*) with digestion levels I and II (n=48, No bait).






Figure A.3 - Diet of the Skipjack tuna *Katsuwonus pelamis* (Numeric frequency - NF and Frequency of occurrence - FO) including all individuals found in the stomachs (n=57, Bait) and excluding bait species (Sand smelt *Atherina sp.*, Bogue *Boops boops*, European pilchard *Sardina pilchardus*, and Longspine snipefish *Macroramphosus scolopax*) with digestion levels I and II (n=54, No Bait).

Table A.3 - Stomach contents of the Skipjack tuna *Katsuwonus pelamis* caught in Cardume in 2018 in the archipelago of Madeira and bait used in each fishing event

Bait	Stomach ID	Content	Number	Digestion level
<i>Sardina pilchardus</i> <i>Atherina</i> sp.	1	<i>Boops boops</i>	3	2-3
		<i>Atherina</i> sp.	1	4
	2	<i>Macroramphosus scolopax</i>	6	3-4
		<i>Atherina</i> sp.	1	4
	3	<i>Scomber colias</i>	2	3
		<i>Macroramphosus scolopax</i>	2	2-4
		Clupeidae	1	4
		<i>Atherina</i> sp.	2	4
	4	<i>Atherina</i> sp.	3	2
		<i>Macroramphosus scolopax</i>	24	3-4
	5	<i>Macroramphosus scolopax</i>	11	3-4
	6	<i>Atherina</i> sp.	5	2-4
		<i>Macroramphosus scolopax</i>	7	3-4
	7	<i>Atherina</i> sp.	8	2-4
		<i>Macroramphosus scolopax</i>	18	3-4
	8	<i>Atherina</i> sp.	4	2-3
		<i>Macroramphosus scolopax</i>	14	3-4
<i>Sardina pilchardus</i> <i>Boops boops</i> <i>Atherina</i> sp.	9	<i>Macroramphosus scolopax</i>	2	4
	10	<i>Boops boops</i>	3	1
	11	<i>Boops boops</i>	1	4
	12	<i>Boops boops</i>	1	4
	13	<i>Scomber colias</i>	1	4

## References

- Aloncle H, Delaporte F (1974) Données nouvelles sur le germon *Thunnus alalunga* Bonnaterre 1788 dans le nord-est Atlantique. Rev Trav Inst Pêches marit 37(4):475-578.
- Alonso H, Granadeiro JP, Dias MP, Catry T, Catry P (2018) Fine-scale tracking and diet information of a marine predator reveals the origin and contrasting spatial distribution of prey. Prog Oceanogr 162:1–12. <https://doi.org/10.1016/j.pocean.2018.02.014>.

## CHAPTER 4

---



Credits: Joana Romero

*This chapter is published as:*

*Joana Romero, Paulo Catry, Dília Menezes, Nádia Coelho, João Paulo Silva, José Pedro Granadeiro (2019) A gull that scarcely ventures on the ocean: Yellow-legged gulls *Larus michahellis atlantis* on the oceanic island of Madeira. *Ardeola* 66: 101-112.*

# **A gull that scarcely ventures on the ocean: Yellow-legged Gulls *Larus michahellis atlantis* on the oceanic island of Madeira**

## **Abstract**

The Yellow-legged Gull *Larus michahellis atlantis* (YLG) is a generalist and opportunistic species that explores different habitats and resources and easily adapts to humanised environments, sometimes posing problems related to aircraft security, public health or predation on other species, for example. We examined the distribution and foraging behaviour of YLGs on the oceanic island of Madeira. GPS-GSM devices were deployed on ten breeding individuals and samples of blood (adults) and feathers (chicks) were collected to examine their stable isotope signatures in relation to those of their prey. Tracked YLGs did not migrate. Birds remained mostly inland during the day (up to 60% of fixes) and barely 5% of the fixes were at sea. Eighty-eight out of 192 gull trips to the sea were in interaction with local fishing vessels, 91% of which involving purse seiners that were active at night. The diet of the adult YLGs consisted of refuse, fish and small petrels, whereas the chick diet included mainly fish. YLGs on this oceanic island are strongly dependent on anthropogenic resources, scarcely relying on natural marine prey. The desired control of this gull population could perhaps be achieved by the reduction of accessibility to organic waste.

**Keywords:** Fishing activity, GPS tracking, Habitat-use, Human waste, Stable isotopes

## Introduction

Populations of large gulls, including Yellow-legged Gulls *Larus michahellis atlantis* (YLG) have been increasing continuously over the past century (eg. Harris, 1970; Duhem *et al.*, 2008). Their generalist and opportunistic behaviour allows them to exploit different environments and resources within both natural and human-modified areas (Moreno *et al.*, 2010; Navarro *et al.*, 2017). Overall, YLGs can be found in both terrestrial and marine environments, the latter mostly used to forage (Arizaga *et al.*, 2010; Ceia *et al.*, 2014), often in association with human activities (*e.g.* at harbours, fishing vessels). Their marine-based diet comes to some extent from natural prey but in many areas is primarily obtained from fishing discards (Oro *et al.*, 1995; Ceia *et al.*, 2014). Interactions between gulls and fishing vessels are a well-known and widespread phenomenon, mainly involving trawlers due to their high discard rates (Camphuysen *et al.*, 1993; Garthe *et al.*, 1996; Karris *et al.*, 2018). Refuse dumps are also important reliable locations for gulls to obtain food (Duhem *et al.*, 2003; Arizaga *et al.*, 2014; Navarro *et al.*, 2016). The plastic behaviour showed by YLGs has led to overpopulation in many areas, which has turned this species into an important safety and public health hazard (Ramos *et al.*, 2009b; Ramos *et al.*, 2010). Its predatory behaviour also poses challenges to the conservation of rarer species (Hernández-Matías & Ruiz, 2003; Matias & Catry, 2010).

YLG populations at mainland coasts and estuaries are well studied but there is comparatively little information regarding their distribution and behaviour on oceanic islands, other than some dietary information (eg. Matias & Catry, 2010; Pedro *et al.*, 2013; López *et al.*, 2016). We examined the foraging behaviour and distribution of YLGs breeding on the Portuguese archipelago of Madeira, a group of oceanic Atlantic islands surrounded by oligotrophic waters. In particular, we aimed to investigate the distribution, habitat-use and daily activity patterns of YLGs, to quantify the relationship between gulls and fishing vessels and to assess the importance of natural prey in their diet. There is no trawler fleet in Madeira (Vallerani *et al.*, 2017), which leads us to hypothesise that the Madeiran YLG population is less dependent on marine resources despite inhabiting an environment surrounded by sea.

## Methods

### *Study area*

The YLG has three main breeding colonies on Madeira island (32°40'08.97'' N, 16°55'28.06'' W), on the uninhabited islets of Desembarcadouro and Chão and on small islets close to Porto Santo island. This work was carried out on Desembarcadouro islet (46ha), which is almost contiguous with Madeira, and on Chão islet (43ha), c. 20km SE of Madeira. The breeding populations of Desembarcadouro and Chão islets have been estimated at about 1,900 and 600 breeding pairs, respectively.

### *GPS-tracking and gull distribution*

In late April 2016 and 2017, six GPS-GSM devices were deployed on incubating birds on Chão islet, and three on Desembarcadouro islet, respectively. A tenth device was deployed only in late May 2017, on Desembarcadouro islet. All birds were marked with a metal ring, weighed and had devices deployed on their backs with a wing teflon harness (Thaxter *et al.*, 2014). The sex of the birds was unknown. The GPS-GSM devices were solar powered and measured 57x26x20mm and weighed 27g (Movetech Telemetry, maximum error ca. 100m), corresponding to c. 3% of the bird weight ( $870 \pm 106$ g, n=15 birds), and recorded fixes every 30 minutes under a normal regime and every two minutes if the battery level was very high. In parallel, we carried out point-counts in January 2017 at strategic sites, to assess the distribution of the YLG around the entire island of Madeira (Supplementary material appendix 1, A1). All procedures were approved by the Animal Welfare Body of the University of Lisbon (2/2016, 8<sup>th</sup> March 2016) and by National Authorities (0.421/2016, 14<sup>th</sup> November 2016).

The distribution of birds in the study area was described using a kernel density map on projected coordinates, using the *adehabitatHR* package (h=0.018, grid=300; Calenge, 2017), under R software (R Core Team, version 1.0.143) (Supplementary material appendix 1, A2).

### *Assessing gull activity in relation to fisheries*

We characterised the activity area of the local fishing fleet using data from the Terrestrial Automatic Identification System (AIS-T, obtained from [www.marinetraffic.com](http://www.marinetraffic.com)) complemented by data from the Vessel Monitoring System (VMS) collected for the study area (provided by the DGRM, see acknowledgements), covering April to September 2016

(Supplementary material appendix 1, A2). This period coincided with the highest number of gulls with at-sea positions.

To quantify the relative utilisation of used areas, four different habitat categories were defined: land (golf fields, urban solid waste treatment plant, wind farm fields, open fields), harbours (at Funchal and Caniçal), coastline (inland and sea area within 500m from the coastline, excluding harbours), and offshore (beyond 500m offshore l). The usage intensity of each habitat category was calculated as the mean proportion of positions in each one at each hour of the day by the study birds, as obtained by the GPS loggers (Supplementary material appendix 1, A2). Fixes were filtered to exclude all positions taken at intervals shorter than one hour. Hence, to estimate the time spent in each habitat category, we considered that each fix corresponded to one hour.

We produced a time-synchronised animation of non-filtered data of gulls and fishing vessels during the previously referred period to investigate the occurrence of gull-fishing vessel interactions (Granadeiro *et al.*, 2011) (Supplementary material appendix 2, Video B4). To describe this behaviour, we recorded the number of gull trips, the number of fishing vessel trips, the number of gull-vessel interactions and the time. For the analysis of the interactions with fishing vessels, we defined a trip as a set of consecutive positions at sea, i.e. farther than 500m from the coast, between two positions on land. An interaction was considered to exist when one gull headed directly towards a fishing vessel and its position was within a 500m radius of the fishing vessel. Gull trips were considered to occur in daylight or in darkness according to the time of local sunrise and sunset (below the horizon) calculated for each day.

#### *Diet assessment*

In late April 2016, we collected blood for stable isotope analysis (SIA) from 15 individuals. Additionally, in late May 2017, we collected 20 regurgitations and 15 samples of growing body feathers from chicks from different broods at Desembarcadouro islet. We calculated frequencies of occurrence as the percentage of regurgitated samples with a given prey type, and numeric frequencies as the number of individuals of each prey type in relation to the total number of individuals. Muscle samples were collected from the regurgitations. Other prey for SIA (known from the literature or personal observations to figure in the YLG diet), including snails, rabbits, Bulwer's Petrels *Bulweria bulwerii* and cow meat were

opportunistically collected (Jorge Camacho *et al.*, 2000; Matias & Catry, 2010; Alonso *et al.*, 2015) (Supplementary material appendix 1, A3).

SIA was carried out using continuous-flow isotope-ratio mass spectrometry (CF-IRMS) (Brand *et al.*, 2015). The contribution of different food sources to the diet of YLGs in Madeira was described using mixing models as implemented in the SIAR package for R (Parnell *et al.*, 2010) (Supplementary material appendix 1, A3).

## Results

### *Distribution of the Yellow-legged Gull*

The deployment of the ten GPS-GSM devices, resulted in an average of  $1132.6 \pm 1107.4$  hourly fixes per individual (range: 13 to 306 days), from 27<sup>th</sup> April 2016 to 23<sup>rd</sup> December 2017.

YLG fixes were concentrated in the south-eastern part of Madeira island. The 50% density contour showed a narrow area of occupancy including the city of Funchal, the Urban Solid Waste Treatment Plant (USWTP) and Santo da Serra golf field (Figure 1). The point-counts conducted on Madeira island (Supplementary material appendix 2, Figure B1) mostly suggested a similar distribution pattern during winter, except for some concentrations in the north-west of the island. A total of 1,662 YLGs were found during these counts.




Figure 1 - Distribution of Yellow-legged Gulls *Larus michahellis atlantis* on Madeira island from 27<sup>th</sup> April 2016 to 23<sup>rd</sup> December 2017. Kernel density contours correspond to 95%



(lighter shade), 75% and 50% (darker shade). USWTP - Urban Solid Waste Treatment Plant. Desembarcadouro and Chão islets are the study breeding colonies.

During the entire annual cycle, GPS-tracked gulls never left Madeira island and Chão islet, keeping within a radius of 56km from Desembarcadouro islet and 70km from Chão islet, and never venturing out into the ocean proper. Excluding commuting movements between Chão islet and Madeira, all positions recorded at sea were within 5km of the coastline, with 61% within 2km of the coastline. Of the GPS positions 94% were terrestrial or less than 2km offshore.

#### *Gull activity in relation to fisheries*

Most fixes (up to 60%) were obtained inland during the day but during the night they were mostly concentrated in the harbours and at-sea areas (Figure 2). Only 6% were at sea (i.e., farther than 500m from the coast), 70% of which were at night.

The fishing activity and the positions of gulls while at sea showed considerable overlap (Supplementary material appendix 2, Figure B2). The analysis of the animations of time-matched gulls and fishing vessels movements in that area, revealed that (at least) 46% of gull trips to the sea involved an interaction with a fishing boat (n=192 trips) and that 91% of those interactions were with three purse seiners. Most (94%) of the interactions (n=89 interactions) occurred between 22:00 and 03:00 (GMT) and purse seiner activity was concentrated between 16:00 and 06:00.

#### *Diet*

Most (80%) of the regurgitations from chicks included marine fish, mainly Blue Jack Mackerel *Trachurus picturatus* and Atlantic Chub Mackerel *Scomber colias* (Table 1). Unidentified fish consisted of pieces with no hard structures that allowed identification and eight small fish found in one regurgitation (Table 1). All food sources were well segregated in at least one of the isotopes ( $\delta^{13}\text{C}$ : ANOVA,  $F_{3,26}=28.5$ ,  $P<0.001$ ;  $\delta^{15}\text{N}$ : ANOVA,  $F_{3,26}=30.2$ ,  $P<0.001$ ) (Figure 3, Supplementary material appendix 2, Figure B3). The results of the mixing models revealed that refuse, marine fish and Bulwer's Petrels constituted the bulk of the diet of adult YLGs (Figure 4a), whereas chicks were mostly fed on fish (Figure 4b).




Figure 2 - Proportion of positions ( $\pm$ SE) of Yellow-legged Gulls *Larus michahellis atlantis* in each habitat category at each hour of the day during the study period, on Madeira island. The shaded areas represent the variation in the sunrise and sunset period throughout the year in Madeira. The sunrise/sunset calculation was done using a  $0^\circ$  inclination of the sun in relation to earth.

Table 1 – Diet of Yellow-legged Gull *Larus michahellis atlantis* chicks on Desembarcadouro islet, in May 2017 (n=20 regurgitations).

Food item	Numeric Frequency (%) (n=29 prey items)	Frequency of Occurrence (%) (n=20 regurgitations)
Fish		
Atlantic Chub Mackerel <i>Scomber colias</i>	13.8	30
Blue Jack Mackerel <i>Trachurus picturatus</i>	10.3	15
Common bogue <i>Boops boops</i>	3.5	5
Unidentified	44.8	30
Insects		
Wart-biter <i>Dectitus albifrons</i>	17.2	10
Refuse		
Chicken meat <i>Gallus domesticus</i>	3.5	5
Potato <i>Solanum tuberosum</i>	3.5	5
Pork meat <i>Sus domesticus</i>	3.5	5




Figure 3 - Distribution of mean ( $\pm$ SD)  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  of adults (blood from April 2016, Chão islet) and chicks (feathers from May 2017, Desembarcadouro islet) of Yellow-legged Gulls *Larus michahellis atlantis* and food categories: Terrestrial herbivores (snails, wart-biters and rabbits), Marine fish (Blue Jack Mackerel, Atlantic Chub Mackerel and Common Bogue), Refuse (chicken, pork and cow meat), and Bulwer's Petrel. Adult and chick stable isotope values are represented after correction for trophic discrimination (i.e. after subtracting 0.38‰ for  $\delta^{13}\text{C}$  and 2.93‰ for  $\delta^{15}\text{N}$  for blood, and 1.97‰ for  $\delta^{13}\text{C}$  and 4.00‰ for  $\delta^{15}\text{N}$  for feathers).




Figure 4 - Proportion of each food category ingested by (a) adults and (b) chicks of Yellow-legged Gull *Larus michahellis atlantis* in April 2016 in Chão islet, and May 2017 in Desembarcadouro islet, respectively. Terrestrial herbivores include snails, wart-biters and rabbits, Marine fish includes Blue Jack Mackerel, Atlantic Chub Mackerel and Common Bogue, Refuse includes chicken, pork and cow meat.

## Discussion

This study investigated the distribution and diet of the YLG on the oceanic island of Madeira. Gulls used mainly human-modified terrestrial environments to feed and rest and made very little use of the marine environment and its natural resources. This limited at-sea activity was mainly associated with purse seiners, which operate near shore and at night. Adult diet included large proportions of refuse, marine fish (probably from fishery discards) and Bulwer's Petrels. The chick diet mainly consisted of marine fish.

### *Distribution of the Yellow-legged gull*

Gulls generally used a relatively small part of Madeira island, mostly occupying human-modified locations, such as the urban solid waste treatment plant (USWTP), golf courses, reservoirs and harbours. Although this is common behaviour for this species, reported in other European populations (Navarro *et al.*, 2017), the YLG population of Madeira stands out for the rare use that it makes of marine environments, which are highly used by continental populations (Ramos *et al.*, 2009a; Ceia *et al.*, 2014). Although our sample size of tracked gulls was small (ten), the distribution obtained was likely to be representative of the population as it was well-matched by the results of the point-counts made around the island during winter, which found 1,662 gulls. The gull distribution and abundance are likely influenced by the greater human presence in the southeast of the island (DREM, 2017) and by greater food availability, mainly of anthropogenic sources, in that area.

Our data confirms that the Madeiran adult population of YLGs is resident. There were no movements farther than *c.* 70km from the breeding colony or farther than 5km from the coastline during the two years that the gulls were tracked.

### *Gull activity in relation to fisheries*

During the day, the gulls mainly used inland areas, whereas at night they moved to harbours, probably to rest. Movements to the sea were occasional and 70% of those trips were between sunset and sunrise, which contrasts with other studies that have reported mostly diurnal activity at sea in synchronisation with fishing vessels present in those areas (González-Solís *et al.*, 1997, 1999). The vessel activity data used in our study was quite complete, including both AIS-T and VMS systems. These surveillance systems are not mandatory in small fishing boats, which were not detected, but such boats are also less likely to produce discards that might attract gulls. Most at-sea positions of gulls took place at night,

when purse seiners comprise the principal fishing gear operating (Vasconcelos *et al.*, 2006). This fleet is active from late afternoon until morning, probably explaining the nocturnal at-sea activity by YLGs. The fishing fleet registered in Madeira constitutes 468 vessels, of which 395 are tuna fishing vessels, 58 use longlines for Black Scabbardfish fishing and three vessels use trolling lines for large pelagic fish. Traps and gillnets are used by five and two vessels, respectively. Purse seiners constitute only 1% of the fishing fleet with five vessels that mainly target Atlantic Chub Mackerel *Scomber colias* and Blue Jack Mackerel *Trachurus picturatus* (Hermida & Delgado, 2016; Vallerani *et al.*, 2017). Interactions with vessels other than purse seiners were scarce, most probably because they produce comparatively few discards or refuse and often operate away from the coast. The low at-sea activity of YLGs in Madeira likely reflects the small size of the purse seiner fleet and the absence of a trawling operation in the area (Camphuysen *et al.*, 1993; Garthe *et al.*, 1996).

### *Diet*

The SIA revealed that adult YLGs consumed mostly refuse, followed by marine fish and Bulwer's Petrels. YLGs frequently prey on small petrels (Neves *et al.*, 2006; Matias & Catry, 2010) and in the region of the studied gull colonies there is a very large aggregation (c. 45,000 pairs) of Bulwer's Petrels (Catry *et al.*, 2014-15). Further, we have witnessed cases of predation on Bulwer's petrels by gulls in the study area. Nevertheless, the estimates of marine fish and Bulwer's Petrel provided by the SI mixing model were negatively correlated (Supplementary material appendix 2, Figure B3), which raises some uncertainty regarding the relative importance of Bulwer's Petrels on the diet of the YLG population from Madeira. Moreover, several other sources of information (e.g. pellet analysis and behavioural observations from unpublished author's information) suggest that the dietary contribution of Bulwer's Petrel may be overestimated by our model at the cost of marine fish, and this issue can only be resolved with more in-depth research. Nevertheless, we believe that the high contribution of refuse fits well with the frequent use of the urban solid waste treatment plant, and that the availability of fish provided by local purse seiners justifies the importance of marine fish, particularly during chick-rearing period. The low use of the marine environment by the gulls in our study and the frequent interactions with fishing vessels suggest that most of the fish consumed is of anthropogenic sources and mainly taken from purse seiners at night, rather than obtained as discards. A high consumption of food from terrestrial sources

by YLGs has also been described on other oceanic islands, namely in the Azores (Neves *et al.*, 2006; Pedro *et al.*, 2013), the Canary islands (Jorge Camacho *et al.*, 2000; López *et al.*, 2016), and on Selvagem Grande Island (Matias & Catry, 2010), where gulls consume mostly refuse, terrestrial invertebrates, seabirds (captured on land) and land snails, respectively. This somewhat contrasts with mainland Europe, where the diet of YLGs is mainly of marine origin, even though a considerable proportion of it probably originates from discards (González-Solís *et al.*, 1997; Ceia *et al.*, 2014; Alonso *et al.*, 2015; Calado *et al.*, 2018; Morera-Pujol *et al.*, 2018). Nonetheless, there are also instances in the western and north-western Mediterranean where terrestrial food sources (refuse and terrestrial invertebrates) are the main food components (Duhem *et al.*, 2003).

In Madeira, YLG chicks were mostly fed marine fish. Adults do not seem to rely on marine resources, so it is likely that purse seiners play an important role in the reproductive success of this population. This seems to be typical for chicks of YLG from various populations (Arizaga *et al.*, 2010, 2013; Ramos *et al.*, 2009b), although variations and alternative prey have also been reported (Pedro *et al.*, 2013; Ceia *et al.*, 2014; Alonso *et al.*, 2015).

The findings of this study confirm that Madeiran YLGs do not exploit natural prey in the pelagic deep-sea environments, as found around the archipelago. This large population apparently thrives due to the existence of anthropogenic food subsidies, mostly obtained inland and also during interactions with purse seiners.

## Acknowledgements

This work was supported by the European Commission through the LIFE project - LIFE RECOVER NATURA (LIFE12 NAT/PT/000195), and Fundação para a Ciência e a Tecnologia (FCT Portugal) through the project Oceanwebs (PTDC/MAR-PRO/0929/2014), the strategic projects UID/MAR/04292/2013 granted to MARE and UID/AMB/50017/2013 granted to CESAM (with co-funding by FEDER, within the PT2020 Partnership Agreement and Compete 2020), and the project IF/00502/2013/CP1186/CT0003 and PTDC/BIAANM/3743/2014. PD/BD/127810/2016 and SFRH/BPD/111084/2015 were granted to JR and JPS, respectively, by FCT. We would like to acknowledge the Instituto das Florestas e Conservação da Natureza, especially Dr. Paulo Oliveira, for permission to undertake experiments with birds and logistical support to carry out the work on the islands

of Desertas and Madeira. Thanks are due to the Direcção Geral de Recursos Naturais, Segurança e Serviços Marítimos (DGRM) for providing the Vessel Monitoring System Data. We also thank the valuable help given by park-wardens Carlos Clemente, Manuel José, Isamberto Silva and volunteers, in the field, and Filipe Moniz, Mafalda Pádua, Gabriel Mendes and Rafael Luís for assistance with laboratory work. We thank to the anonymous reviewer and the editor for the helpful review of the manuscript.

## References

- Alonso, H., Almeida, A., Granadeiro, J.P. & Catry, P. (2015). Temporal and age-related dietary variations in a large population of yellow-legged gulls *Larus michahellis*: implications for management and conservation. *Eur. J. Wildl. Res.*, 61: 819-829. <https://doi.org/10.1007/s10344-015-0958-9>
- Arizaga, J., Aldalur, A., Herrero, A., Cuadrado, J.F., Díez, E. & Crespo, A. (2014). Foraging distances of a resident yellow-legged gull (*Larus michahellis*) population in relation to refuse management on a local scale. *Eur. J. Wildl. Res.*, 60: 171-175. <https://doi.org/10.1007/s10344-013-0761-4>
- Arizaga, J., Aldalur, A., Herrero, A., Cuadrado, J.F., Mendiburu, A. & Sanpera, C. (2010). High importance of fish prey in the diet of Yellow-legged Gull *Larus michahellis* chicks from the southeast Bay of Biscay. *Seabird*, 23: 1-6.
- Arizaga, J., Jover, L., Aldalur, A., Cuadrado, J.F., Herrero, A. & Sanpera, C. (2013). Trophic ecology of a resident Yellow-legged Gull (*Larus michahellis*) population in the Bay of Biscay. *Mar. Environ. Res.*, 87–88: 19-25. <https://doi.org/10.1016/j.marenvres.2013.02.016>
- Brand, W.A., Douthitt, C.B., Fourel, F., Maia, R., Rodrigues, C., Maguas, C. & Prohaska, T. (2015). Gas Source Isotope Ratio Mass Spectrometry (IRMS). In Prohaska, T., Irrgeher, J., Zitek, A. & Jakubowski, N. (eds): *Sector Field Mass Spectrometry for Elemental and Isotopic Analysis*. pp. 500-549. <https://doi.org/10.1039/9781849735407-00500>
- Calado, J.G., Matos, D.M., Ramos, J.A., Moniz, F., Ceia, F.R., Granadeiro, J.P. & Paiva, V.H. (2018). Seasonal and annual differences in the foraging ecology of two gull

- species breeding in sympatry and their use of fishery discards. *J. Avian Biol.*, 49: 1-12.  
<https://doi.org/10.1111/jav.01463>
- Calenge, C. (2017). Home Range Estimation in R: the adehabitatHR Package. The Comprehensive R Archive Network. Available at: <https://cran.r-project.org/web/packages/adehabitatHR/adehabitatHR.pdf> (accessed 1 March 2018).
- Camphuysen, C.J., Ensor, K., Furness, R.W., Garthe, S., Hüppop, O., Leaper, G., Offringa, H. & Tasker, M.L. (1993). Seabirds feeding on discards in winter in the North Sea. EC DG XIV research contract 92/3505. NIOZ Rapport 1993 – 8, Netherlands Institute for Sea Research, Texel: 140.
- Catry, P., Dias, M., Catry, T., Pedro, P., Tenreiro, P. & Menezes, D. (2014-15). Bulwer's petrels breeding on the Desertas Islands (Madeira): improved estimates indicate the NE Atlantic population to be much larger than previously thought. *Airo*, 23: 10-14.
- Ceia, F.R., Paiva, V.H., Fidalgo, V., Morais, L., Baeta, A., Crisóstomo, P., Mourato, E., Garthe, S., Marques, J.C. & Ramos, J.A. (2014). Annual and seasonal consistency in the feeding ecology of an opportunistic species, the yellow-legged gull *Larus michahellis*. *Mar. Ecol. Prog. Ser.*, 497: 273-284. <https://doi.org/10.3354/meps10586>
- DREM (2017). Estatísticas Demográficas da região Autónoma da Madeira - Ano 2017. Available at: <https://estatistica.madeira.gov.pt/download-now-3/social-gb/popcondsoc-gb/demografia-gb/demografia-publicacoes-gb/finish/223-demografia-publicacoes/10054-estatisticas-demograficas-da-ram-2017.html> (accessed 15 September 2018).
- Duhem, C., Roche, P., Vidal, E. & Tatoni, T. (2008). Effects of anthropogenic food resources on yellow-legged gull colony size on Mediterranean islands. *Popul. Ecol.*, 50: 91-100.  
<https://doi.org/10.1007/s10144-007-0059-z>
- Duhem, C., Vidal, E., Legrand, J. & Tatoni, T. (2003). Opportunistic feeding responses of the Yellow-legged Gull *Larus michahellis* to accessibility of refuse dumps. *Bird Study*, 50: 61-67. <https://doi.org/10.1080/00063650309461291>
- Galarza, A., Herrero, A., Domínguez, J.M., Aldalur, A. & Arizaga, J. (2012). Movements of Mediterranean Yellow-legged Gulls *Larus michahellis* to the Bay of Biscay. *Ringling & Migration*, 27: 26-31. <https://doi.org/10.1080/03078698.2012.691060>



- Garthe, S., Camphuysen, C.J. & Furness, R.W. (1996). Amounts of discards by commercial fisheries and their significance as food for seabirds in the North Sea. *Mar. Ecol. Prog. Ser.*, 136: 1-11. <https://doi.org/10.3354/meps136001>
- Granadeiro, J.P., Phillips, R.A., Brickle, P. & Catry, P. (2011). Albatrosses Following Fishing Vessels: How Badly Hooked Are They on an Easy Meal? *PLoS ONE*, 6: e17467. <https://doi.org/10.1371/journal.pone.0017467>
- González-Solís, J., Oro, D., Jover, L., Ruiz, X. & Pedrocchi, V. (1997). Trophic niche width and overlap of two sympatric gulls in the southwestern mediterranean. *Oecologia*, 112: 75-80. <https://doi.org/10.1007/s004420050285>
- González-Solís, J., Ruiz, X. & Jover, L. (1999). Fisheries and daily activity cycles of Audouin's *Larus audouinii* and Yellow-legged Gulls *L.cachinnans* breeding at the Chafarinas Islands (Moroccan coast). *Die Vogelwarte*, 40: 52-56.
- Harris, M.P. (1970). Rates and Causes of Increases of some British Gull Populations. *Bird Study*, 17: 325-335. <https://doi.org/10.1080/00063657009476275>
- Hermida, M. & Delgado, J. (2016). High trophic level and low diversity: Would Madeira benefit from fishing down? *Mar. Policy*, 73: 130-137. <https://doi.org/10.1016/j.marpol.2016.07.013>
- Hernández-Matías, A., & Ruiz, X. (2003). Predation on common tern eggs by the yellow-legged gull at the Ebro Delta. *Scientia Marina*, 67: 95-101.
- Jorge Camacho, M. A., Schuster, C. & Acosta Díaz, C. (2000). Dieta primaveral de la gaviota patiamarilla, *Larus cachinnans*, en Alegranza, islas Canarias. *Vieraea*, 28: 31-37.
- Karris, G., Ketsilis-Rinis, V., Kalogeropoulou, A., Xirouchakis, S., Machias, A., Maina, I., Kavadas, S. (2018). The use of demersal trawling discards as a food source for two scavenging seabird species: a case study of an eastern Mediterranean oligotrophic marine ecosystem. *Avian Res.* 9: 1-14
- López, H., Pérez, A.J., Rumeu, B. & Nogales, M. (2016). Trophic strategies of Yellow-legged Gull *Larus michahellis* on oceanic islands surrounded by deep waters. *Bird Study*, 63: 337-345. <https://doi.org/10.1080/00063657.2016.1194804>
- Matias, R. & Catry, P. (2010). The diet of Atlantic Yellow-legged Gulls (*Larus michahellis atlantis*) at an oceanic seabird colony: estimating predatory impact upon breeding petrels. *Eur. J. Wildl. Res.*, 56: 861–869. <https://doi.org/10.1007/s10344-010-0384-y>

- Moreno, R., Jover, L., Munilla, I., Velando, A. & Sanpera, C. (2010). A three-isotope approach to disentangling the diet of a generalist consumer: the yellow-legged gull in northwest Spain. *Mar. Biol.*, 157: 545-553. <https://doi.org/10.1007/s00227-009-1340-9>
- Morera-Pujol, V., Ramos, R., Pérez-Méndez, N., Cerdà-Cuéllar, M., & González-Solís, J. (2018). Multi-isotopic assessments of spatio-temporal diet variability: the case of two sympatric gulls in the western Mediterranean. *Marine Ecology Progress Series*, 606: 201-214.
- Navarro, J., Grémillet, D., Afán, I., Ramírez, F., Bouten, W., & Forero, M. G. (2016). Feathered detectives: real-time GPS tracking of scavenging gulls pinpoints illegal waste dumping. *PloS ONE*, 11: e0159974.
- Navarro, J., Grémillet, D., Ramirez, F.J., Afán, I., Bouten, W. & Forero, M.G. (2017). Shifting individual habitat specialization of a successful predator living in anthropogenic landscapes. *Mar. Ecol. Prog. Ser.*, 578: 243-251. <https://doi.org/10.3354/meps12124>
- Neves, V.C., Murdoch, N. & Furness, R.W. (2006). Population status and diet of the Yellow-legged Gull in the Azores. *Arquipélago. Life and Marine Sciences*, 23A: 59-73.
- Oro, D., Bosch, M. & Ruiz, X. (1995). Effects of a trawling moratorium on the breeding success of the Yellow-legged Gull *Larus cachinnans*. *Ibis*, 137: 547-549. <https://doi.org/10.1111/j.1474-919X.1995.tb03265.x>
- Parnell, A.C., Inger, R., Bearhop, S. & Jackson, A.L. (2010). Source Partitioning Using Stable Isotopes: Coping with Too Much Variation. *PLoS ONE*, 5: e9672. <https://doi.org/10.1371/journal.pone.0009672>
- Pedro, P.I., Ramos, J.A., Neves, V.C. & Paiva, V.H. (2013). Past and present trophic position and decadal changes in diet of Yellow-legged Gull in the Azores Archipelago, NE Atlantic. *Eur. J. Wildl. Res.*, 59: 833-845. <https://doi.org/10.1007/s10344-013-0737-4>
- Ramos, R., Ramírez, F., Sanpera, C., Jover, L. & Ruiz, X. (2009a). Feeding ecology of yellow-legged gulls *Larus michahellis* in the western Mediterranean: a comparative assessment using conventional and isotopic methods. *Mar. Ecol. Prog. Ser.*, 377: 289–297. <https://doi.org/10.3354/meps07792>

- Ramos, R., Ramírez, F., Sanpera, C., Jover, L. & Ruiz, X. (2009b). Diet of Yellow-legged Gull (*Larus michahellis*) chicks along the Spanish Western Mediterranean coast: the relevance of refuse dumps. *J. Ornithol.*, 150: 265-272. <https://doi.org/10.1007/s10336-008-0346-2>
- Ramos, R., Cerdà-Cuellar, M., Ramírez, F., Jover, L., & Ruiz, X. (2010). Influence of refuse sites on the prevalence of *Campylobacter* spp. and *Salmonella* serovars in seagulls. *Applied and environmental microbiology*, 76: 3052-3056.
- Thaxter C.B., Ross-Smith V.H., Clark, J.A., Clark, N.A., Conway, G.J., Marsh, M., Leat, E.H.K. & Burton, N.H.K. (2014). A trial of three harness attachment methods and their suitability for long-term use on Lesser Black-backed Gulls and Great Skuas. *Ringling & Migration*, 29: 65-76. <https://doi.org/10.1080/03078698.2014.995546>
- Vasconcelos, J., Alves, A., Gouveia, E., Faria, G. (2006). Age and growth of the blue jack mackerel, *Trachurus picturatus* Bowdich, 1825 (Pisces: Teleostei) off Madeira archipelago. *Arquipélago. Life and Marine Sciences*, 23A: 47-57.

## Supplementary Material 1

### *Supplementary material appendix 1, A1: Methods for point-counts of gulls in winter*

We carried out 19 point-counts at strategic sites around the island of Madeira (14 points) and at five other locations further inland to assess the distribution of the Yellow-legged Gull *Larus michahellis atlantis* (YLG) on that island. Counting points were initially selected from a 1km grid all around the island, but some quadrats that were not easily accessed by car were not counted. This work took place during two days in January 2017 and involved counting (with a telescope 20x) all individuals visible, whether landed, flying or feeding, within 1,000m. No distinction was made between adults and immature birds.

### *Supplementary material appendix 1, A2: Filtering of the gull tracking data, and AIS-T and VMS data*

We deployed GPS-GSM devices on ten YLGs in Madeira island with the aim of describing the distribution of the population. Some technical problems arose causing for larger intervals between positions taken, leading to the need of data regularisation. Hourly positions were more frequent and congruent throughout all GPS devices; therefore, we regularised all trajectories by linear interpolation of fixes. Also, different GPS units recorded data over different periods. Therefore, some of the analyses were based on randomly sampled fixes from each individual, based on the GPS with the smallest number of fixes (n=176). To study the areas used by the gulls in Madeira, both for foraging and resting, we excluded the fixes at colonies. At-sea positions between the breeding colony, Chão islet and the main island were completely linear and represented commuting movements between the islands, not foraging activity, so were also excluded. The previous conditions were applied to all tracking data analyses.

VMS and AIS-T data were used to characterise the activity area of fishing vessels in the vicinity of Madeira island at the time positions of YLGs were present in the sea. VMS data were preferably used while AIS-T data complemented the database with a few fishing vessels that were not present in the VMS dataset. The sampling intervals of VMS and AIS-T data differed among vessels and therefore we regularised the data by only keeping vessel positions that differed by at least one hour. Furthermore, we filtered these positions for speeds under 4 knots. The AIS-T and VMS data represented a total of 123 vessels and 10,854 vessel positions, and the analysed timespan included 6 gulls. We then constructed a 50%, 75% and

95% kernel density of fishing vessels in activity, using filtered AIS-T and VMS data and the *adehabitatHR* package ( $h=0.03$ ,  $grid=300$ ; Calenge, 2017), under R software (R Core Team, version 1.0.143).

*Supplementary material appendix I, A3: Stable Isotope Analysis and treatment of blood, feathers and muscle*

To assess the diet of the YLGs in the Archipelago of Madeira, we used stable isotope analysis of adults (blood) and chicks (feathers) of YLGs and their preys (muscles), collected in April 2016 and May 2017. The feathers collected from chicks had just grown, which meant that the isotopic signature represented the diet consumed during the chick rearing period. Because feathers of adults start moulting later in August (Monteiro *et al.*, 1996), after the chick rearing, they represented a different time period than that of the chick feathers. Therefore, blood of adults was collected for comparison of the same period as the chicks.

Spontaneous regurgitations of chicks provided some diet samples, and prey were identified to the lowest taxonomic level, using hard structures like vertebrae and/or otoliths, and an extensive reference collection. Among prey species identified were fish (*Scomber colias*, *Trachurus picturatus*, *Boops boops*), wart-biters (*Dectitus albifrons*), and pork and chicken meat.

Bulwer's Petrels have never been found in chick regurgitations or ever described as a dietary item, unlike in adults (Ramos *et al.*, 2009a, Arizaga *et al.*, 2010, 2013, Alonso *et al.*, 2015). Therefore, Bulwer's Petrels were not considered to represent prey in the chick diet.

Feathers were washed with 10% NaOH solution to clean them from external particles (Bearhop *et al.*, 2000, Arizaga *et al.*, 2013), and then thoroughly washed with distilled water. Blood samples from adults did not undergo any treatment before isotopic analysis (Bearhop *et al.*, 2000, Granadeiro *et al.*, 2013). Prior to isotopic analysis of Carbon and Nitrogen, muscle samples from prey were treated with a 2:1 chloroform-methanol solution to extract the lipids of the tissues.

SIA was carried out using continuous-flow isotope-ratio mass spectrometry (CF-IRMS) (Brand *et al.*, 2015). Results are represented in parts per thousand (‰) relative to the Pee Dee Belemnite (PDB) for  $\delta^{13}\text{C}$ , and atmospheric nitrogen for  $\delta^{15}\text{N}$ . The contribution of different food sources to the diet of YLG in Madeira was described using mixing models as

implemented in the SIAR package for R (Parnell *et al.*, 2010). Prey were grouped according to their trophic guild: terrestrial herbivores (including snails, wart-biters and rabbits), refuse (chicken, pork and cow meat), marine fish (Atlantic chub mackerels, blue jack mackerels and bogues), and Bulwer's Petrel. We could not find discrimination factors for diet-blood and diet-feather for YLG, and therefore we calculated average values derived from literature (Hobson & Clark, 1992, Mizutani *et al.*, 1992, Bearhop *et al.*, 1999, 2002, Ogden *et al.*, 2004, Cherel *et al.*, 2005, Williams *et al.*, 2007, Ramos *et al.*, 2009b): diet-blood:  $2.93 \pm 0.45\text{‰}$  for  $\delta^{15}\text{N}$  and  $0.39 \pm 0.96\text{‰}$  for  $\delta^{13}\text{C}$ , and diet-feather:  $4.00 \pm 1.07\text{‰}$  for  $\delta^{15}\text{N}$  and  $1.97 \pm 1.54\text{‰}$  for  $\delta^{13}\text{C}$ .

## References

- Alonso, H., Almeida, A., Granadeiro, J.P. & Catry, P. (2015). Temporal and age-related dietary variations in a large population of yellow-legged gulls *Larus michahellis*: implications for management and conservation. *Eur. J. Wildl. Res.*, 61: 819-829. <https://doi.org/10.1007/s10344-015-0958-9>
- Arizaga, J., Aldalur, A., Herrero, A., Cuadrado, J.F., Mendiburu, A. & Sanpera, C. (2010). High importance of fish prey in the diet of Yellow-legged Gull *Larus michahellis* chicks from the southeast Bay of Biscay. *Seabird*, 23: 1-6.
- Arizaga, J., Jover, L., Aldalur, A., Cuadrado, J.F., Herrero, A. & Sanpera, C. (2013). Trophic ecology of a resident Yellow-legged Gull (*Larus michahellis*) population in the Bay of Biscay. *Mar. Environ. Res.*, 87–88: 19-25. <https://doi.org/10.1016/j.marenvres.2013.02.016>
- Bearhop, S., Phillips, R.A., Thompson, D.R., Waldron, S. & Furness, R.W. (2000). Variability in mercury concentrations of great skuas *Catharacta skua*: The influence of colony, diet and trophic status inferred from stable isotope signatures. *Mar. Ecol. Prog. Ser.*, 195: 261-268. <https://doi.org/10.3354/meps195261>
- Bearhop, S., Thompson, D.R., Waldron, S., Russell, I.C., Alexander, G. & Furness, R.W. (1999). Stable isotopes indicate the extent of freshwater feeding by cormorants *Phalacrocorax carbo* shot at inland fisheries in England. *J. Appl. Ecol.*, 36: 75-84. <https://doi.org/10.1046/j.1365-2664.1999.00378.x>
- Bearhop, S., Waldron, S., Votier, S.C. & Furness, R.W. (2002). Factors That Influence Assimilation Rates and Fractionation of Nitrogen and Carbon Stable Isotopes in Avian

- Blood and Feathers. *Physiol. Biochem. Zool.*, 75: 451-458.  
<https://doi.org/10.1086/342800>
- Brand, W.A., Douthitt, C.B., Fourel, F., Maia, R., Rodrigues, C., Maguas, C. & Prohaska, T. (2015). Gas Source Isotope Ratio Mass Spectrometry (IRMS). In Prohaska, T., Irrgeher, J., Zitek, A. & Jakubowski, N. (eds) *Sector Field Mass Spectrometry for Elemental and Isotopic Analysis*. pp. 500-549.  
<https://doi.org/10.1039/9781849735407-00500>
- Calenge, C. (2017). Home Range Estimation in R: the adehabitatHR Package. The Comprehensive R Archive Network. Available at: <https://cran.r-project.org/web/packages/adehabitatHR/adehabitatHR.pdf> (accessed 1 March 2018).
- Cherel, Y., Hobson, K.A. & Hassani, S. (2005). Isotopic Discrimination between Food and Blood and Feathers of Captive Penguins: Implications for Dietary Studies in the Wild. *Physiol. Biochem. Zool.*, 78: 106-115. <https://doi.org/10.1086/425202>
- Granadeiro, J.P., Brickle, P. & Catry, P. (2013). Do individual seabirds specialize in fisheries' waste? The case of black-browed albatrosses foraging over the Patagonian Shelf. *Anim. Conserv.*, 17: 19-26. <https://doi.org/10.1111/acv.12050>
- Hobson, K.A. & Clark, R.G. (1992). Assessing Avian Diets Using Stable Isotopes II: Factors Influencing Diet-Tissue Fractionation. *Condor*, 94: 189-197.  
<https://doi.org/10.2307/1368808>
- Mizutani, H., Fukuda, M. & Kabaya, Y. (1992).  $^{13}\text{C}$  and  $^{15}\text{N}$  Enrichment Factors of Feathers of 11 Species of Adult Birds. *Ecology*, 73: 1391-1395.  
<https://doi.org/10.2307/1940684>
- Monteiro, L.R., Ramos, J.A., Furness, R.W. & Delnevo, A.J. (1996). Movements, Morphology, Breeding, Molt, Diet and Feeding of Seabirds in the Azores. *Colon. Waterbirds*, 19: 82-97. <https://doi.org/10.2307/1521810>
- Ogden, L.J.E., Hobson, K.A. & Lank, D.B. (2004). Blood Isotopic ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) Turnover and Diet-Tissue Fractionation Factors in Captive Dunlin (*Calidris alpina pacifica*). *Auk*, 121: 170-177. [https://doi.org/10.1642/0004-8038\(2004\)121\[0170:BICANT\]2.0.CO;2](https://doi.org/10.1642/0004-8038(2004)121[0170:BICANT]2.0.CO;2)

- Parnell, A.C., Inger, R., Bearhop, S. & Jackson, A.L. (2010). Source Partitioning Using Stable Isotopes: Coping with Too Much Variation. PLoS ONE, 5: e9672. <https://doi.org/10.1371/journal.pone.0009672>
- Ramos, R., Ramírez, F., Sanpera, C., Jover, L. & Ruiz, X. (2009a). Diet of Yellow-legged Gull (*Larus michahellis*) chicks along the Spanish Western Mediterranean coast: the relevance of refuse dumps. J. Ornithol., 150: 265-272. <https://doi.org/10.1007/s10336-008-0346-2>
- Ramos, R., Ramírez, F., Sanpera, C., Jover, L. & Ruiz, X. (2009b). Feeding ecology of yellow-legged gulls *Larus michahellis* in the western Mediterranean: a comparative assessment using conventional and isotopic methods. Mar. Ecol. Prog. Ser., 377: 289-297. <https://doi.org/10.3354/meps07792>
- Williams, C.T., Buck, C.L., Sears, J. & Kitaysky, A.S. (2007). Effects of nutritional restriction on nitrogen and carbon stable isotopes in growing seabirds. Oecologia, 153: 11-18. <https://doi.org/10.1007/s00442-007-0717-z>



## Supplementary Material 2




Figure B1 - Total numbers of Yellow-legged Gulls *Larus michahellis atlantis* at each point-count site (black points) in January 2017, in Madeira island. Desembarcadouro and Chão islet are the breeding colonies studied.




Figure B2 – Kernel density estimates of fishing activity (vessels travelling at <4 knots) around Madeira island, from 1<sup>st</sup> April 2016 to 30<sup>th</sup> September 2016 and at-sea positions of Yellow-legged Gulls *Larus michahellis atlantis* during the study period. The kernel density is represented by the 95%, 75% and 50% contours, with increasing shading. White points represent at-sea gull locations during the study period.




Figure B3 – Matrixplot of the proportions of food categories consumed by adult Yellow-legged Gulls *Larus michahellis atlantis* in Madeira. Terrestrial herbivores (snails, wart-biters and rabbits), Marine fish (Blue Jack Mackerel, Atlantic Chub Mackerel and Common Bogue), Refuse (chicken, pork and cow meat), and Bulwer’s Petrel.

Video B4 – Example of the animation of time-matched Yellow-legged Gulls *Larus michahellis atlantis* and fishing vessels movements in the Archipelago of Madeira between April and September 2016. Fishing vessels are represented by black points and gulls by red points. Access: <https://bioone.org/journals/ardeola/volume-66/issue-1/arla.66.1.2019.sc4/A-Gull-that-Scarcely-Ventures-on-the-Ocean--Yellow/10.13157/arla.66.1.2019.sc4.short>

## CHAPTER 5

---



**Credits: Joana Romero**

*This chapter is published as:*

*Joana Romero, Paulo Catry, Hany Alonso, José Pedro Granadeiro (2021) Seabird diet analysis suggests sudden shift in the pelagic communities of the subtropical Northeast Atlantic. Marine Environmental Research, 165: 105232.*

# Seabird diet analysis suggests sudden shift in the pelagic communities of the subtropical Northeast Atlantic

## Abstract

The dynamics of the subtropical pelagic ecosystems of the Northeast Atlantic are still poorly known due to the high costs associated with sampling large oceanic areas. Top predators can be used as alternative low-cost samplers and indicators of the temporal variability of such systems. To study the variation in the composition of pelagic species through time in the broad Canary current region, we analysed foraging trips and regurgitations of Cory's shearwaters *Calonectris borealis* nesting on Selvagens Islands, in 2008-2011 and 2016-2018. Fisheries data, oceanographic variables and the North Atlantic Oscillation were explored as possible explanatory variables for trends in behaviour and diet. Cory's shearwaters diet, complemented by fisheries data, revealed marked changes in the composition of the pelagic fish communities. In 2016 there was a peak in the abundance of the Atlantic chub mackerel *Scomber colias*, followed by an explosive increase in the abundance of the Longspine snipefish *Macroramphosus scolopax* in 2017 and 2018, as deduced from the diet composition of the Cory's shearwater, and supported by fisheries data, in the broad oceanic area surrounding the Selvagens Islands. Oceanographic variables did not show fluctuations correlated with these marked shifts in pelagic fish availability, the causes of which remain largely unknown. This study highlights the importance of the Atlantic chub mackerel and of the Longspine snipefish in the Madeira/Canary region and exemplifies the efficiency of avian predators in revealing rapid changes in pelagic communities of oceanic domains. Such trends and variations need to be better monitored and understood to measure the impact of ongoing global changes and to sustainably manage the marine environment and resources.

**Keywords:** Seabirds, Ecosystem change, Pelagic, Canary Current, *Calonectris borealis*, *Macroramphosus scolopax*, *Scomber colias*, Community shift

## Introduction

The high-seas of the subtropical eastern North Atlantic are a nutrient-poor region, contrasting with the richer waters of the strong coastal upwelling of West Africa associated with the Canary current (Cropper et al., 2014). The existence of seamounts and islands in an otherwise deep-ocean region enhances conditions for many pelagic organisms to thrive in this area (Morato et al., 2008; Pitcher et al., 2007).

Even though predators at high trophic levels dominate the pelagic community with a large number of species in this region, all trophic levels are characterized by a low biomass when comparing with the more productive coastal region (Pitcher et al., 2007). Among these forage fish are the Atlantic chub mackerel *Scomber colias* (hereafter simply chub mackerel) and the Blue jack mackerel *Trachurus picturatus* which are some of the most abundant and with higher productivity rates of this type of ecosystem (Hermida and Delgado, 2016). Other vital, but less studied marine organisms, include cephalopods and mesopelagic fish which sustain not only deep-sea fish (Fock et al., 2002a) but also epipelagic feeders, like seabirds (Waap et al., 2017). In addition, their daily vertical migration creates an important link between the depths and the epipelagic domain (Salvanes and Kristoffersen, 2001).

Small pelagic fish have been studied extensively in coastal shelf areas where they often represent commercially important fisheries. In neritic environments, oceanographic variables (e.g. Thiaw et al. 2017) and environmental indexes have been suggested as explanations for the variations and shifts in the reproduction or migration of marine species (e.g. Edwards et al. 2013). The only other pelagic habitat which has been further studied are seamounts (Pitcher et al., 2007). On the other hand, the open ocean beyond shelf waters is among the least known biomes on Earth because of the logistic and monetary challenges of sampling these areas. Data on the ecology and temporal fluctuations of pelagic species in areas such as the oceanic region around the Madeira and Canary archipelagos, off NW Africa, are still very limited. Here, there have been some attempts to study the composition, abundance and distribution of pelagic and demersal species with the help of research or fishing vessels (Christiansen et al., 2009; Delgado et al., 2018; Fock et al., 2002b). However, these studies were mostly species-directed and few presented a multi-specific approach.

Top predators with high mobility, such as cetaceans, tuna, and seabirds, have been successfully used as tools to sample large areas of the marine environment (e.g. Grebmeier

et al. 2006; Pethybridge et al. 2015; Church et al. 2018). Studies on their diet and foraging behaviour enable scientists to gather useful information on the composition and the spatial and temporal distribution of the trophic groups that constitute their prey (Velarde et al., 1994). The Cory's shearwater *Calonectris borealis* has been the target of long-term studies which have provided important information on their feeding ecology and behaviour (e.g. Alonso et al., 2018, 2012). This pelagic seabird nests in several islands of the Macaronesian region, including the Selvagem islands, a small group of oceanic islands ca. 300 km south of the Madeira archipelago. During the breeding season, birds from Selvagem typically travel several hundred kilometres in a single foraging trip, targeting pelagic areas around the island, seamounts and also neritic areas over the African continental shelf (Alonso et al., 2018). They are generalist predators, feeding on a variety of (mostly shoaling) pelagic fish and squid species caught during the day near the surface of the ocean (Alonso et al., 2014; Granadeiro et al., 1998a). Cory's shearwaters are also known to occasionally interact with underwater predators, like tunas and cetaceans, which drive prey to the surface and make them available to seabirds (Veit and Harrison, 2017).

As an abundance and widespread species in this region, these seabirds are potential indicators of important variations in epipelagic communities that might pinpoint changes in the structure of prey communities. In the present study, we describe the interannual variability of the pelagic community in the broad region surrounding the Selvagens islands, using the Cory's shearwater as a biological sampler. We combined dietary information with the location of their main foraging areas and fisheries data in order to assess the temporal variations in the composition and distribution of the most common pelagic species in the region. Concurrently, we examined the changes in oceanographic variables, to assess the extent to which they might explain the variation of the community in this oceanic area. Finally, we examine the relationship between the observed changes in the diet and the potential consequences for the reproductive success of this species.

## Methods

Fieldwork was carried out at Selvagem Grande island (30° 09' N, 15° 52' W, Figure 1) where data was collected during the chick-rearing seasons of 2008-2011, and 2016-2018.




Figure 1 – (A) Foraging areas of Cory's shearwaters *Calonectris borealis* during the breeding season in Selvagem Grande in 2008-2011 and 2016-2018, as revealed by the present study (shadowed area corresponds to 95% Kernel Density Estimation). (B) Domains used by Cory's shearwaters during their foraging trips (shadowed areas correspond to 50% Kernel Density Estimation) and examples of foraging trips to Pelagic areas (▲), Seamounts (●), and the African coast (■). Isobathic lines of 200m, 1000m and 2000m are represented as black lines.

## Diet

### *Sampling and identification of prey*

A total of 827 regurgitations were collected between July-late August of 2008-2011 and 2016-2018. Breeding individuals were captured upon their arrival at the nest to feed their chick at dusk and their stomach content was collected by stomach flushing with salt water (Wilson, 1984). Individuals were then marked with colour (livestock paint crayon) to avoid any subsequent disturbance. The stomach content was then carefully sieved to remove excess water and immediately stored in a container and frozen until analysis in the laboratory. Less

digested fish prey were identified morphologically to the lowest taxonomic level, using identification guides. Digested items (levels III and IV, following Alonso et al., 2018) were ascribed to species based on our reference collection of fish skeletons and otoliths from the NE Atlantic (over 700 specimens of ca. 100 species).

Whenever the taxonomic identification of a prey by hard structures was not possible, a muscle tissue sample was collected from the remaining bones, stored in 96% alcohol, and frozen to be identified by DNA barcoding (Alonso et al., 2014). A total of 103 muscle samples belonging to 56 fish and 47 cephalopod prey were collected from Cory's shearwaters regurgitations. DNA was extracted from these samples using the E.Z.N.A. Tissue DNA kit (Omega Bio-tek). The 3' end region of the 16S rRNA gene of fish and cephalopod DNA was amplified with the universal primers 16Sar and 16SSbr (Palumbi, 1996) using optimized PCR conditions (Alonso et al., 2014). The PCR products were sequenced in both directions in outsourcing (Macrogen Inc). The resulting sequences were used in BLAST (NCBI) searches and similarity values higher than 98% were considered as a positive identification for the queried sample.

For the purpose of this study we did not distinguish between different Chub mackerel species and refer to them as *Scomber* sp., as several prey items were only identified to genus level. Nevertheless, only chub mackerel *Scomber colias* was identified to species level within the samples. Eroded cephalopod beaks were not included in the analysis of the diet because they tend to remain in the stomach of seabirds for long periods (weeks) after ingestion, resulting in an over-representation of this group when they are included.

#### *Data analysis*

All fish found whole in the regurgitations were weighted (wet weight,  $\pm 0.1$ g) and measured (standard length,  $\pm 1$ mm). Whenever that was not possible, fish size was estimated from the size of specific vertebrae using published equations (Granadeiro and Silva, 2000). The standard length (SL) of Pilotfish was estimated from the length of the first caudal vertebrae (CV);  $1^{st} CV$ ;  $SL = 1.602 + 1^{st} CV \times 29.998$ ,  $r^2=0.98$ ,  $n=14$  (SL=30–118mm) (H. Alonso, unpubl).

Numeric frequencies (NF, number of individuals of a given prey type as a percentage of the total number of prey items), and frequencies of occurrence (FO, number of samples



with a given prey type as a percentage of the total number of samples) were calculated for each prey species (or lowest taxonomic level identified for a prey item) and year.

A Permutational Multivariate Analysis of Variance (PERMANOVA) was performed to check for significant differences in the diet of the Cory's shearwater among years. The PERMANOVA was carried out using the R *vegan* package with Jaccard similarities on prey presence/absence in regurgitations (FO) (Oksanen et al., 2019). In order to compare the diet diversity of this species among years, the Shannon-Wiener diversity index ( $H'$ ) was calculated for numeric frequency. To avoid bias due to the different number of samples collected in each year, we resampled the smallest number of samples collected ( $n=30$ ) and carried it out 1000 times for each year. Means and standard deviations were then calculated from those values.

### **GPS deployment and foraging trip analysis**

GPS-loggers were deployed in a total of 150 breeding Cory's shearwaters during the chick rearing season of 2009-2011 (12, 24, and 43 individuals, respectively), 2016 (43) and 2018 (28). The weight of GPS devices (iGotU GPS loggers, 17g) represented ca. 2% of the average weight of an adult Cory's shearwater, therefore unlikely to have deleterious effect on the behaviour of birds (Phillips et al., 2003). GPS loggers were attached to four central tail feathers with TESA tape, and deployed in the nests at dusk, while the adults were visiting the chicks. Upon return from the feeding trip, the bird was recaptured, and the GPS retrieved. Most trips lasted between one and three days (79%,  $n=306$ ), although some lasted up to 11 days.

Positional data were downloaded from the loggers and divided into separate feeding trips, each starting and ending in Selvagem Island. A total of 306 trips were discriminated from the data. Different GPS loggers were programmed to collect fixes between 10min and 60min intervals. Consequently, in order to allow for comparison among birds, all trips were processed to only include hourly positions, discarding all remaining fixes.

Area utilization was calculated with Kernel Density Estimation (KDE) using the R software package *adehabitatHR* (Calenge, 2015). The home-range and the main foraging areas were considered as those comprising 95% and 50% of the positions, respectively. We compared home ranges and foraging areas of different years using the Utilization Distribution Overlap Index (UDOI), made available through function *kerneloverlap*.

The 50% KDE created by the combination of all trips, revealed 4 main foraging domains: Selvagem (2000m bathymetric line around the Selvagem archipelago); Pelagic (circle with a 160km radius around Selvagem and with depths between 2000 and 4000m); Seamounts (2000m bathymetric line around the seamounts) and African coast (area defined by latitudes 20-35°N and 80km offshore) (Alonso et al., 2018, 2012) (Figure1B). Due to different oceanographic conditions, the African Coast area was further subdivided into Strong Permanent Upwelling Region (SUR, 20-26°N) and Weak Permanent Upwelling Region (WUR, 26-35°N) (Cropper et al., 2014), creating five domains with different environmental features. One of these domains was assigned to each feeding trip as its destination. The destination was assigned as the domain with the highest number of positions of that trip (and the most convoluted part of the track), which was usually also the area with the farthest locations from the colony. Birds sometimes crossed more than one oceanographic domain, but one of these was much more used than the others.

The frequency of occurrence of each foraging domain was calculated to quantify its use by Cory's shearwaters each year. Permutational multivariate analysis of variance (PERMANOVA) was used to test for interannual variability in foraging areas and was carried out using the *vegan* package with Jaccard similarities on destination of each foraging trip (Oksanen et al., 2019). To calculate the maximum distances from the colony reached by each individual, distances between each position and Selvagem were calculated using the function *spDistsNI* from the package *sp* (Pebesma et al., 2018). Differences among years in maximum distances from the colony were tested using a one-way ANOVA. All statistical analyses were performed using the Software R (version 3.5.2).

### **Fisheries and environmental data**

The total annual landings of chub mackerel from Madeira Island from local purse-seiners were obtained from the Regional Direction of Fisheries of the Autonomous Region of Madeira and used as an indicator of the abundance of this species between 2008 and 2018.

Monthly 4km AQUAMODIS Chlorophyll a (Chla) and SST data were downloaded from the ERDDAP data server (<https://coastwatch.pfeg.noaa.gov/erddap/index.html>). Monthly 0.125° zonal (u) and meridional (v) wind components at 10 meter were downloaded from the European Centre for Medium-Range Weather Forecasts (ECMWF) using the ERA-Interim server (<https://apps.ecmwf.int/datasets/>). The NAO index was obtained from the

Climate Data Guide server (<https://climatedataguide.ucar.edu/climate-data/hurrell-north-atlantic-oscillation-nao-index-station-based>). Data on these variables were obtained for January to April (hereafter “Winter”, corresponding to the strongest period of upwelling off the African coast and, thus, the highest Chl a values) and June to September (hereafter “Summer”, corresponding to our study period and second peak in Chl a) between 2008-2018, to assess seasonal and interannual variability (Davenport et al., 1999). Each variable was averaged within the domains. To do that, the median value of each month was obtained for each domain, and then we calculated the mean for each season. Mean Chl a values were then log transformed for further analysis. Climatic (NAO) and environmental (Chl a, SST, Wind speed and direction) variables were analysed for correlation with the prey occurrence and with the frequency of occurrence of Cory’s shearwaters in different domains throughout the study period. Analyses of variance (ANOVAs) were performed to assess differences among years, seasons and domains.

### **Breeding success**

Throughout the breeding seasons of 2008-2011 and 2016-2018, in Selvagem Grande, between 296 and 349 breeding pairs of Cory’s shearwaters were followed each year as part of a long-term demographic study. Successes and failures were registered and the hatching success (percentage of laid eggs that hatched; Cory’s shearwaters lay one single egg per year), fledging success (percentage of hatched chicks that survived to fledging age), and the breeding success (hatching success  $\times$  fledging success = percentage of chicks fledged per egg laid) were calculated as a measure of reproductive performance.

## **Results**

### *Foraging ecology of the Cory’s shearwater*

We counted a total of 2775 prey individuals in regurgitations, 79% of which were successfully identified to species, genus or family level, resulting in 47 identified species/genus of 32 identified families (see Supplementary Materials, Table A.1). Fish were the most consumed prey across all years, both in terms of number of individuals (NF range = 68.7-99.2%) and in occurrence (FO range = 87.1-100%). Cephalopods were always present throughout the study period (NF range = 0.8-26.7%; FO range = 10.0-50.0%) (see Supplementary Materials, Table A.1). Overall, chub mackerel was the most common prey found in the regurgitations of Cory’s shearwater from 2008 to 2011 and in 2016, followed

by Pilotfish *Naucrates ductor*, European pilchard *Sardina pilchardus* and Flying-neon squid *Ommastrephes bartramii* (Figure 2). In 2017, Longspine snipefish *Macroramphosus scolopax* appeared as the most numerous prey (NF=61.5%, Figure 2), occurring in 53% of the samples. This situation persisted in 2018 with snipefish dominating the diet (NF=95.3% and FO=87.5%). We found significant differences between years in the diet composition of this shearwater (PERMANOVA:  $F_{6,820} = 13.202$ ,  $r^2 = 0.08$ ,  $p < 0.01$ ).




Figure 2- Interannual variability (numeric frequency) of the most common prey of the Cory's shearwater *Calonectris borealis* from Selvagem Grande (2008, n=471; 2009, n=703; 2010, n=633; 2011, n=137; 2016, n=183; 2017, n=156; 2018, n=492). Detailed data in Supplementary Materials, Table A.1.

In 2017 and 2018 the Shannon-Wiener diversity index reduced abruptly to  $H=1.36 \pm 0.24$  and  $H=0.24 \pm 0.05$ , respectively, compared with the previous years' mean ranges:  $H=2.19-2.61$  in 2008-2011, and  $H=1.75 \pm 0.20$  in 2016.

The average standard length of the Longspine snipefish was 78.5mm (range: 51.7-115.9mm, n=516), with no differences among the 3 years (ANOVA:  $F_{2,512}=0.92$ ,  $p=0.39$ , See Supplementary materials, Figure A.1). Cory's shearwater fed on Pilotfish with standard lengths of 90.7mm (range: 24.3-225.6mm). See Supplementary materials for further information on the standard length of prey of Cory's shearwaters (Table A.2).

The destination of foraging trips also presented important differences among years (PERMANOVA:  $F_{5,300} = 11.73$ ,  $r^2 = 0.16$ ,  $p < 0.01$ , Figure 3). From 2009 to 2011, the African coast was the most used foraging destination ( $52 \pm 11\%$ ). However, in 2016, most trips targeted the Selvagem domain and its surroundings (FO=28% and FO=36%, respectively), and in 2018, the Pelagic domain was the most frequently used (FO=82%). Overall, trips in 2016 and 2018 were closer to the colony comparing to other years, presenting average maximum distances from the colony of 121.6 and 91.6km, respectively. Even though, 2016 presented a higher number of shorter trips than in 2018. Trips from 2016 and 2018 were significantly different from trips from 2010, 2011 and 2013 (ANOVA:  $F_{5,312} = 12.21$ ,  $p < 0.01$  followed by post-hoc test Tukey HSD, Figure 4).




Figure 3 - Percentage of use of different oceanic domains (FO%) by Cory's shearwaters *Calonectris borealis* during the chick-rearing phase of 2009-2011 ( $n=20$ ,  $34$ ,  $46$ ,  $28$ , respectively), 2016 ( $n=103$ ), 2018 ( $n=75$ ). Data from Weak Permanent Upwelling Region and Strong Permanent Upwelling Region were combined ("African coast").

The total annual landings of chub mackerels in the archipelago of Madeira from 2008 to 2018 showed no down trend that could imply a reduction in the availability of the species (Figure 5). It is worth noting that in 2018, when this fish almost disappeared from the diet of Cory's shearwaters, catch levels by human fisheries were not unusually low (Figure 5). The

correlation between the numeric frequency of chub mackerels and landings showed to be positive but not quite attaining statistical significance ( $r=0.63$ ,  $p=0.13$ ).




Figure 4 - Average (SD) maximum distance (Km) reached by Cory's shearwaters *Calonectris borealis*, in each trip per year.




Figure 5 - Variation in the total annual landings and mean catch value (224 tonnes, dashed line) of Atlantic chub mackerel *Scomber colias* in Madeira island for the decade of 2008-2018.

#### *Environmental and oceanographic characteristics of foraging areas*

Overall, the defined domains showed a similar annual pattern in the SST, Chl a and wind components, but their mean values differed, except the Selvagem and Pelagic domains

which were indistinguishable (see Supplementary Materials, Table A.3 and Figure A.2). No correlation was found between any of the oceanographic and meteorological (NAO) variables studied and the proportions of various prey in the diet of Cory's shearwaters (see Supplementary Materials, Table A.4) either in Winter or Summer.

### *Breeding success*

Breeding success only varied slightly during the studied period of 2008-2011 (72.1-75.4% in 2008-2010 and 63.8% in 2011) and 2016-2018 (71.3-76.7%).

## **Discussion**

Cory's Shearwaters are opportunistic feeders which used different foraging domains within our study area, ranging far from their breeding colony (Figure 1, see also Alonso et al., 2018). We took advantage of this to gather information on prey in these waters for a total of 7 years using the combination of GPS-tracking and diet data, which provided useful indications concerning the likely origin of their pelagic prey (Alonso et al., 2018). In 2016, the population of chub mackerel showed a peak of abundance in the waters surrounding Selvagens. More noticeably, since 2017 Longspine snipefish suddenly became the dominant prey-item of Cory's shearwaters, mostly captured in the pelagic domain adjacent to Selvagens islands. Fisheries data indicate that the dietary shift of this seabird was not associated with any noticeable decrease of chub mackerel in the region. Hence, our Cory's shearwater diet dataset strongly suggests a major shift in the epipelagic fish communities of this little-known region, with a sudden and dramatic expansion of a previously scarce species, the Longspine snipefish.

From 2008 to 2011, the main prey of Cory's shearwaters from Selvagem Grande colony was the chub mackerel. Chub mackerels are widely distributed in the NW African region, in deep-sea areas, seamounts, and continental and island shelves (Alonso et al., 2018; Pitcher et al., 2007; Stromme et al., 2006; Wienerroither et al., 2009). Such large distribution might explain its importance in the diet of Cory's shearwaters (Alonso et al., 2018). A positive correlation (which failed to reach significance, possibly due to the small number of sampled years) between the landings of chub mackerel and its numeric frequency supports the hypothesis that the diet of Cory's shearwaters may provide broad indication of the abundance of this species. However, it would be important to sample more years and for a longer period of time to validate this assumption. While it is difficult to know where chub mackerels are

most abundant, it is possible to state that they were very abundant in the Selvagem domain in 2016. In this year, Cory's shearwaters frequently foraged around Selvagem Grande (FO%=28%) and adjacent pelagic waters (FO%=36%) and fed mostly on chub mackerels (NF=55% and FO=70%). The peak in total annual landings of chub mackerels observed in Madeira (this study) and in the African coast (Morocco to Senegal, including the Canary islands, 400000 tonnes; FAO, 2018) in that same year fit well with the peak that we recorded around Selvagem islands. The increase of the population at that time is further confirmed by acoustic surveys performed along the African coast in 2016-2017 (FAO, 2019).

From 2009 to 2011, the African coast was the domain where Cory's shearwaters foraged the most. There, the community of forage fish is mainly composed by European pilchards, which represent 45% of catch contribution of small pelagic fish (FAO, 2018). From 2008 to 2010, there was a slight increase in the total annual catches of the European pilchard (FAO, 2018) which coincides with the increase of European pilchards in Cory's shearwater diet. In 2016, the proportion of pilchards in the diet did not follow the new increase of landings described in FAO (2018). This can be explained by a reduction in the use of the African coast by Cory's shearwaters due to the apparent increase of chub mackerel around Selvagem.

Pilotfish which are generally found in association with floating structures (juveniles) or large marine species (adults) in pelagic ecosystems (e.g. Riera et al. 1999), were also a common pelagic prey of Cory's shearwaters. Throughout the period of 2008-2011 and 2016-2017, this seabird preyed upon Pilotfish with sizes between 24 and 230mm, which correspond to juveniles of up to 5 month old (Vassilopoulou et al., 2005). The near-constant presence of this age-group in the diet of Cory's shearwaters during this period is an indicator that they were feeding on new recruits every year.

Besides fish, cephalopods were also an important prey group in the diet of the Cory's shearwater. In our study, 30% of the cephalopods found in the stomachs of this seabird were Neon-flying squid *Ommastrephes bartramii* or belonged to the *Ommastrephidae* family. This is the most common cephalopod family in the Canary Current System, and the Neon-flying squid is among the most common species off North Africa (Hastie et al., 2009). The small variation in the occurrence of flying squids in the diet of the Cory's shearwater in the study



years of 2008-2017 indicates a regular presence of this species in the pelagic waters off NW Africa.

In 2017, Cory's shearwaters from the Selvagem colony shifted their diet to feed mostly on Longspine snipefish. In the following year, the importance of this fish in the diet of this seabird further increased to 95.3%, and the pelagic areas around the archipelago of Selvagens replaced the Selvagem domain and adjacent waters as the main foraging grounds of the Cory's shearwater. The high abundance of the chub mackerel in the African coast and the Canary islands, in 2017 and 2018 (FAO, 2019) implies that the consumption of the Longspine snipefish was not a result of a shortage of the traditional prey, but a choice. Therefore, we conclude that the trophic shift observed for this seabird reflects an increase in the availability of Longspine snipefish in the pelagic waters around Selvagem in 2017/2018. Such an increase very close to the nesting colony probably allowed Cory's shearwaters to have access to high quantities of food with low effort, even if less caloric (Martins et al., 2004), rather than targeting alternative, more nutritious prey farther away, which agrees with the classic optimal foraging theory and central place foraging (Bartumeus and Catalan, 2009). Thus, should Longspine snipefish have been abundant close to the colony in previous years, then these seabirds would not have travelled to the African coast to forage. Despite the possible link between a change in the foraging area and a shift in prey consumed, the foraging behaviour of seabirds is known to be influenced by extrinsic factors that are related to prey availability and not by a random choice of foraging areas (Shealer, 2002). Furthermore, between 2016 and 2018, the foraging areas remained the same but with a shift in the diet of Cory's shearwaters from chub mackerels to snipefish, which suggests that the foraging area was not the reason for the shift in diet. For this reason, the shift in the diet is much more likely to be a result of the shift in the prey community of that specific area than solely a change in the foraging area.

Snipefish found in the regurgitations of Cory's shearwater had a mean length of  $78.5 \pm 59.1$  mm, which did not vary during the study years. These sizes correspond to snipefish slightly younger than one year (Ehrich, 1976; Lopes and Farinha, 1996), possibly resulting from the spawning of December-March (Lopes and Farinha, 1996). The yearly occurrence of snipefish with these standard lengths in the diet of the Cory's shearwater allows us to conclude that the birds were preying on new recruits every year which, consequently, leads

us to hypothesize that snipefish were reproducing successfully in the region from 2016 to 2018.

The distribution of the Longspine snipefish in the North Atlantic ranges in latitude from the Iberian Peninsula down to North Mauritania, including varied environments such as shelves and upper slopes, and seamounts (Ehrich et al., 1987). Marked fluctuations in Longspine snipefish abundance have been recorded in other regions of the NE Atlantic. On the Portuguese continental shelf, the only area monitored regularly through systematic surveys (Borges, 2000; Lopes et al., 2006; Lopes and Farinha, 1996; Marques et al., 2005), the snipefish population presented high abundances in the 1970's (annual landings: 10,000 tonnes in 1973, and 33,000 tonnes in 1978) (Morais, 1981), and late 1990's/ early 2000's (acoustic estimates: 500 000 tonnes in 1998 to 175 000 tonnes in 2003) (Marques et al., 2005), with low abundances in the 1980's (Marques et al., 2005). Seemingly, these fluctuations coincided with abrupt fluctuations in other areas. Outbursts were described in the coast of Morocco (Arístegui et al., 2004) and in the Azores (Granadeiro et al., 1998; Ramos et al., 1998) in the 70's (acoustic survey: 1,000,000 tonnes in 1976 to disappear in the 80's) and the 90's (1994-1995), respectively. In 2005, dietary data from Cory's shearwaters provided some evidence of the decrease in abundance of snipefish in the Azores comparing with the 90's (Xavier et al., 2011). The Longspine snipefish was considered abundant in the Seine Seamount (located northeast of Madeira) in 2003-2005, and in the Meteor Seamount (located in the middle of the Atlantic, west of Madeira) in 1967, 1970 and 1998 (Christiansen et al., 2009; Fock et al., 2002b; Zidowitz and Fock, 2004).

Studies have shown that the appearance and disappearance of Longspine snipefish throughout time can alter considerably the trophic web, with predators suddenly shifting their diet to this prey once its numbers increase (e.g. Morato et al. 1999; Silva 1999). Snipefish is an important prey (at some times and locations the main prey) of seabirds, sharks, dolphins, tunas, rays, hakes and John Dory *Zeus faber* (e.g. Granadeiro et al. 1998; Morato et al. 1999, 2003; Silva 1999; Zidowitz and Fock 2004). To what extent snipefish can influence the functioning of the ecosystems when their numbers change is still uncertain. In our study, Cory's shearwaters, which like many seabird species are sensitive to prey fluctuations (Becker et al., 2007; Kowalczyk et al., 2014), did not show signs of having been affected by this major shift in their main prey. Despite a radical shift in diet, between 2016 and 2018, the

breeding success remained largely unchanged. This also suggests that snipefish can be a valuable resource for pelagic predators.

What drives the Longspine snipefish wide population fluctuations is still unknown, and we did not find changes in oceanographic variables that correlated with the increase of this species' abundance reported here. The sudden proliferation of some marine taxa can sometimes be associated with the disappearance of a dominant species (predators or competitors) due to fisheries or environmental changes (Caddy and Rodhouse, 1998; Gulland and Garcia, 1984). The increase of snipefish in our study was apparently not linked with the decrease of any other abundant species, according to fisheries data (FAO, 2019). So, the reasons for such successful consecutive recruitments is yet to be discovered.

The Cory's shearwater is considered to be a generalist predator, but it is important to point out that Cory's shearwaters are surface feeders and shallow divers, reaching an average depth during foraging trips of only 2.7m (Mougin and Mougin, 1998). Hence, they cannot be considered totally unbiased samplers of the whole marine environment. Their diet will thus reflect availability in surface waters, rather than overall abundance of fish stocks. Nevertheless, it is reasonable to assume that availability in surface waters correlates with overall abundance, even if imperfectly so. The action of subsurface predators can bring more prey to surface waters, and should their abundance drastically change, this could have an important influence in the diet of Cory's shearwaters. However, we know of no evidence for drastic changes in the community of underwater predators which might explain the very major shift in the diet of the Cory's shearwaters observed in this study.

This study exemplifies how biological samplers like Cory's shearwaters can provide novel information about important changes in epipelagic ecosystems, that would otherwise be difficult to detect. Previous studies have already shown the usefulness of seabirds as bioindicators of their foraging areas and prey, unveiling annual fluctuations of fish stocks (e.g. Scopel et al., 2018) and changes in the food web and in oceanographic conditions (e.g. Montevecchi, 2007). The major shifts in the epipelagic fish community around Selvagens archipelago revealed by this study and the lack of known environmental correlates for such shifts underscore our poor understanding of the dynamics of key-species in little known subtropical pelagic waters. Such trends and variations need to be better monitored and

understood if we are to measure the impact of ongoing global changes and to sustainably manage the broad marine environment and resources.

## **Acknowledgements**

This work was supported by Fundação para a Ciência e a Tecnologia (FCT Portugal) through the project Oceanwebs (PTDC/MAR-PRO/0929/2014) and Seamigrant (PTDC/BIA-ANM/3743/2014), the strategic projects UIDB/04292/2020 and UIDP/04292/2020 granted to MARE, and UIDB/50017/2020 and UIDP/50017/2020 granted to CESAM. PD/BD/127810/2016 was granted to JR by FCT. We would like to acknowledge Serviço do Parque Natural da Madeira and Instituto das Florestas e Conservação da Natureza, specially Dr. Paulo Oliveira and Dr. Carolina Santos, for providing the required permits to undertake experiments with birds and logistical support to carry out the work at the islands of Selvagens. All procedures were carried out following a positive recommendation from the Animal Welfare Regulatory Body of the Faculty of Sciences of Lisbon University and were carried out under licenses issued by the Direção Geral de Alimentação e Veterinária, license number DGAV 0421/2016. Thanks are due to Direção Regional de Pescas da Madeira, specially to Dr. Graça Faria, for providing data on the annual landings of small pelagic fishes. Filipe Moniz for helping during the field and laboratory work. We also thank to Letizia Campioni, Diana Barão, Teresa Catry, Maria Alho and Marie Claire for helping during field work and giving further insights for this study, and to park-wardens Carlos Clemente, Carlos Santos, Jaques da Mata, João Paulo Mendes, Maurício Baptista, Marco Camacho and Fernando. No conflicts of interest.

## **References**

- Alonso, H., Granadeiro, J.P., Dias, M.P., Catry, T., Catry, P., 2018. Fine-scale tracking and diet information of a marine predator reveals the origin and contrasting spatial distribution of prey. *Prog. Oceanogr.* 162, 1–12. <https://doi.org/10.1016/j.pocean.2018.02.014>
- Alonso, H., Granadeiro, J.P., Paiva, V.H., Dias, A.S., Ramos, J.A., Catry, P., 2012. Parent-offspring dietary segregation of Cory's shearwaters breeding in contrasting environments. *Mar. Biol.* 159, 1197–1207. [https://doi.org/10.1007/s00227-012-1900-](https://doi.org/10.1007/s00227-012-1900-2)

- Alonso, H., Granadeiro, J.P., Waap, S., Xavier, J., Symondson, W.O.C., Ramos, J.A., Catry, P., 2014. An holistic ecological analysis of the diet of Cory's shearwaters using prey morphological characters and DNA barcoding. *Mol. Ecol.* 23, 3719–3733. <https://doi.org/10.1111/mec.12785>
- Arístegui, J., Álvarez-Salgado, X.A., Barton, E.D., Figueiras, F.G., Hernández-León, S., Roy, C., Santos, A.M.P., 2004. Oceanography and Fisheries of the Canary Current/Iberian Region of the Eastern North Atlantic (18a, E), in: *The Sea*. pp. 877–932.
- Bartumeus, F., Catalan, J., 2009. Optimal search behavior and classic foraging theory. *J. Phys. A: Math. Theor.* 42: 12pp. <https://doi.org/10.1088/1751-8113/42/43/434002>
- Becker, B.H., Peery, M.Z., Beissinger, S.R., 2007. Ocean climate and prey availability affect the trophic level and reproductive success of the marbled murrelet, an endangered seabird. *Mar. Ecol. Prog. Ser.* 329:267–279. <https://doi.org/10.3354/meps329267>
- Borges, L., 2000. Age and growth of the snipefish, *Macrorhamphosus spp.*, in the Portuguese continental waters. *J. Mar. Biol. Assoc. United Kingdom* 80, 147–153. <https://doi.org/10.1017/S0025315499001666>
- Cabral, H.N. and Murta, A.G., 2002. The diet of blue whiting, hake, horse mackerel and mackerel off Portugal. *J. Appl. Ichthyol.*, 18:14–23.
- Caddy, J.F., Rodhouse, P.G., 1998. Cephalopod and groundfish landings: evidence for ecological change in global fisheries? *Rev. Fish Biol. Fish.* 8, 431–444. <https://doi.org/10.1023/A:1008807129366>
- Calenge, C., 2015. Home Range Estimation in R: the adehabitatHR Package. R vignette. <https://doi.org/10.1111/j.1365-2656.2006.01186.x>
- Christiansen, B., Martin, B., Hirsch, S., 2009. The benthopelagic fish fauna on the summit of Seine Seamount, NE Atlantic: Composition, population structure and diets. *Deep. Res. Part II Top. Stud. Oceanogr.* 56, 2705–2712. <https://doi.org/10.1016/j.dsr2.2008.12.032>
- Church, G.E., Furness, R.W., Tyler, G., Gilbert, L., Votier, S.C., 2018. Change in the North Sea ecosystem from the 1970s to the 2010s: great skua diets reflect changing forage fish, seabirds, and fisheries. *ICES J. Mar. Sci.* <https://doi.org/10.1093/icesjms/fsy165>

- Cropper, T.E., Hanna, E., Bigg, G.R., 2014. Spatial and temporal seasonal trends in coastal upwelling off Northwest Africa, 1981-2012. Deep. Res. Part I Oceanogr. Res. Pap. 86, 94–111. <https://doi.org/10.1016/j.dsr.2014.01.007>
- Davenport, R., Neuer, S., Hernandez-Guerra, A., Rueda, M.J., Llinas, O., Fischer, G., Wefer, G., 1999. Seasonal and interannual pigment concentration in the Canary Islands region from CZCS data and comparison with observations from the ESTOC. Int. J. Remote Sens. 20, 1419–1433. <https://doi.org/10.1080/014311699212803>
- Delgado, J., Amorim, A., Gouveia, L., Gouveia, N., 2018. An Atlantic journey: The distribution and fishing pattern of the Madeira deep sea fishery. Reg. Stud. Mar. Sci. <https://doi.org/10.1016/j.rsma.2018.05.001>
- Edwards, M., Beaugrand, G., Helaouët, P., Alheit, J., Coombs, S., 2013. Marine Ecosystem Response to the Atlantic Multidecadal Oscillation. PLoS One 8, 1–5. <https://doi.org/10.1371/journal.pone.0057212>
- Ehrich, S., 1976. Zur Taxonomie, Ökologie und Wachstum von *Macroramphosus scolopax* (Linnaeus, 1758) (Pisces, Syngnathiformes) aus dem subtropischen Nordostatlantik. Berichte der Dtsch. wissenschaftlichen Kommission für Meeresforsch. 24, 251–266.
- Ehrich, S., John, H.-C., Westhaus-Ekau, P., 1987. Southward extension of the reproductive range of *Macroramphosus scolopax* in the upwelling area off North-West Africa. South African J. Mar. Sci. 5, 95–105. <https://doi.org/10.2989/025776187784522298>
- FAO, 2019. Report of the FAO Working Group on the Assessment of Small Pelagic Fish off Northwest Africa. Banjul, the Gambia, 26 June–1 July 2018. FAO Fisheries and Aquaculture Report No. R1247. Rome. Licence: CC BY-NC-SA 3.0 IGO. Downloaded from: <http://www.fao.org/publications/card/en/c/CA4121B>
- FAO, 2018. Fishery Committee for the Eastern Central Atlantic, Status summary for small pelagic stocks in the northern area of the eastern central Atlantic - CECAF. Abidjan, Côte d'Ivoire, 23–26 October 2018. FAO Fisheries and Aquaculture Report. CECAF/SSCVIII/2018/2. Downloaded from: <http://www.fao.org/fi/static-media/MeetingDocuments/CECAF/CECAF-SSC8/default.htm>
- Fock, H.O., Matthiessen, B., Zidowitz, H., Westernhagen, H. V., 2002a. Diel and habitat-dependent resource utilisation by deep-sea fishes at the Great Meteor seamount: niche

- overlap and support for the sound scattering layer interception hypothesis. *Mar. Ecol. Prog. Ser.* 244, 219–233. <https://doi.org/10.3354/meps244219>
- Fock, H.O., Uiblein, F., Von Westernhagen, H., 2002b. Biodiversity and species–environment relationships of the demersal fish assemblage at the Great Meteor Seamount (subtropical NE Atlantic), sampled by different trawls. *Mar. Biol.* 141, 185–199. <https://doi.org/10.1007/s00227-003-1281-7>
- Granadeiro, J.P., Monteiro, L.R., Furness, R.W., 1998a. Diet and feeding ecology of Cory’s shearwater *Calonectris diomedea* in the Azores, north-east Atlantic. *Mar. Ecol. Prog. Ser.* 166, 267–276. <https://doi.org/10.3354/meps166267>
- Granadeiro, J.P., Nunes, M., Silva, M. C., Furness, R. W., 1998b. Flexible foraging strategy of Cory’s shearwater, *Calonectris diomedea*, during the chick-rearing period. *Animal Behaviour*. 56, 1169–1176.
- Granadeiro, J.P., Silva, M.A., 2000. The use of otoliths and vertebrae in the identification and size-estimation of fish in predator-prey studies. *Cybium* 24, 383–393.
- Grebmeier, J.M., Overland, J.E., Moore, S.E., Farley, E. V., Carmack, E.C., Cooper, L.W., Frey, K.E., Helle, J.H., McLaughlin, F.A., McNutt, S.L., 2006. A major ecosystem shift in the northern Bering sea. *Science*. 311, 1461–1464. <https://doi.org/10.1126/science.1121365>
- Gulland, J.A., Garcia, S., 1984. Observed Patterns in Multispecies Fisheries, in: *Exploitation of Marine Communities*. Springer, Heidelberg, Berlin, pp. 155–190. [https://doi.org/10.1007/978-3-642-70157-3\\_7](https://doi.org/10.1007/978-3-642-70157-3_7)
- Hastie, L., Pierce, G., Wang, J., Bruno, I., Moreno, A., Piatkowski, U., Robin, J., 2009. Cephalopods in the North-eastern Atlantic: Species, biogeography, ecology, exploitation and conservation, in: *Oceanography and Marine Biology: An Annual Review*. pp. 111–190. <https://doi.org/10.1201/9781420094220.ch3>
- Hermida, M., Delgado, J., 2016. High trophic level and low diversity: Would Madeira benefit from fishing down? *Mar. Policy* 73, 130–137. <https://doi.org/10.1016/j.marpol.2016.07.013>
- Kowalczyk, N., Chiaradia, A., Preston, T., Reina, R.D., 2014. Linking dietary shifts and reproductive failure in seabirds: A stable isotope approach. *Functional Ecology*, 28:755–765. <https://doi.org/10.1111/1365-2435.12216>

- Lopes, M., Murta, A.G., Cabral, H.N., 2006. The ecological significance of the zooplanktivores, snipefish *Macroramphosus spp.* and boarfish *Capros aper*, in the food web of the south-east North Atlantic. J. Fish Biol. 69, 363–378. <https://doi.org/10.1111/j.1095-8649.2006.01093.x>
- Lopes, P.C., Farinha, A., 1996. Occurrence of the eggs and larvae of the snipefish *Macroramphosus scolopax* in the western coast of the Iberian Peninsula. Int. Counc. Explor. Sea (CM Pap. Reports) CM 1996/L.; 1–10.
- Marques, V., Chaves, C., Morais, A., Cardador, F., Stratoudakis, Y., 2005. Distribution and abundance of snipefish (*Macroramphosus spp.*) off Portugal (1998–2003). Sci. Mar. 69, 563–576. <https://doi.org/10.3989/scimar.2005.69n4563>
- Martins, I., Pereira, J.C., Ramos, J.A., Jørgensen, S.E., 2004. Modelling the effects of different quality prey fish species and of food supply reduction on growth performance of Roseate Tern chicks. Ecol. Modell. 177, 95–106. <https://doi.org/10.1016/j.ecolmodel.2004.02.002>
- Matallanas, J., 1982. Aspectos generales del regimen alimentario de *Macroramphosus scolopax* (Linnaeus 1758) (Pisces Macroramphosidae) en las costas catalanas (Mediterraneo occidental). Cah.Biol.Mar. 243–252.
- Montevecchi W.A., 2007. Binary dietary responses of northern gannets *Sula bassana* indicate changing food web and oceanographic conditions. Mar. Ecol. Prog. Ser. 352:213–220. <https://doi.org/10.3354/meps07075>
- Morais, R., 1981. Sobre a pescaria e biologia do apara-lápis ou trombeteiro. Bol. Inst. Nac. Invest. Pescas 6, 5–35.
- Morato, T., Solà, E., Grós, M., Menezes, G., 1999. Diets of forkbeard (*Phycis phycis*) and conger eel (*Conger conger*) off the Azores during spring of 1996 and 1997. Archipelago Life and M, 51–64.
- Morato, T., Solà, E., Grós, M.P., Menezes, G., 2003. Diets of thornback ray (*Raja clavata*) and tope shark (*Galeorhinus galeus*) in the bottom longline fishery of the Azores, northeastern Atlantic. Fish. Bull. 101, 590–602.
- Morato, T., Varkey, D.A., Damaso, C., Machete, M., Santos, M., Prieto, R., Santos, R.S., Pitcher, T.J., 2008. Evidence of a seamount effect on aggregating visitors. Mar. Ecol. Prog. Ser. 357, 23–32. <https://doi.org/10.3354/meps07269>



- Mougin, J.L., Mougin M.C., 1998. Maximum diving depths of Cory's shearwater in the course of its feeding trips during incubation. *Revue D Ecologie-La Terre Et La Vie* 53: 69–76.
- Oksanen, J., F. Guillaume Blanchet, R.K., Legendre, P., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., M. Henry H. Stevens, H.W., 2019. Community Ecology Package. R vignette.
- Palumbi, S.R., 1996. Nucleic acids II: the polymerase chain reaction, in: *Molecular Systematics*. pp. 205–247
- Pebesma EJ, Bivand RS (2005). Classes and methods for spatial data in R. *R News*, 5(2), 9–13. <https://CRAN.R-project.org/doc/Rnews/>.
- Pethybridge, H.R., Young, J.W., Kuhnert, P.M., Farley, J.H., 2015. Using stable isotopes of albacore tuna and predictive models to characterize bioregions and examine ecological change in the SW Pacific Ocean. *Prog. Oceanogr.* 134, 293–303. <https://doi.org/10.1016/j.pocean.2015.03.001>
- Phillips, R.A., Xavier, J.C., Croxall, J.P., 2003. Effects of Satellite Transmitters on Albatrosses and Petrels. *Auk* 120, 1082–1090.
- Pitcher, T.J., Morato, T., Hart, P.J.B., Clark, M.R., Haggan, N., Santos, R.S., 2007. *Seamounts: Ecology, Fisheries & Conservation*, Blackwell Publishing. <https://doi.org/10.1002/9780470691953>
- Ramos, J.A., Solá, E., Monteiro, L.R., Ratcliffe, N., 1998. Prey delivered to roseate tern chicks in the Azores. *J. F. Ornithol.* 69, 419–429.
- Riera, F., Grau, A., Grau, A.M., Pastor, E., Quetglas, A., Pou, S., 1999. Ichthyofauna associated with drifting floating objects in the Balearic Islands (western Mediterranean). *Sci. Mar.* 63, 229–235. <https://doi.org/10.3989/scimar.1999.63n3-4239>
- Salvanes, A.G.V., Kristoffersen, J.B., 2001. Mesopelagic Fishes, in: *Encyclopedia of Ocean Sciences*. Academic Press, pp. 1711–1717. <https://doi.org/10.1016/B978-012374473-9.00012-6>
- Shealer, D.A., 2002. Foraging behaviour and food of seabirds, in: *Biology of Marine birds*. CRC Press, pp. 137-177.

- Silva, A., 1999. Feeding habits of John Dory, *Zeus faber*, off the Portuguese continental coast. J. Mar. Biol. Assoc. United Kingdom 79, 333–340. <https://doi.org/10.1017/S002531549800037X>
- Stromme, T., Charouki, N., Alvheim, O., 2006. Survey of the pelagic fish resources off North West Africa, Part III: Cape Blanc – Cape Sim.
- Thiaw, M., Auger, P.-A., Ngom, F., Brochier, T., Faye, S., Diankha, O., Brehmer, P., 2017. Effect of environmental conditions on the seasonal and inter-annual variability of small pelagic fish abundance off North-West Africa: The case of both Senegalese sardinella. Fish. Oceanogr. 26, 583–601. <https://doi.org/10.1111/fog.12218>
- Vassilopoulou, V., Siapatis, A., Christides, G., Bekas, P., 2005. The biology and ecology of juvenile pilotfish (*Naucrates ductor*) associated with Fish Aggregating Devices (FADs) in eastern Mediterranean waters. Mediterr. Mar. Sci. 5, 61–70.
- Veit, R.R. and Harrison, N.M., 2017. Positive interactions among foraging seabirds, marine mammals and fishes and implications for their conservation. Front. Ecol. Evol., 5:121. <https://doi.org/10.3389/fevo.2017.00121>
- Velarde, E., Tordesillas, M. de la S., Rocio Esquivel, L.V., 1994. Seabirds as indicators of important fish populations in the Gulf of California. CalCofi Rep. 35, 137–143.
- Waap, S., Symondson, W.O.C., Granadeiro, J.P., Alonso, H., Serra-Gonçalves, C., Dias, M.P., Catry, P., 2017. The diet of a nocturnal pelagic predator, the Bulwer's petrel, across the lunar cycle. Sci. Rep. 7. <https://doi.org/10.1038/s41598-017-01312-3>
- Wienerroither, R., Uiblein, F., Bordes, F., Moreno, T., 2009. Composition, distribution, and diversity of pelagic fishes around the Canary Islands, Eastern Central Atlantic. Mar. Biol. Res. 5, 328–344. <https://doi.org/10.1080/17451000802478055>
- Wilson, R., 1984. An improved stomach pump for penguins and other seabirds. J. F. Ornithol. 55, 109–112.
- Xavier, J.C., Magalhães, M., Mendonça, A.S., Antunes, M., Carvalho, N., Machete, M., Santos, R.S., Paiva, V.H., Hamer, K.C., 2011. Changes in diet of Cory's shearwaters *Calonectris diomedea* breeding in the Azores. Mar. Ornithol. 39, 129–134. <https://doi.org/10.3354/meps166267>

Zidowitz, H., Fock, H.O., 2004. Trophic ecology of *Zenopsis conchifer* (Lowe, 1852) at the Great Meteor Seamount (subtropical North-east Atlantic). Arch. Fish. Mar. Res. 51, 279–286.

## Supplementary Materials

Table A.2 - Numeric frequency (NF%, number of prey in parenthesis) and Frequency of occurrence (FO%, number of samples in parenthesis) of prey species found in the regurgitations of Cory's shearwaters *Calonectris borealis* during the chick rearing season of 2008-2011 and 2016-2018, in Selvagem Grande island.

Prey Species	NF%							FO%						
	2008 (471)	2009 (703)	2010 (633)	2011 (137)	2016 (183)	2017 (156)	2018 (492)	2008 (199)	2009 (278)	2010 (194)	2011 (38)	2016 (48)	2017 (30)	2018 (40)
<b>TELEOSTEI</b>	<b>84.1</b>	<b>72.3</b>	<b>68.7</b>	<b>83.2</b>	<b>88.5</b>	<b>84.6</b>	<b>99.2</b>	<b>89.9</b>	<b>87.1</b>	<b>89.7</b>	<b>97.4</b>	<b>95.8</b>	<b>90</b>	<b>100</b>
<i>Belonidae</i>														
<i>Belone belone</i>				1.5	0.5						5.3	2.1		
<i>Belone sp</i>	0.8	0.1	1.4					2.0	0.4	3.6				
<i>Caproidae</i>														
<i>Capros aper</i>			0.2							0.5				
<i>Carangidae</i>														
<i>Naucrates ductor</i>	14.9	14.4	9.6	19.7	6.6	5.1		15.1	16.9	13.4	28.9	4.2	6.7	
<i>Trachurus picturatus</i>	0.2	0.1			1.6		1.8	0.5	0.4			4.2		12.5
<i>Trachurus sp</i>	1.1	1.7	3.5	1.5	4.4			2.5	3.2	4.6	5.3	8.3		
<i>Unidentified</i>	0.2	0.4	0.5					0.5	1.1	1.0				
<i>Centriscidae</i>														
<i>Macroramphosus scolopax</i>			0.3		2.2	61.5	95.3			1.0		4.2	53.3	87.5
<i>Clupeidae</i>														
<i>Sardina pilchardus</i>	3.8	4.0	11.7	13.1	6.0	0.6		7.5	8.6	20.1	18.4	10.4	3.3	
<i>Sardinella sp</i>		0.1							0.4					
<i>Unidentified</i>	0.4	0.4	0.6					1.0	1.1	2.1				
<i>Coryphaenidae</i>														
<i>Coryphaena equiselis</i>		0.1			0.5				0.4			2.1		
<i>Coryphaena hippurus</i>				0.7							2.6			

<i>Coryphaena sp</i>	0.2		0.2				0.5		0.5		
Diretmidae											
<i>Diretmus argenteus</i>			0.6						2.1		
Engraulidae											
<i>Engraulis encrasicolus</i>	0.8	3.6	0.5	2.2			1.5	3.2	1.0	5.3	
Exocoetidae	7.6	4	2.7	0.7	1.6		13.1	7.1	7.7	7.9	6.3
<i>Cheilopogon exsiliens</i>		0.4						1.1			
<i>Cheilopogon melanurus</i>		0.1						0.4			
<i>Cheilopogon pinnatibarbatus</i>		0.3						0.7			
<i>Cheilopogon sp</i>	0.6	0.3	0.3				1.5	0.7	1.0		
<i>Exocoetus volitans</i>	0.6				1.1		1.5				4.2
<i>Exocoetus sp</i>	3.4	0.4	1.6	0.7			7.0	1.1	5.2	2.6	
Unidentified	3.0	2.4	0.8		0.5		6.0	4.0	2.6		2.1
Halosauridae											
Unidentified	0.6	0.1	0.2	0.7			1.0	0.4	0.5	2.6	
Microstomatidae											
<i>Nansenia sp</i>				0.7						2.6	
Molidae											
<i>Ranzania laevis</i>			1.1						3.6		
Myctophidae											
<i>Diaphus splendidus</i>		0.1						0.4			
<i>Lampadena luminosa</i>			0.2						0.5		
Unidentified	0.6		0.8	1.5	0.5	1.9	1.5		2.1	5.3	2.1 6.7
Neoscopelidae											
<i>Neoscopelus macrolepidotus</i>	0.2			1.5			0.5			2.6	
Nettastomatidae											
<i>Nettastoma melanurum</i>				0.7						2.6	
Opisthoproctidae											
<i>Opisthoproctus soleatus</i>				0.7						2.6	

Scombridae														
<i>Auxis rochei</i>		0.6					0.4		1.4					5.0
<i>Katsuwonus pelamis</i>	0.2	0.6	0.9					0.5	1.4	3.1				
<i>Scomber colias</i>	16.6	16.1	4.6	9.5	32.2	10.3	1.6	23.6	29.5	10.8	23.7	54.2	46.7	20.0
<i>Scomber sp</i>	19.7	18.2	16.3	11.7	21.9	0.6		30.2	28.4	30.4	26.3	45.8	3.3	
<i>Scomber</i>	36.3	34.3	20.9	21.2	54.1	10.9	1.6	45.7	47.5	35.1	39.5	68.8	50.0	20.0
Scomberesocidae														
<i>Scomberesox saurus</i>						0.6								3.3
<i>Scomberesox sp</i>	4.2	0.4	2.7		1.1			3.0	1.1	4.6		2.1		
Sparidae														
<i>Boops boops</i>		0.1	0.5		4.9				0.4	1.0		4.2		
<i>Dentex macrophthalmus</i>				7.3							5.3			
Sternoptuchidae														
<i>Argyropelecus aculeatus</i>	0.4							0.5						
Synaphobranchidae														
<i>Unidentified</i>	0.4	0.3	1.3	2.2				1.0	0.7	3.6	7.9			
Trichiuridae														
<i>Unidentified</i>	1.9	0.9	1.6		0.5	0.6		4.5	2.2	4.6		2.1	3.3	
Unidentified Fish	8.9	5.8	7.0	7.3	3.8	3.2		19.6	14.4	20.6	23.7	12.5	13.3	
<b>CEPHALOPOD</b>	<b>13.6</b>	<b>26.7</b>	<b>20.1</b>	<b>16.8</b>	<b>9.3</b>	<b>14.7</b>	<b>0.8</b>	<b>26.6</b>	<b>36.7</b>	<b>45.9</b>	<b>50</b>	<b>27.1</b>	<b>33.3</b>	<b>10</b>
Chiroteuthidae														
<i>Chiroteuthis sp</i>		0.4							1.1					
Cranchiidae														
<i>Taonius pavo</i>	0.2		0.5					0.5		1.0				
Grimalditeuthidae														
<i>Grimalditeuthis bonplandi</i>	0.2							0.5						
Histiotteuthidae														
<i>Histioteuthis arcturi</i>	0.4	0.4	1.9					1.0	1.1	5.7				
<i>Histioteuthis meleanogroteuthis</i>			0.2						0.7	0.5				

<i>Histioteuthis sp</i>	0.4	0.3						1.0						
<i>Unidentified</i>				3.6	1.6					13.2	6.3			
Mastigoteuthidae														
<i>Mastigoteuthis sp</i>		0.1							0.4					
<i>Unidentified</i>			0.6							2.1				
Neuteuthidae														
<i>Neoteuthis sp</i>			0.2							0.5				
Octopoteuthidae														
<i>Taningia danae</i>	0.2	0.1						0.5	0.4					
Ommastrephidae	4.5	10.2	2.4	4.5	1.1	9.0		9.5	15.1	6.2	18.4	4.2	16.7	
<i>Ommastrephes bartramii</i>	3.8	10.0	2.4	1.5		9.0		8.0	14.7	6.2	5.3		16.7	
<i>Unidentified</i>	0.6	0.3		5.1	1.1			1.5	0.7		15.8	4.2		
Onychoteuthidae														
<i>Ancistrotheuthis lichtensteinii</i>		0.1							0.4					
Sepiidae														
<i>Unidentified</i>			0.2							0.5				
Unidentified Cephalopod	7.6	14.9	14.2	6.6	6.6	5.8	0.8	15.6	26.3	35.1	23.7	22.9	20.0	10.0
<b>CRUSTACEA</b>	<b>2.3</b>	<b>0.9</b>	<b>11.2</b>	<b>0</b>	<b>2.2</b>	<b>0.6</b>	<b>0</b>	<b>2.5</b>	<b>1.8</b>	<b>14.4</b>	<b>0</b>	<b>6.3</b>	<b>3.3</b>	<b>0</b>
Decapoda			0.3							0.5				
Isopoda		0.6	9.8		1.1				1.1	10.8		4.2		
Unidentified Crustacea	2.3	0.3	1.1		1.1	0.6		2.5	0.7	3.1		4.2	3.3	




Figure A.1 - Size distribution of the Longspine snipefish *Macroramphosus scolopax* captured by the Cory's shearwater *Calonectris borealis* of Selvagem Grande breeding colony from 2016 to 2018 (2016,  $n=4$ ; 2017,  $n=96$ ; 2018,  $n=469$ ).

Table A.3 - Mean standard length ( $\pm$ SD) and range values (in mm) of each prey species and sampling years of Cory's shearwater *Calonectris borealis* breeding in Selvagem Grande.

Prey	Year	N	Mean	SD	Range
<b>Fish</b>					
<i>Belone belone</i>	2016	1	421.2		
<i>Belone</i> sp.	2008	1	415.7		
	2009	1	408.6		
	2010	6	402.1	13.9	383.9-422.8
<i>Boops boops</i>	2009	1	205.5		
	2010	3	124.6	44.7	93.6-175.9
<i>Capros aper</i>	2010	1	57.4		
<i>Cheilopogon exsiliens</i>	2009	3	268.2	12.8	254.0-279.0
<i>Cheilopogon malanurus</i>	2009	1	212.8		
<i>Cheilopogon pinnatibarbatulus</i>	2009	1	249.8		
<i>Cheilopogon</i> sp.	2008	3	205.9	71.9	128.2-270.0
	2009	2	226.3	32.2	203.5-249.0
	2010	2	124.7	41.9	95.1-154.3
<i>Engraulis encrasicolus</i>	2008	2	128.7	4.9	125.2-132.2
	2009	20	141.6	8.1	125.2-156.2
	2010	2	140.4	10.9	132.7-148.1
<i>Exocoetidae</i>	2008	8	219.7	28.9	152.6-249.8
	2009	11	194.1	37.6	114.0-226.7
	2010	3	234.1	18.5	213.7-249.8
<i>Exocoetus</i> sp.	2008	14	216.1	5.3	209.0-228.2
	2009	3	219.8	10.2	221.0-231.0
	2010	9	229.9	12.9	213.7-249.8
	2016	2	214.3	2.6	212.4-216.1



<i>Exocoetus volitans</i>	2008	2	225.7	7.9	220.0-231.3
	2016	2	214.3	2.6	212.4-216.1
<i>Macroramphosus scolopax</i>	2010	1	115.9		
	2016	4	80.8	4.1	78.8-86.9
	2017	75	79.7	10.2	54.6-103.2
	2018	436	78.2	9.2	51.7-111.7
<i>Naucrates ductor</i>	2008	61	103.0	27.2	51.8-225.6
	2009	94	89.2	29.2	24.3-168.4
	2010	59	78.5	27.9	35.7-157.0
	2011	24	97.9	32.9	28.4-204.3
	2016	11	98.8	22.7	60.6-133.1
	2017	6	64.4	14.0	48.9-81.9
<i>Sardina pilchardus</i>	2008	10	192.6	31.5	130.0-234.3
	2009	16	199.3	30.0	135.5-225.4
	2010	50	164.2	23.6	112.1-252.0
	2011	13	123.4	32.8	73.4-185.9
	2016	9	128.0	41.4	89.9-179.2
<i>Scomber</i> sp.	2008	148	185.2	30.4	89.2-270.1
	2009	209	208.1	21.5	162.8-273.3
	2010	102	197.2	18.3	155.0-253.1
	2011	21	214.0	20.7	187.3-270.4
	2016	80	193.7	27.8	155.3-272.5
	2017	9	209.6	31.2	151.4-243.1
	2018	4	227.3	26.4	207.9-266.1
<i>Trachurus picturatus</i>	2008	1	169.4		
	2009	1	217.0		
	2016	3	158.9	7.5	150.8-165.4
	2018	8	153.8	3.7	145.9-157.3
<i>Trachurus</i> sp.	2008	4	229.4	64.6	151.1-309.3
	2009	10	156.4	48.0	76.2-240.2
	2010	19	163.7	17.7	135.4-201.8
	2011	1	291.2		
	2016	4	162.9	19.7	145.9-189.8
<b>Cephalopods</b>					
<i>Grimalditeuthis bonplandi</i>	2008	1	82.2		
<i>Histioteuthis arcturi</i>	2008	2	59.7	18.8	46.4-73.0
<i>Ommastrephes bartramii</i>	2008	18	71.2	29.8	28.0-108.9
	2009	56	54.6	24.4	14.9-109.1
	2010	8	58.4	28.4	20.0-105.0
	2011	2	38.0	24.0	21.0-55.0
	2017	2	75.5	0.7	75.0-76.0
<i>Ommastrephidae</i>	2008	1	63.5		
	2009	1	65.6		
	2011	2	22.0	1.4	21.0-23.0
	2016	1	102.0		
<i>Taonius pavo</i>	2008	1	380.9		

Table A.3 - ANOVA results for differences in oceanographic variables among different domains (Selvagem, Pelagic, Seamounts, Strong Permanent Upwelling Region and Weak Permanent Upwelling Region) in Summer and Winter. Significant differences are marked in bold.


	Domains		Year		Domains x Year	
	ANOVA	p-value	ANOVA	p-value	ANOVA	p-value
<b>Summer</b>						
<i>Chlorophyll a</i>	F <sub>4,165</sub> =2379.879	<b>&lt;0.01</b>	F <sub>10,165</sub> =8.851	<b>&lt;0.01</b>	F <sub>40,165</sub> =1.797	<b>&lt;0.01</b>
<i>Sea Surface Temperature</i>	F <sub>4,165</sub> =30.589	<b>&lt;0.01</b>	F <sub>10,165</sub> =1.250	0.263	F <sub>40,165</sub> =0.239	1
<i>Wind speed</i>	F <sub>4,165</sub> =10.349	<b>&lt;0.01</b>	F <sub>10,165</sub> =3.088	<b>&lt;0.01</b>	F <sub>40,165</sub> =0.253	1
<i>Zonal velocity</i>	F <sub>4,165</sub> =19.337	<b>&lt;0.01</b>	F <sub>10,165</sub> =3.219	<b>&lt;0.01</b>	F <sub>40,165</sub> =0.427	0.998
<i>Meridional velocity</i>	F <sub>4,165</sub> =13.334	<b>&lt;0.01</b>	F <sub>10,165</sub> =3.448	<b>&lt;0.01</b>	F <sub>40,165</sub> =0.253	0.999
<b>Winter</b>						
<i>Chlorophyll a</i>	F <sub>4,165</sub> =715.005	<b>&lt;0.01</b>	F <sub>10,165</sub> =7.649	<b>&lt;0.01</b>	F <sub>40,165</sub> =0.691	0.915
<i>Sea Surface Temperature</i>	F <sub>4,165</sub> =31.637	<b>&lt;0.01</b>	F <sub>10,165</sub> =21.608	<b>&lt;0.01</b>	F <sub>40,165</sub> =0.761	0.844
<i>Wind speed</i>	F <sub>4,165</sub> =18.633	<b>&lt;0.01</b>	F <sub>10,165</sub> =5.253	<b>&lt;0.01</b>	F <sub>40,165</sub> =0.238	1
<i>Zonal velocity</i>	F <sub>4,165</sub> =8.202	<b>&lt;0.01</b>	F <sub>10,165</sub> =7.372	<b>&lt;0.01</b>	F <sub>40,165</sub> =0.368	1
<i>Meridional velocity</i>	F <sub>4,165</sub> =13.575	<b>&lt;0.01</b>	F <sub>10,165</sub> =4.944	<b>&lt;0.01</b>	F <sub>40,165</sub> =0.370	1

Table A.4 – Pearson's correlation between the oceanographic variables in the Pelagic domain (circle with a 160km radius around Selvagem and with depths between 2000 and 4000m, see text) and the numeric frequency of the main prey in the Cory's shearwater diet. Winter: January-April, Summer: June-September.


	Winter		Summer	
	Correlation	p-value	Correlation	p-value
<b>Chlorophyll a</b>				
<i>Chub mackerel</i>	-0.29	0.5	-0.05	0.9
<i>Cephalopods</i>	-0.38	0.4	-0.42	0.4
<i>Pilotfish</i>	-0.045	0.3	-0.52	0.2
<i>Longspine snipefish</i>	0.67	0.1	0.54	0.2
<b>Sea Surface Temperature</b>				
<i>Chub mackerel</i>	0.14	0.8	0.01	0.9
<i>Cephalopods</i>	-0.15	0.7	0.25	0.6
<i>Pilotfish</i>	-0.09	0.8	-0.08	0.9
<i>Longspine snipefish</i>	-0.21	0.6	0.23	0.6
<b>Wind speed</b>				
<i>Chub mackerel</i>	0.10	0.8	0.14	0.8

<i>Cephalopods</i>	-0.31	0.5	-0.13	0.8
<i>Pilotfish</i>	-0.42	0.3	-0.06	0.9
<i>Longspine snipefish</i>	0.40	0.4	-0.14	0.8
<b>Zonal velocity</b>				
<i>Chub mackerel</i>	-0.39	0.4	-0.18	0.7
<i>Cephalopods</i>	0.22	0.6	-0.17	0.7
<i>Pilotfish</i>	-0.01	0.9	-0.15	0.7
<i>Longspine snipefish</i>	-0.06	0.9	0.28	0.5
<b>Meridional velocity</b>				
<i>Chub mackerel</i>	-0.02	0.9	-0.10	0.8
<i>Cephalopods</i>	0.22	0.6	0.23	0.6
<i>Pilotfish</i>	0.29	0.5	0.12	0.8
<i>Longspine snipefish</i>	-0.38	0.4	0.09	0.9


(a)




(b)




(c)




(d)




(e)




(f)




(g)



(h)



(i)



(j)




Figure A.2- Interannual variation of oceanographic variables in the 5 domains (Selvagem, Pelagic, Seamounts, SPUR: Strong Permanent Upwelling Region and WPUR: Weak Permanent Upwelling Region). Values correspond to the average values ( $\pm$ SD) of the months from each season (Winter: January-April; Summer: June-September). a) Winter and (b) Summer Chlorophyll a, (c) Winter and (d) Summer Sea Surface Temperature, (e) Winter and (f) Summer Wind speed, (g) Winter and (h) Summer Zonal velocity, and (i) Winter and (j) Summer Meridional velocity.

## CHAPTER 6

---



Credits: Aurore Ponchon

# **Food web structure of the subtropical oceanic archipelago of Madeira, eastern North Atlantic**

## **Abstract**

Oceanic environments, as the surroundings of the archipelago of Madeira, are still in need of base line information on their structure and functioning. This is particularly important in the scope of increasing impacts of fisheries and global changes. In this study, we aim to develop a food web model for the area within the marine Exclusive Economic Zone of the archipelago of Madeira, using the Ecopath with Ecosim software. A total of 57 functional groups were defined, representing both coastal and open ocean areas, and epipelagic and deep-sea realms. The Pedigree index defined the model as having reasonable quality. The total biomass of the Madeira system was calculated as  $70 \text{ t} \cdot \text{km}^{-2}$ , with lower trophic levels, like primary producers, zooplankton, other crustaceans and invertebrates, and small pelagic and mesopelagic species, comprising a significant part of its biomass. The food web was characterized by a more linear-like food chain in opposition to a more web-like food chain, with a large proportion of more specialized organisms, like dolphins, shearwaters, and large pelagic fish. Despite the low mean trophic level of the system, the mean trophic level of fisheries was 4.2, targeting mainly top predators, like tunas and Black scabbardfish, which were also the components with most impact in the ecosystem.

**Keywords:** Top predators, Ecopath with Ecosim, Madeira, Food web, Fisheries



## Introduction

The continuous demand for resources has made it increasingly important to understand the ocean as an ecosystem, but such task requires a wealth of information including basic knowledge of its food webs. Food webs are virtual representations of trophic relationships and of the flow of energy and matter through an ecosystem (Cohen et al. 1993). They are useful frameworks to assess the magnitude and importance of trophic relationships with high heuristic value for ecological theories (Link 2002). The study of food webs also allows to depict shifts in the ecosystem, and enables comparisons among similar ecosystems (Cohen et al. 1993). Understanding the structure of food webs and their associated abiotic and biotic factors has far-reaching implications for the conservation of species and their communities, in the sense that the knowledge obtained can be used for management applications (Christensen et al. 2008).

The marine ecosystem has been highly impacted by fisheries in the last 50 to 60 years, when significant examples of technological innovation occurred in larger sized industrial fishing fleets (Valdemarsen 2001), and which continues to deplete stocks at an alarming rate (Worm 2016), despite the warnings and the creation of strategic policies which encourage the implementation of an ecosystem-based approach to fisheries management that would take into account the environmental impacts of fishing, like the European Union Marine Strategy Framework Directive (MSFD; Directive 2008/56/EC, European Parliament 2008). Under sudden environmental and anthropogenic impacts, some species might adapt, while others will tend to migrate to more suitable areas. However, the remainder will eventually disappear, which can further alter the ecosystem functionality and increase the extent of shifts in the community structure and in food webs (Crespo & Dunn 2017, Hillebrand et al. 2018).

In the past decades, the impacts and shifts brought about by global changes and human activities on open oceans have been increasingly noticed. Even though open oceans are often oligotrophic and considered species-poor ecosystems in comparison to coastal areas, they still house complex trophic links of which little is known yet. Studies using stable isotope and fatty acid analysis have been used to describe these food webs (e.g. Bode et al. 2007, Denda et al. 2017, Madgett et al. 2019). However, these are limited to the species sampled and lack power with increasing complexity of the web, as they can carry some uncertainties and errors, like species with multiple organic inputs and consumers which cannot be discerned by one or two stable isotope tracers, or because

isotope ratios are limited and can change according to species composition, metabolic pathway, season or geographic region (Michener & Kaufman 2007).

One way to achieve an ecosystemic view is by means of a mass-balanced model. The Ecopath with Ecosim (EwE) software provides for one of such approaches based on the work of Polovina (1984). The model implemented by EwE considers the biomass and energy flows among different functional groups, and can include fisheries as a particular functional group within the ecosystem (Christensen & Walters 2004). The model has been applied to different aquatic ecosystems (Coll  ter et al. 2015), focusing on assessment of ecosystem maturity (Chea et al. 2016), temporal variations (Heymans et al. 2004) or impacts of climate change (Tam et al. 2008), fisheries (Torres et al. 2019) or of invasive species (Arias-Gonz  lez et al. 2011).

The archipelago of Madeira is a group of oceanic islands located in the subtropical East Atlantic region, far from the influence of the productive Canary Current Upwelling System. The area comprising this archipelago and that of the Selvagem islands, located ca. 300 km south of the Madeira archipelago, provides a great variety of habitats like intertidal and shelf areas, which represent a small proportion of all habitats, and deep-ocean areas, which are the dominant habitat and with the highest ecological and economic importance (e.g. Hern  ndez & Ortega 2000, Freitas et al. 2012, Farias et al. 2013, Alonso et al. 2018). Madeira, like many other oceanic ecosystems, is characterized by oligotrophic and, therefore, low productivity waters. Still, several predators use these waters, including tunas, seabirds, cetaceans, turtles, small pelagic fish and other deep-sea fish. The fact that there is already a considerable amount of information about the ecology and distribution of many species and groups offer the opportunity to build a mass-balance model for the region, including information about the local fisheries (e.g. Hern  ndez & Ortega 2000, Granadeiro et al. 2006, Alonso et al. 2014, Alves et al. 2015, Gouveia et al. 2019, Raposo et al. 2019). Providing the importance of the oceanic region of Madeira for these top predators, it is important to establish baseline information on the Madeira food web structure. This will allow to depict changes in their biomasses and relationships, but also to model alternative futures according to different predictions on fisheries or global changes.

The aim of this study is to provide a model of the ecosystem of the Exclusive Economic Zone (EEZ) of Madeira using the EwE software. This model synthesises the most recent information available concerning the trophic relationships in the Madeira region, describing the main energy and matter flows among the most important functional

groups of the ecosystem. We also include recent data concerning the prevalence of fisheries in the region and its current impact in the structure of the ecosystem.

## Methods

### *Study Area*

This study is focused on the Exclusive Economic Zone (EEZ) of Madeira (hereby referred to as Madeira system), which is an ensemble of groups of islands covering 454,495 km<sup>2</sup>, constituted by the archipelago of Madeira, with two main islands (Madeira and Porto Santo) and a group of islands, Desertas, located ca. 20 km to southeast, and the Selvagens islands, at ca. 300 km south of the main group (Figure 1). These volcanic islands are characterized by narrow continental shelves, with the rest of the region being dominated by extensive abyssal plains ca. 4000 m deep. In the north and northeast of Madeira, the abyssal plain is interrupted by the Tore-Madeira Rise, a chain of seamounts ca. 1000 km long and 50 km wide. Seamounts are areas of increased biodiversity (Pitcher et al. 2007) sustained by current-topography interactions, which induces upwelling and enhances primary production (Genin 2004). The Azores Front is the main mesoscale oceanographic feature in the subtropical northeast Atlantic, which allied to the North Atlantic subtropical gyre and the Canary Current, affects primary production, and the depth and intensity of late-winter ocean-mixing in the region (Fründt & Waniek 2012). For these reasons, the corridor created by the Tore-Madeira Rise, the archipelago of Madeira and the Canary Islands, is considered an important fishing area for Portuguese fishing vessels (Campos et al. 2019).

### *Functional groups and fisheries*

The archipelago of Madeira is acknowledged for its biodiversity and fishing tradition. Seabirds, tunas, and many cetaceans are seasonal visitors that use Madeira during the warmer months (Spring to beginning of Autumn). Pelagic forage fish, mainly composed by mackerels and sardines, are prey to these migratory species but also to resident top predators, such as marlins (Istiophoridae) and sharks. Other resident species in these waters include the deep-sea scabbardfish. The vast deep waters surrounding Madeira island harbour two species, the Black scabbardfish *Aphanopus carbo* and Intermediate scabbardfish *A. intermedius*, which together contributed with the highest proportion of total landings in the archipelago between 2000-2013 (Hermida & Delgado 2016). However, since then, the landings of tunas have dominated the fisheries in Madeira again, as it was before 2000 (Gouveia et al. 2017). The tuna pole-and-line fishery is only

present between March and November (Gouveia & Mejuto 2003), but it is the most important revenue for fishermen in Madeira, mainly targeting Bigeye tuna *Thunnus obesus* and Skipjack tuna *Katsuwonus pelamis* (Hermida & Delgado 2016). Purse seining is the third most important type of fishery in Madeira in terms of catch and economic value, and targets small pelagic fishes, like Blue jack mackerel *Trachurus picturatus*, Atlantic chub mackerel *Scomber colias*, and to a lesser extent European pilchard *Sardina pilchardus*. Small fisheries are also reported in Madeira but have less economic importance and represent very low catches. These include the bottom longline and handline and recreational fisheries, which also target top predators, like large pelagic species and squids (Hermida & Delgado 2016). Overall, the Madeira fleet is dominated by artisanal fishing vessels.




Figure 2 - Exclusive Economic Zone (EEZ) of the archipelago of Madeira (grey line), the study area. Bathymetric lines correspond to 200, 1000 and 2000m.

### Modelling Approach

The food web model presented in this study was constructed using the EwE software version 6.6.3 ([www.ecopath.org](http://www.ecopath.org)). This software is a food web modelling facility that provides a trophic steady mass-balanced snapshot of the ecosystem (Ecopath), but can also create temporal dynamics (Ecosim) resulting from time varying biomass and/or

harvest rates (Christensen et al. 2008). In this study, we will use the Ecopath facility alone.

There are two master equations in the parameterization of EwE. The first describes the production term and ensures a mass balance between groups by exploring the energy transfer in each functional group so the inputs will equal the outputs:

$$B_i(P/B)_i EE_i - \sum_{j=1}^n B_j(Q/B)_i DC_{ji} - Y_i - BA_i - E_i = 0 \quad (1)$$

Where  $B_i$  is the mean biomass of the functional group  $i$  (in  $t \cdot km^{-2}$ );  $P/B_i$  is the production/biomass ratio of the functional group  $i$  ( $year^{-1}$ );  $EE_i$  is the fraction of the functional group  $i$  consumed by a predator or caught in fisheries (i.e., the proportion of the production that is used in the system, designated as Ecotrophic Efficiency);  $Q/B_i$  is the food consumption per biomass unit of the functional group  $i$  ( $year^{-1}$ );  $DC_{ij}$  is the fraction of the functional group  $i$  consumed by predator  $j$ ;  $Y_i$  is the total fishery catch rate of the functional group  $i$  ( $t \cdot km^{-2}$ );  $E_i$  is the net migration rate of the functional group  $i$  ( $year^{-1}$ ); and  $BA_i$  is the biomass accumulation rate for the functional group  $i$  ( $year^{-1}$ ).

The second equation is based on the principle of conservation of matter and describes the energy balance for each functional group:

$$Q = P + R + UAF \quad (2)$$

Where  $Q$  is the consumption of a functional group (in  $t \cdot km^{-2} \cdot year^{-1}$ ),  $P$  is the sum of production of the same group ( $t \cdot km^{-2} \cdot year^{-1}$ ),  $R$  is its respiration, and  $UAF$  is the unassimilated food by that group.

Ecopath can estimate missing parameters and ensure the energy balance within the different functional groups by using the links between the production of each functional group and the consumption of all functional groups (Christensen et al. 2008). The software includes routines which consider the estimation uncertainty (generally designated as pedigree) associated with the model inputs of four parameters: biomass, production/biomass and consumption/biomass ratios, and ecotrophic efficiency of the different functional groups (Christensen & Walters 2004, Plaganyi 2007). At least three of these parameters need to be provided to the model so it can estimate the missing parameter. Additional information on diet composition and fisheries is also needed to complete the model (Christensen & Walters 2004).

### *Model Structure and Parameterization*

The present model represents the food web of the EEZ of the archipelago of Madeira, including the vertical layers and from coastal to pelagic domains of this ecosystem.

The species included in this model comprised the most common and best documented species for the EEZ of Madeira, including those that are targeted by commercial fisheries (Hermida & Delgado 2016). Whenever data on biomass, production, consumption, and diet of relevant species were not available for the region, we resorted to the information available for similar ecosystems, preferably in the east Atlantic region. These species were grouped according to their diet and feeding ecology, and primary producers of the ecosystem, in 56 functional groups, representing seabirds (3 groups), marine mammals (6), sea turtles (1), fish (32), cephalopods (3), other invertebrates (6), algae (1), and plankton (4) (see Supplementary Materials, Table S1). The detritus was also included in this list. Some groups composed of one or several species were specifically defined considering their economic importance in the region. Other groups were split into different size classes, whether because good quality data were available for these groups or because predator dietary data was fine enough to justify prey-size stratification.

Due to their economic and social importance, Black scabbardfish (including both *Aphanopus carbo* and *A. intermedius*), Bigeye tuna, Skipjack tuna *Katsuwonus pelamis*, Atlantic chub mackerel *Scomber colias* and Blue jack mackerel *Trachurus picturatus*, were included as individual functional groups. The Sperm whale *Physeter macrocephalus* and the Monk seal *Monachus monachus* were also separated into single species functional groups due to their particular ecology and behaviour, more specifically due to the use of deep waters and coastal areas, respectively. Multi species groups were arranged according to biological and ecological traits, such as diet (herbivores, carnivores), domain (epipelagic, mesopelagic, demersal) or size (see Supplementary Materials, Table S1).

Biomass data for many groups was not available for our study region and was therefore estimated by the model. In the remaining cases (23%), ecotrophic efficiency was the parameter to be estimated by the model. Whenever possible, parameters P/B, Q/B or P/Q were either based on recent and local literature (e.g. Morissette et al. 2010, Alves et al. 2018) or obtained from models of similar oceanographic and ecological areas (e.g. Morato et al. 2016). We also used data on production and consumption from studies elsewhere in the Atlantic (e.g. Piroddi et al. 2015), whenever possible, and on information on lengths and growth coefficients available in Fishbase (Froese & Pauly 2020) to

calculate natural mortality. A similar approach was applied to diet composition data. For further information on the input parameters used in the model, see Supplementary material, Table S1 and Table S2. The period of 2005-2010 was chosen as reference to build the model as most data was originated from those years.

Fisheries data from Madeira were obtained from Direção Regional de Estatística da Madeira (DREM; [estatistica.madeira.gov.pt](http://estatistica.madeira.gov.pt)), Instituto Nacional de Estatística (INE; [www.ine.pt](http://www.ine.pt)), and local studies (e.g. Gouveia et al. 2019). Currently, there are six gear types in Madeira, purse seining (small pelagic fish), handline and pole-line fishing (large pelagic fish, tuna, demersal fish), drifting longlines (deepwater species, scabbardfish), set or bottom longlines (finfish, demersal fish), fishing pots and traps (finfish, various demersal fish, crustaceans and cephalopods), and game fishing (marlin) (DGPA 2010, Vallerani et al. 2017). Catch values were estimated based on a dataset of 17 years (2000-2017; see Supplementary material, Table S3). Discard values were estimated based on studies carried out in the Madeira region (Martins & Ferreira 1995, Severino 2004) and studies from the Canary Islands (Pajuelo et al. 2010).

Ecopath incorporates a tool to describe the origin and quality of the data input to the model, enabling an informal assessment of the reliability of the model. The Pedigree index is attributed to each input parameter and each functional group, varying between 0 and 1, with lower values representing non-local and highly uncertain data, such as estimations by the model, and higher values representing local and precise data (Christensen & Walters 2004). The final index can vary between 0 and 1, indicating a poor or high-quality final model, respectively.

#### *Model outputs*

The model provides the (average) Trophic Level (TL) of each functional group based on the diet composition of all groups, and the Omnivory index (OI), which is defined by the degree of trophic specialization of each consumer (Christensen et al. 2008). The Connectance index describes whether the food web is web or linear-like by measuring the number of food links in a system in relation to the total number of possible links (Gardner & Ashby 1970). The System Omnivory Index (SOI) is defined by the average Omnivory index of all consumers weighted by the logarithm of their consumption, which can indicate the trophic specialization of the whole system. Diet overlap was estimated according to the similarities in the diet among different functional groups in the model, following Pianka's (1974) equation.

To assess the most important and impacting/impacted functional groups in the Madeira ecosystem, we used the Keystoneness index and the Mixed trophic impact (MTI) routine, respectively. The Keystoneness index identifies the species that have large impacts on other species or functional groups, despite their relative low abundances (Paine 1995, Libralato et al. 2006). The MTI also enables the assessment of the impact of fisheries in each functional group and in the food web.

## Results

### *Model balancing and Parameters of the balanced model*

The Pedigree index for the Madeira System was estimated at 0.39 (Table 1). Final input parameters of each functional group used for the Madeira ecosystem model are described in Table 2. Ecotrophic efficiency values varied greatly according to the proportion of the production of each functional group used by the system. The lowest values corresponded to marine mammals, seabirds and pelagic sharks, while intermediate trophic levels and other functional groups targeted by fisheries presented the highest values.

Table 1 – Summary statistics and ecosystem attributes of the Madeira system

Parameter	Value	Units
<b>Ecosystem properties</b>		
Sum of all consumption	1066.338	t·km <sup>-2</sup> ·y <sup>-1</sup>
Sum of all exports	755.208	t·km <sup>-2</sup> ·y <sup>-1</sup>
Sum of all respiratory flows	570.879	t·km <sup>-2</sup> ·y <sup>-1</sup>
Sum of all flows into detritus	1141.623	t·km <sup>-2</sup> ·y <sup>-1</sup>
Sum of all production	1608.277	t·km <sup>-2</sup> ·y <sup>-1</sup>
Total system throughput (TST)	3534.049	t·km <sup>-2</sup> ·y <sup>-1</sup>
Total net primary production	1326.086	t·km <sup>-2</sup> ·y <sup>-1</sup>
Net system production	755.207	t·km <sup>-2</sup> ·y <sup>-1</sup>
Total biomass (excluding detritus)	70.747	t·km <sup>-2</sup> ·y <sup>-1</sup>
<b>Ecosystem maturity</b>		
Total primary production/total respiration	2.323	
Total primary production/total biomass	18.744	
Total biomass/total throughput	0.020	
<b>Food web structure</b>		
Mean Transfer Efficiency	15.102	%
Connectance Index	0.135	
System Omnivory Index	0.232	
<b>Model reliability</b>		
Pedigree index	0.39	



### *Summary statistics*

The summary statistics and flow indices are presented in Table 1. The model estimated a total biomass (excluding detritus) of  $70.75 \text{ t}\cdot\text{km}^{-2}$ , of which 22.8% were primary producers and 26% were zooplanktonic species. Invertebrates were by far the group with the highest biomass (36.8%), of which molluscs and other benthos constituted the major proportion. Cephalopods, crabs and shrimps are also accounted in this group. Fish followed with 14.2% of all biomass. Air-breathing fauna, such as seabirds, marine mammals and sea turtles only represented 0.15% of the biomass found in this ecosystem.

### *Trophic levels*

The food web structure, interactions and flow diagram of the Madeira ecosystem is represented in Figure 2. The Madeira ecosystem model was composed by five trophic levels (TL), with an average TL (biomass-weight average) of  $2.19 \pm 0.79$  (Table 2). The maximum trophic level was 5.03 (Sperm whale). The deep-sea sharks and Killer whale followed with  $\text{TL}=4.69$ , pelagic sharks with  $\text{TL}=4.57$  and the Black scabbardfish with  $\text{TL}=4.56$ . Other functional groups with a  $\text{TL} > 4$  included species of large pelagic fish, marine mammals, seabirds, tunas, and cephalopods. Small pelagic and mesopelagic fish had TL between 2.86 and 3.36, and TL between 2.96 and 3.08, respectively. Baleen whales, coastal birds, small petrels and sea turtles had TLs between 3.72 and 3.98.

### *Predation and Omnivory index*

Excluding detritus and primary producers, zooplanktonic species were the most consumed functional groups in the Madeira system, both in number of predators (29) and average proportion in the diet of all predators of the ecosystem (19%; see Supplementary materials, Table S4). Myctophids and small mesopelagic followed with the next higher number of predators (26 and 16). Small pelagic fishes (including Chub mackerel and Blue jack mackerel) had an average proportion in the diet of predators similar to mesopelagic fish (10% and 9%, respectively). Small/medium mesopelagic cephalopods and shrimps were also among the functional groups with the highest number of predators (21 and 20, respectively) and with a higher average proportion in the diet of predators (5% and 3%, respectively; see Supplementary materials, Table S4).

Among top predators, dolphins, shearwaters, tunas, and large pelagic fish had the most similar diet when analysing the resolution used in this model. Toothed whales and Sperm whales also fed on similar prey.




Figure 2 – Ecopath flow diagram representing the food web structure and trophic levels of the Madeira system. The different sizes of the circles represent the logarithmic biomass of each functional group. The colour of each link corresponds to the proportion of the functional group on the diet of its consumer; light blue corresponds to proportions close to 0 and red to proportions close to 100%.

Overall, omnivory index values were low, with the maximum value reaching 0.55 for medium reef-associated fish (Table 2). Chub mackerel followed with the next highest value. The index among top predators varied between 0.07 for turtles and 0.39 for Monk seal (Table 2).

#### *Mixed Trophic Impact and Keystoneness index*

Detritus, macroalgae, phytoplankton, small-sized zooplankton and small pelagic fish had the highest averaged positive impacts on other species (see Supplementary materials, Figure S1). On the other hand, baleen whales, dolphins, and medium bathypelagic fish (>1000m) were the functional groups with the highest averaged negative impacts. The model also identified top predators, like dolphins, large demersal fish, Killer whales, and pelagic sharks as keystone groups in the Madeira ecosystem (see Supplementary materials, Table S3).

#### *Fisheries*

The average total catches in Madeira system between 2000-2017 was 0.019 t·km<sup>-2</sup>, targeting a mean trophic level of 4.26, and with a gross efficiency of the fishery at 0.000015.

Tunas and other large pelagic fish presented the highest mortality rate by fisheries (F/Z >90%), followed by Black scabbardfish (67%), and medium pelagic fish (46%; Table 2).

Table 2 - Input and output parameters of the Madeira Ecosystem model. Estimated values by Ecopath are in bold. P: Production (t·km<sup>-2</sup>·yr<sup>-1</sup>); B: Biomass; Q: Consumption (t·km<sup>-2</sup>·yr<sup>-1</sup>); EE: Ecotrophic efficiency; F: Fishing mortality; Z or P/B: Total mortality; OI: Omnivory Index. \* fished species

Group name	Trophic level	Habitat (%)	Biomass (t·km <sup>-2</sup> )	P/B (yr <sup>-1</sup> )	Q/B (yr <sup>-1</sup> )	EE	P/Q (yr <sup>-1</sup> )	F/Z	OI
Baleen whales	3.72	100	0.0409	0.01	6.26	<b>0.00</b>	<b>0.0016</b>		<b>0.36</b>
Toothed whales	4.48	100	0.0039	0.04	9.92	<b>0.27</b>	<b>0.0040</b>		<b>0.14</b>
Sperm whale	5.03	100	0.0555	0.05	5.03	<b>0.00</b>	<b>0.0099</b>		<b>0.17</b>
Killer whales	4.69	100	0.0001	0.02	7.76	<b>0.00</b>	<b>0.0026</b>		<b>0.20</b>
Dolphins	4.29	100	0.0086	0.05	13.74	<b>0.26</b>	<b>0.0036</b>		<b>0.11</b>
Monk seal	4.22	1	2.0·10 <sup>-7</sup>	0.08	13.15	<b>0.00</b>	<b>0.0061</b>		<b>0.40</b>
Shearwaters	4.37	100	0.0001	0.09	42.72	<b>0.00</b>	<b>0.0021</b>		<b>0.17</b>
Small petrels	3.90	100	3.5·10 <sup>-5</sup>	0.19	91.24	<b>0.42</b>	<b>0.0021</b>		<b>0.29</b>
Coastal birds	3.76	15	4.1·10 <sup>-6</sup>	0.05	69.73	<b>0.00</b>	<b>0.0007</b>		<b>0.10</b>
Turtles	3.98	100	0.0005	0.15	3.50	<b>0.93</b>	<b>0.0429</b>	<b>0.1530</b>	<b>0.07</b>
Pelagic sharks	4.57	100	<b>0.0005</b>	0.27	1.44	0.40	<b>0.1875</b>	<b>0.4000</b>	<b>0.30</b>
Deep-sea sharks*	4.69	95	<b>0.0387</b>	0.36	2.01	0.60	<b>0.1791</b>	<b>0.0417</b>	<b>0.30</b>

Rays	3.93	90	<b>0.0004</b>	0.22	2.29	0.60	<b>0.0961</b>		<b>0.22</b>
Pelagic L*	4.53	100	<b>0.0002</b>	0.73	5.27	0.95	<b>0.1385</b>	<b>0.9500</b>	<b>0.15</b>
Bigeye tuna*	4.26	100	<b>0.0016</b>	1.88	6.00	0.95	<b>0.3133</b>	<b>0.9208</b>	<b>0.18</b>
Skipjack tuna*	4.10	100	<b>0.0027</b>	1.43	10.40	0.95	<b>0.1375</b>	<b>0.9163</b>	<b>0.27</b>
Other tunas*	4.27	100	<b>0.0003</b>	0.96	8.07	0.95	<b>0.1190</b>	<b>0.9500</b>	<b>0.23</b>
Pelagic M*	4.22	100	<b>0.0001</b>	0.87	3.93	0.95	<b>0.2214</b>	<b>0.4613</b>	<b>0.05</b>
Epipelagic S/M	3.16	100	<b>0.0124</b>	0.83	5.35	0.95	<b>0.1551</b>		<b>0.53</b>
Pelagic S*	2.86	100	<b>0.1102</b>	1.98	16.03	0.95	<b>0.1235</b>	<b>0.0002</b>	<b>0.38</b>
Chub mackerel*	3.34	100	<b>0.0293</b>	1.98	5.60	0.95	<b>0.3536</b>	<b>0.0163</b>	<b>0.14</b>
Blue Jack mackerel*	3.36	100	<b>0.0317</b>	1.76	6.50	0.95	<b>0.2708</b>	<b>0.0257</b>	<b>0.27</b>
Reef-associated L	3.95	15	<b>4.0·10<sup>-7</sup></b>	0.44	3.42	0.95	<b>0.1287</b>		<b>0.12</b>
Reef-associated M*	2.78	15	<b>0.0008</b>	1.26	12.14	0.95	<b>0.1038</b>	<b>0.0040</b>	<b>0.55</b>
Reef-associated S	2.87	15	<b>0.0081</b>	2.49	17.53	0.95	<b>0.1420</b>		<b>0.47</b>
Demersal L*	4.19	30	<b>0.0056</b>	0.57	2.88	0.95	<b>0.1979</b>	<b>0.0243</b>	<b>0.28</b>
Demersal M*	3.98	30	<b>0.0050</b>	0.34	3.67	0.95	<b>0.0926</b>	<b>0.0026</b>	<b>0.06</b>
Demersal S*	3.73	30	<b>0.0101</b>	1.23	6.30	0.95	<b>0.1952</b>	<b>0.0013</b>	<b>0.28</b>
Benthic S/M	3.54	30	<b>0.0078</b>	1.08	5.87	0.95	<b>0.1840</b>		<b>0.17</b>
Mesopelagic S	3.08	100	<b>1.1647</b>	2.27	14.34	0.95	<b>0.1583</b>		<b>0.15</b>
Myctophids	2.96	100	<b>2.9788</b>	1.76	13.88	0.95	<b>0.1268</b>		<b>0.23</b>
Benthopelagic L*	4.32	100	<b>0.0020</b>	0.57	2.04	0.95	<b>0.2794</b>	<b>0.0558</b>	<b>0.26</b>
Benthopelagic M1*	3.53	50	<b>0.0112</b>	1.01	4.29	0.95	<b>0.2354</b>	<b>0.0053</b>	<b>0.38</b>
Benthopelagic M2*	4.01	100	<b>0.1949</b>	0.30	3.28	0.95	<b>0.0915</b>	<b>0.0003</b>	<b>0.04</b>
Benthopelagic S	3.11	100	<b>0.1381</b>	1.20	5.52	0.95	<b>0.2174</b>		<b>0.14</b>
Bathypelagic M	4.14	100	<b>1.8069</b>	0.56	4.46	0.95	<b>0.1256</b>	<b>7.1·10<sup>-5</sup></b>	<b>0.13</b>
Bathydemersal L*	4.17	30	<b>0.0164</b>	0.42	2.98	0.95	<b>0.1409</b>	<b>0.0036</b>	<b>0.35</b>
Bathydemersal M1	3.54	30	<b>0.1697</b>	0.30	2.23	0.95	<b>0.1345</b>	<b>2.7·10<sup>-5</sup></b>	<b>0.20</b>
Bathydemersal M2	3.48	90	<b>1.5556</b>	0.35	2.42	0.95	<b>0.1446</b>	<b>0.0001</b>	<b>0.12</b>
Bathydemersal S	3.17	90	<b>1.6837</b>	0.70	5.91	0.95	<b>0.1184</b>		<b>0.13</b>
Black scabbard fish*	4.56	30	<b>0.0259</b>	0.53	2.52	0.95	<b>0.2103</b>	<b>0.6667</b>	<b>0.29</b>
Epipelagic cephalopods S*	4.07	100	<b>0.0147</b>	8.50	25.00	0.95	<b>0.3400</b>	<b>1.7·10<sup>-5</sup></b>	<b>0.21</b>
Mesopelagic cephalopods L	4.33	100	<b>0.0824</b>	2.50	10.00	0.95	<b>0.2500</b>		<b>0.29</b>
Mesopelagic cephalopods S/M	3.47	100	<b>0.3267</b>	4.45	16.86	0.95	<b>0.2639</b>		<b>0.14</b>
Jellyfish	3.08	100	<b>0.0265</b>	13.87	50.48	0.95	<b>0.2748</b>		<b>0.44</b>
Crabs	2.59	100	<b>0.1792</b>	1.60	10.00	0.95	<b>0.1600</b>		<b>0.37</b>
Shrimps	2.86	100	<b>3.6210</b>	1.45	11.74	0.95	<b>0.1235</b>		<b>0.51</b>
Echinoderms	2.70	100	<b>1.5917</b>	1.72	6.91	0.85	<b>0.2489</b>		<b>0.05</b>
Molluscs	2.39	100	<b>8.7528</b>	2.24	5.50	0.90	<b>0.4073</b>		<b>0.30</b>
Other benthos	2.63	100	<b>8.6220</b>	2.74	12.25	0.90	<b>0.2237</b>		<b>0.42</b>
Sessile filter feeders	2.20	100	<b>2.8427</b>	0.80	9.00	0.50	<b>0.0889</b>		<b>0.02</b>
Suprabenthic zooplankton	2.05	100	<b>3.7398</b>	16.00	42.00	0.95	<b>0.3810</b>		<b>0.16</b>
Zooplankton L	2.28	100	<b>3.7805</b>	8.70	29.00	0.95	<b>0.3000</b>		<b>0.24</b>
Zooplankton S	2.00	100	10.9000	11.21	43.29	<b>0.69</b>	<b>0.2590</b>		<b>0</b>
Macroalgae	1.00	10	<b>6.8237</b>	27.95		0.50			<b>0</b>
Phytoplankton	1.00	100	9.3200	121.82		<b>0.31</b>			<b>0</b>
Detritus	1.00	100	1000			<b>0.34</b>			<b>0</b>

## Discussion

This study represents a snapshot of the ecosystem state of Madeira system in 2005-2010 and constitutes the first mass-balance model for this region. The lower trophic levels (primary producers, zooplankton, crabs and shrimps, benthic invertebrates and sessile filter-feeders, and small pelagic and mesopelagic fish) represented the highest proportion of biomass in this ecosystem, and also the most predated groups. The structure of the food web is characterized by a more linear-like food chain in opposition to a web-like food chain, with a high proportion of specialists, like dolphins and other epipelagic feeders. Some of these top predators were also considered keystone species, i.e. with high impact on the structure and functioning of the ecosystem. Fisheries showed to have significant impacts on top predators, like tunas and Black scabbardfish.

### *Balancing of the model*

The large variety of functional groups defined aimed at including the highest number of species from which we could find appropriate information, belonging to all vertical layers and distances to the coast. Like in many other regions (e.g. Couce-Montero et al. 2015, Morato et al. 2016, Veiga-Malta et al. 2019), despite the existence of local and good quality data for most groups, many input parameters still had to be taken from studies from nearby regions, other models or, ultimately, estimated by the model. The Pedigree index estimated a value of 0.39 which is lower than the average calculated by Colléter et al. (2015), 0.472 (n=34 models), but in the upper range of the intermediate pedigree (0.200-0.399), as considered by Morissette (2007), who assessed the quality of a set of 50 models. Even so, we are aware of the necessity of refining the information provided for some of the parameters of the model, in order to improve the quality of the model.

### *Ecosystem structure*

The total biomass estimated by the model,  $70.75 \text{ t} \cdot \text{km}^{-2}$ , was similar to the Gulf of Cadiz ( $80.02 \text{ t} \cdot \text{km}^{-2}$ ; Torres et al. 2013), but higher than for the Azores EEZ system ( $24.73 \text{ t} \cdot \text{km}^{-2}$ ; Morato et al. 2016) and lower than that for the Portuguese shelf ( $116 \text{ t} \cdot \text{km}^{-2}$ ; Veiga-Malta et al. 2019). The higher biomass values are most probably due to the influence of upwelling processes in the Portuguese coastal shelf, which enhance biological productivity and provide conditions to support higher abundances of organisms. The Madeira system is dominated by lower trophic levels in terms of total biomass, like benthonic invertebrates and sessile filter feeders, primary producers, and

zooplankton. Fish only corresponded to 14% of all biomass, cephalopods corresponded to 0.5%, and air-breeding species only to 0.15%. This type of composition was also observed in other oceanic ecosystems in the same region, the Gran Canaria and the Azores EEZ systems (Couce-Montero et al. 2015, Morato et al. 2016), but differed from coastal systems, where fish represented the highest proportion of biomass (Torres et al. 2019, Veiga-Malta et al. 2019, Paradell et al. 2020).

The structure of a food web can be characterized by a more linear or more web-like food chains, depending on the number of links of each species, i.e. depending on the overall diet specialization. This can be assessed by two indexes estimated by the model, the System Omnivory Index (SOI) and the Connectance Index (CI). The Madeira system showed a more linear food web pattern (SOI=0.23 and CI=0.13), with relatively more specialists than generalists (the Omnivory Index varied between 0.01-0.54), which agrees with the low complexity of the food web. The Atlantic chub mackerel, an opportunistic species with a varied diet of planktonic and fish species in Madeira (Romero et al. submitted - Chapter 2), had among the highest omnivory indexes in the ecosystem, 0.53. On the other hand, dolphins, turtles, shearwaters, tunas, and large pelagic fish presented a low omnivory index ( $<0.2$ ), showing a highly specialized diet composed of few small epipelagic species. These indexes presented comparable values to other nearby systems, in which similar taxonomic resolutions were used, both oceanic (Morato et al. 2016) and coastal (Heymans et al. 2004, Essekhyr et al. 2019, Torres et al. 2019, Veiga-Malta et al. 2019), and also to the average values estimated for the east Atlantic (Heymans et al. 2014).

Zooplanktonic species, myctophids, other small mesopelagic and epipelagic fish as well as cephalopods are the most predated species in this ecosystem. The MTI analysis has further indicated them as having high positive impacts on the other functional groups, since they are the main prey of many predators. Yet, some top predators (dolphins, Killer whales, pelagic sharks, and large demersal fish) also had important impacts on the biomass of several species and in the ecosystem, overall, as the MTI and Keystoneness index suggested. Despite their overall low biomass, these groups were composed by animals with high nutrition needs and, therefore, with high use of marine resources. Thus, top predators have a strong influence on ecosystem dynamics and on the abundance of other species, maintaining the community organization and biodiversity (Mills et al. 1993, Paine 1995). Dolphins, shearwaters, tunas, and other large pelagic fish had among the higher prey overlap between functional groups. These are epipelagic consumers in the

oceanic waters of Madeira, feeding mainly of mackerels and other small pelagic fish, on cephalopods, and few small mesopelagic fish (Veiga et al. 2011, Alonso et al. 2014, Dromby 2018, Romero et al. submitted - Chapter 3 and 5). Even the Bigeye tuna, which is majorly mesopelagic elsewhere (Ohshimo et al. 2018, da Silva et al. 2019), is mostly an epipelagic feeder in Madeira (Romero et al. Submitted - Chapter 3). Toothed and Sperm whales occupy a different niche, feeding mostly on mesopelagic species, like cephalopod and myctophids (Clarke et al. 1993, Dromby 2018). Such information is valuable because it gives an overview of the most impacting and impacted species and on possible shifts in the food web in the face of continued fishing or global changes.

Overall, the TLs of several functional groups for the Madeira system were similar to the ones defined in other models in nearby regions, like the Gran Canaria system (Couce-Montero et al. 2015), the Azores EEZ (Morato et al. 2016), and the Portuguese shelf systems (Torres et al. 2019, Veiga-Malta et al. 2019). Unfortunately, the diet of many deep-sea species from the northeast Atlantic is not well studied, which led to differences in the estimation of the TL. Further studies on the diet of these groups are needed to obtain a better model of the food web structure in the ecosystem of Madeira.

### *Fisheries*

In Madeira, fisheries target mostly top predators, like tunas and scabbardfish (Hermida & Delgado 2016), which are common in oceanic regions like the Azores (Morato et al. 2016). This is why the average trophic level of fisheries in Madeira is so high, even higher than in the Azores (3.95; Morato et al. 2016). In Madeira, fisheries of intermediate forage fish are less developed than in coastal areas due to their low profitability in terms of cost/efficiency of the fisheries and cost at the market (Hermida & Delgado 2016). The total fisheries catch for the Madeira system is expected to be low ( $0.019 \text{ t}\cdot\text{km}^{-2}$ ) due to the low ratio of fishing areas in relation to the study area. This estimation is similar to the catches observed in the Azores EEZ where a similar study area is analysed ( $0.02 \text{ t}\cdot\text{km}^{-2}$ ; Morato et al. 2016), but lower than the values observed in coastal systems (e.g. Torres et al. 2019, Veiga-Malta et al. 2019, Paradell et al. 2020), which present high productivity and biomasses, but also smaller study areas. Consequently, the gross efficiency of fisheries in Madeira system also had lower estimations than systems with higher catches (e.g. Couce-Montero et al. 2015, Veiga-Malta et al. 2019).

Fisheries were the most important factor of mortality of tunas and other large pelagic fish, and of Black scabbardfish in the oceanic region of the archipelago of

Madeira. However, our model suggested that the ecosystem should remain in equilibrium if fisheries are to continue exploiting the marine resources of Madeira at the same rate. While this might be true in the case of the Black scabbardfish, as its spawning grounds are in the region of Madeira (Farias et al. 2013), for tunas, which are migratory species, the health of the population will mostly depend on its status in the spawning and nursing areas. Nevertheless, while the Skipjack tuna population status remains unknown, the Atlantic Bigeye tuna stock has been considered overexploited by ICCAT (2018), which may raise concerns on the sustainability of this fisheries. A steep reduction in the stock of any of these three groups of species will have significant impacts on the structure of the food web, most probably impacting the epipelagic and deep-sea ecosystems, and consequently also impacting the economy of the archipelago of Madeira.

## **Conclusion**

As done in models from other ecosystems (e.g. Vasconcellos & Watson 2004), we attempted to build a comprehensive representation of the Madeira system, putting together the highest possible number of representative groups. However, for some there was less regional information available, such as for deep-sea species and lower trophic levels (plankton, crustaceans). These gaps were filled by data from other similar regions, which probably resulted in a relatively low Pedigree Index and, thus, lower suitability and quality of the model. To correct this, it will be important to estimate the biomass of the most important species in the Madeira system but also study the diet and reproduction of these groups, to raise the confidence of the model.

This study is a first effort to build a model of the food web structure of Madeira and a step forward towards the understanding of this ecosystem. It has also given us valuable insights on the most important trophic levels, on the interactions among species, and a brief look at some of the impacts that fisheries may have over the marine ecosystem of Madeira. A more refined version of this model will not only allow to create a well-sustained ecosystem-based fishery management but will also allow to predict possible outcomes of different fisheries scenarios, and even of global changes, providing the opportunity to define the most adequate strategies towards a more sustainable marine resource exploitation.



## References

- Alonso H, Granadeiro JP, Dias MP, Catry T, Catry P (2018) Fine-scale tracking and diet information of a marine predator reveals the origin and contrasting spatial distribution of prey. *Prog Oceanogr* 162:1–12
- Alonso H, Granadeiro JP, Waap S, Xavier J, Symondson WOC, Ramos JA, Catry P (2014) An holistic ecological analysis of the diet of Cory's shearwaters using prey morphological characters and DNA barcoding. *Mol Ecol* 23:3719–3733
- Alves F, Dinis A, Nicolau C, Ribeiro C, Kaufmann M, Fortuna C, Freitas L (2015) Survival and abundance of short-finned pilot whales in the archipelago of Madeira, NE Atlantic. *Mar Mammal Sci* 31:106–121
- Alves F, Ferreira R, Fernandes M, Halicka Z, Dias L, Dinis A (2018) Analysis of occurrence patterns and biological factors of cetaceans based on long-term and fine-scale data from platforms of opportunity: Madeira Island as a case study. *Mar Ecol* 39:1–13
- Arias-González JE, González-Gándara C, Luis Cabrera J, Christensen V (2011) Predicted impact of the invasive lionfish *Pterois volitans* on the food web of a Caribbean coral reef. *Environ Res* 111:917–925
- Bode A, Alvarez-Ossorio MT, Cunha ME, Garrido S, Peleteiro JB, Porteiro C, Valdés L, Varela M (2007) Stable nitrogen isotope studies of the pelagic food web on the Atlantic shelf of the Iberian Peninsula. *Prog Oceanogr* 74:115–131
- Campos A, Lopes P, Fonseca P, Figueiredo I, Henriques V, Gouveia N, Delgado J, Gouveia L, Amorim A, Araujo G, Drago T, Santos A dos (2019) Portuguese fisheries in seamounts of Madeira-Tore (NE Atlantic). *Mar Policy* 99:50–57
- Chea R, Guo C, Grenouillet G, Lek S (2016) Toward an ecological understanding of a flood-pulse system lake in a tropical ecosystem: Food web structure and ecosystem health. *Ecol Modell* 323:1–11
- Christensen V, Walters CJ (2004) Ecopath with Ecosim: Methods, capabilities and limitations. *Ecol Modell* 172:109–139
- Christensen V, Walters C, Pauly D, Forrest R (2008) Ecopath with Ecosim version 6: User Guide.
- Clarke MR, Martins HR, Pascoe P (1993) The diet of sperm whales off the Azores. *Philos Trans R Soc B Biol Sci* 339:67–82

- Cohen AJE, Beaver R a, Cousins SH, Deangelis DL, Goldwasser L, Heong KL, Holt RD, Kohn AJ, Lawton JH, Martinez N, O'Malley R, Page LM, Patten BC, Pimm SL, Polis GA, Rejmánek M, Schoener TW, Schoenly K, Sprules WG, Teal JM, Ulanowicz RE, Warren PH, Wilbur HM, Yodzis P (1993) Improving Food Webs Stable. *Ecology* 74:252–258
- Colléter M, Valls A, Guitton J, Gascuel D, Pauly D, Christensen V (2015) Global overview of the applications of the Ecopath with Ecosim modeling approach using the EcoBase models repository. *Ecol Modell* 302:42–53
- Couce-Montero L, Christensen V, Castro JJ (2015) Effects of small-scale and recreational fisheries on the Gran Canaria ecosystem. *Ecol Modell* 312:61–76
- Crespo GO, Dunn DC (2017) A review of the impacts of fisheries on open-ocean ecosystems. *ICES J Mar Sci* 74:2283–2297
- Denda A, Stefanowitsch B, Christiansen B (2017) From the epipelagic zone to the abyss: Trophic structure at two seamounts in the subtropical and tropical Eastern Atlantic - Part I zooplankton and micronekton. *Deep Res Part I Oceanogr Res Pap* 130:63–77
- DGPA (2010) Data Collection for the Fisheries Sector. pp 92
- Dromby M (2018) Assessment of the ecophysiological conditions and trophic ecology of two cetacean species (*Tursiops truncatus* and *Globicephala macrorhynchus*) in Madeira Archipelago, using RNA/DNA ratios and Carbon/Nitrogen stable isotopes
- Essekhyr H, Khalil K, Damsiri Z, Derhy G, Elkalay K (2019) Trophic interactions in the coastal ecosystem of Morocco: An Ecopath approach. *Community Ecol* 20:161–171
- Farias I, Morales-Nin B, Lorange P, Figueiredo I (2013) Black scabbardfish, *Aphanopus carbo*, in the northeast Atlantic: Distribution and hypothetical migratory cycle. *Aquat Living Resour* 26:333–342
- Freitas L, Dinis a., Nicolau C, Ribeiro C, Alves F (2012) New records of cetacean species for Madeira archipelago with an updated checklist. *Bol do Mus Munic do Funchal (História Nat* 62:25–43
- Froese R, Pauly D (2020) Fish Base. World Wide Web Electronic Publication. <https://www.fishbase.in/home.htm>
- Fründt B, Waniek J (2012) Impact of the azores front propagation on deep ocean particle flux. *Open Geosci* 4:531–544

- Gardner MR, Ashby WR (1970) Connectance of large dynamic (Cybernetic) systems: Critical values for stability. *Nature* 228:784
- Genin A (2004) Bio-physical coupling in the formation of zooplankton and fish aggregations over abrupt topographies. *J Mar Syst* 50:3–20
- Gouveia L, Amorim A, Alves A, Hermida M (2017) Updated Fishery Statistics for Bigeye, Skipjack and Albacore Tunas From Madeira Archipelago. *Collect Vol Sci Pap ICCAT* 73:1547–1560
- Gouveia L, Amorim A, Alves A, Hermida M (2019) Updated fishery statistics of tuna species caught off Madeira archipelago. *Collect Vol Sci Pap ICCAT* 75:2081–2097
- Gouveia L, Mejuto J (2003) Seasonality and interannual variability in catches of skipjack tuna (*Katsuwonus pelamis*) and bigeye tuna (*Thunnus obesus*) in the area around the Archipelago of Madeira. *Col Vol Sci Pap ICCAT* 55:1853–1867
- Granadeiro JP, Dias MP, Rebelo R, Santos CD, Catry P (2006) Numbers and Population Trends of Cory's Shearwater *Calonectris diomedea* at Selvagem Grande, Northeast Atlantic. *Waterbirds* 29:56–60
- Hermida M, Delgado J (2016) High trophic level and low diversity: Would Madeira benefit from fishing down? *Mar Policy* 73:130–137
- Hernández JJC, Ortega ATS (2000) Synopsis of Biological Data on The Chub Marckerel (*Scomber japonicus* Houttuyn, 1782). Rome, FAO
- Heymans JJ, Coll M, Libralato S, Morissette L, Christensen V (2014) Global patterns in ecological indicators of marine food webs: A modelling approach. *PLoS One* 9
- Heymans JJ, Shannon LJ, Jarre A (2004) Changes in the northern Benguela ecosystem over three decades: 1970s, 1980s, and 1990s. *Ecol Modell* 172:175–195
- Hillebrand H, Brey T, Gutt J, Hagen W, Metfies K, Meyer B, Lewandowska A (2018) Climate Change: Warming Impacts on Marine Biodiversity. *Handb Mar Environ Prot*:353–373
- ICCAT (2018) Report of the 2018 ICCAT Bigeye tuna stock assessment meeting.
- Libralato S, Christensen V, Pauly D (2006) A method for identifying keystone species in food web models. *Ecol Modell* 195:153–171
- Link J (2002) Does food web theory work for marine ecosystems? *Mar Ecol Prog Ser* 230:1–9
- Madgett AS, Yates K, Webster L, McKenzie C, Moffat CF (2019) Understanding marine food web dynamics using fatty acid signatures and stable isotope ratios: Improving

- contaminant impacts assessments across trophic levels. *Estuar Coast Shelf Sci* 227:106327
- Martins R, Ferreira C (1995) Line Fishing for Black Scabbardfish (*Aphanopus carbo* Lowe, 1839) and Other Deep Water Species in the Eastern Mid Atlantic to the North of Madeira. *Deep Fish North Atl Ocean Slope*:323–335
- Michener RH, Kaufman L (2007) Stable isotope ratios as tracers in marine food webs: An update. In: Blackwell.p 238–282
- Mills LS, Soulé ME, Doak DF (1993) The Keystone-Species Concept in Ecology and Conservation. *Bioscience* 43:219–224
- Morato T, Lemey E, Menezes G, Pham CK, Brito J, Soszynski A, Pitcher TJ, Heymans JJ (2016) Food-web and ecosystem structure of the open-ocean and deep-sea environments of the azores, NE Atlantic. *Front Mar Sci* 3:1–13
- Morissette L (2007) Complexity, cost and quality of ecosystem models and their impact on resilience: a comparative analysis, with emphasis on marine mammals and the gulf of St. Lawrence
- Morissette L, Kaschner K, Gerber LR (2010) Ecosystem models clarify the trophic role of whales off Northwest Africa. *Mar Ecol Prog Ser* 404:289–302
- Ohshimo S, Hiraoka Y, Sato T, Nakatsuka S (2018) Feeding habits of bigeye tuna (*Thunnus obesus*) in the North Pacific from 2011 to 2013. *Mar Freshw Res* 69:585–606
- Paine RT (1995) Diversity on Refining the Concept of Keystone. *Conserv Biol* 9:962–964
- Pajuelo JG, González JA, Santana JI (2010) Bycatch and incidental catch of the black scabbardfish (*Aphanopus* spp.) fishery off the Canary Islands. *Fish Res* 106:448–453
- Paradell O, López BD, Methion S, Rogan E (2020) Food-web interactions in a coastal ecosystem influenced by upwelling and terrestrial runoff off North-West Spain. *Mar Environ Res* 157:1–15
- Pianka ER (1974) E aijNj. *Proc Natl Acad Sci U S A* 71:2141–2145
- Piroddi C, Coll M, Steenbeek J, Moy DM, Christensen V (2015) Modelling the Mediterranean marine ecosystem as a whole: Addressing the challenge of complexity. *Mar Ecol Prog Ser* 533:47–65
- Pitcher TJ, Morato T, Hart PJB, Clark MR, Haggan N, Santos RS (2007) Seamounts: Ecology, Fisheries & Conservation.
- Plaganyi EE (2007) Models for an ecosystem approach to fisheries. Rome, FAO

- Polovina JJ (1984) Model of a Coral Reef Ecosystem. *Coral Reefs* 3:1–11
- Raposo C, Patrício AR, Catry P, Dellinger T, Granadeiro JP (2019) Evidence for trophic differences between live and bycatch oceanic juvenile loggerhead sea turtles. *Mar Biol* 166:1–8
- Severino B (2004) Faculdade de Ciências do Mar e do Ambiente Contributo para o estudo da espécie *Centrophorus squamosus* e sua importância na pescaria de *Aphanopus carbo* na Madeira. Universidade do Algarve
- Silva GB da, Hazin HG, Hazin FHV, Vaske-Jr T (2019) Diet composition of bigeye tuna (*Thunnus obesus*) and yellowfin tuna (*Thunnus albacares*) caught on aggregated schools in the western equatorial Atlantic Ocean. *J Appl Ichthyol* 35:1111–1118
- Tam J, Taylor MH, Blaskovic V, Espinoza P, Ballón RM, Díaz E, Wosnitza-mendo C, Argüelles J, Purca S, Ayón P, Quipuzcoa L, Gutiérrez D, Goya E, Ochoa N, Wolff M (2008) Trophic modeling of the Northern Humboldt Current Ecosystem, Part I: Comparing trophic linkages under La Niña and El Niño conditions. *Prog Oceanogr* 79:352–365
- Torres MÁ, Coll M, Heymans JJ, Christensen V, Sobrino I (2013) Food-web structure of and fishing impacts on the Gulf of Cadiz ecosystem (South-western Spain). *Ecol Modell* 265:26–44
- Torres MA, Fonseca P, Erzini K, Cerveira Borges T, Campos A, Castro M, Santos J, Esmeralda Costa M, Marçalo A, Oliveira N, Vingada J (2019) Modelling the impact of deep-water crustacean trawl fishery in the marine ecosystem off Portuguese Southwestern and South Coasts: I) the trophic web and trophic flows.
- Valdemarsen JW (2001) Technological trends in capture fisheries. *Ocean Coast Manag* 44:635–651
- Vallerani M, Martí C, Ojamaa P (2017) Research for PECH Committee - Fisheries in Madeira.
- Vasconcellos M, Watson R (2004) Mass-balance models of oceanic systems in the Atlantic. In: *West African marine ecosystems: models and fisheries impacts*. Fisheries Centre Research Reports.p 171–214
- Veiga-Malta T, Szalaj D, Angélico MM, Azevedo M, Farias I, Garrido S, Lourenço S, Marçalo A, Marques V, Moreno A, Oliveira PB, Paiva VH, Prista N, Silva C, Sobrinho-Gonçalves L, Vingada J, Silva A (2019) First representation of the trophic structure and functioning of the Portuguese continental shelf ecosystem: Insights into the role of sardine. *Mar Ecol Prog Ser* 617–618:323–340

- Veiga P, Xavier JC, Assis C a., Erzini K (2011) Diet of the blue marlin, *Makaira nigricans*, off the south coast of Portugal. *Mar Biol Res* 7:820–825
- Worm B (2016) Averting a global fisheries disaster. *Proc Natl Acad Sci U S A* 113:4895–4897

## Supplementary Materials

Table S4 - Input data and references used for each functional group for the Madeira EEZ ecosystem model. S – Small, M – Medium, L – Large

<b>1. Baleen whales</b> ( <i>Balaenoptera borealis</i> , <i>Balaenoptera brydei</i> , <i>Balaenoptera physalus</i> )		
<b>Biomass</b>	0.040888 t·km <sup>-2</sup>	Density estimates based on boat surveys (Freitas et al. 2014)
<b>P/B</b>	0.01 year <sup>-1</sup>	Individual body weight (Aguilar 2009, Horwood 2009, Kato & Perrin 2009)
<b>Q/B</b>	6.26 year <sup>-1</sup>	Morissette et al. 2010
<b>EE</b>	Estimated by the model	
<b>Diet</b>	Kawamura 1980, Tershy et al. 1993, Konishi et al. 2009	
<b>2. Toothed whales</b> ( <i>Globicephala macrorhynchus</i> , <i>Grampus griseus</i> , <i>Kogia breviceps</i> , <i>Mesoplodon densirostris</i> , <i>Ziphius cavirostris</i> )		
<b>Biomass</b>	0.003906 t·km <sup>-2</sup>	Density and abundance estimates based on boat surveys (INCOGEO 2010, Freitas et al. 2014, Nicolau et al. 2014, Sambolino et al. 2017)
<b>P/B</b>	0.04 year <sup>-1</sup>	Individual body weight (Trites & Pauly 1998)
<b>Q/B</b>	9.92 year <sup>-1</sup>	Alves et al. 2014, Morissette et al. 2010
<b>EE</b>	Estimated by the model	
<b>Diet</b>	Martins et al. 1985, Hernández-García et al. 1994, Santos et al. 2001, MacLeod et al. 2003, Blanco et al. 2006, Santos et al. 2006, Beatson 2007, Santos et al. 2007, Ozturk et al. 2007, Mintzer et al. 2008, Fernández et al. 2009, West et al. 2009, Bearzi et al. 2011, West et al. 2017	
<b>3. Sperm whale</b> ( <i>Physeter macrocephalus</i> )		
<b>Biomass</b>	0.055496 t·km <sup>-2</sup>	Density and abundance estimates based on boat surveys (INCOGEO 2010, Nicolau et al. 2014, Fais et al. 2016)
<b>P/B</b>	0.05 year <sup>-1</sup>	Individual body weight (Trites & Pauly 1998)
<b>Q/B</b>	5.03 year <sup>-1</sup>	Morissette et al. 2010
<b>EE</b>	Estimated by the model	
<b>Diet</b>	Clarke & MacLeod 1974, Clarke et al. 1993, Fernández et al. 2009	
<b>4. Killer whales</b> ( <i>Orcinus orca</i> , <i>Pseudorca crassidens</i> )		
<b>Biomass</b>	0.000253 t·km <sup>-2</sup>	Abundance estimate based on occasional observations reported by boat surveys and other studies (Freitas et al. 2004b, Alves et al. 2006, Freitas et al. 2012, Nicolau et al. 2014, Alves et al. 2018)
<b>P/B</b>	0.02 year <sup>-1</sup>	Individual body weight (Trites & Pauly 1998)
<b>Q/B</b>	7.76 year <sup>-1</sup>	Morissette et al. 2010
<b>EE</b>	Estimated by the model	

Diet		Sekiguchi et al. 1992, Alonso et al. 1999, Hernández-García et al. 2002
<b>5. Dolphins</b> ( <i>Delphinus delphis</i> , <i>Stenella frontalis</i> , <i>Stenella coeruleoalba</i> , <i>Steno bredanensis</i> , <i>Tursiops truncatus</i> )		
<b>Biomass</b>	0.008612 t·km <sup>-2</sup>	Density and abundance estimates based on boat surveys (INCOGEO 2010, Santos et al. 2012, Freitas et al. 2014, Nicolau et al. 2014, Sambolino et al. 2017)
<b>P/B</b>	0.05 year <sup>-1</sup>	Individual body weight (Trites & Pauly 1998)
<b>Q/B</b>	13.74 year <sup>-1</sup>	Morissette et al. 2010
<b>EE</b>	Estimated by the model	
<b>Diet</b>		Blanco et al. 2001, Clua & Grosvalet 2001, Ringelstein et al. 2006, Pusineri et al. 2007, Dinis et al. 2008, Perrin et al. 2008, Brophy et al. 2009, Fernández et al. 2009, Perrin 2009, Aznar et al. 2017, Giménez et al. 2017, Giménez et al. 2018
<b>6. Monk seal</b> ( <i>Monachus monachus</i> )		
<b>Biomass</b>	0.000023 t·km <sup>-2</sup>	Abundance estimate based on local observations (Pires et al. 2008, 2011, Karamanlidis et al. 2016)
<b>P/B</b>	0.08 year <sup>-1</sup>	Individual body weight (Trites & Pauly 1998)
<b>Q/B</b>	13.15 year <sup>-1</sup>	Piroddi et al. 2015
<b>EE</b>	Estimated by the model	
<b>Diet</b>		Karamanlidis et al. 2011, Pierce et al. 2011, Tonay et al. 2016
<b>7. Shearwaters</b> ( <i>Ardenna gravis</i> , <i>Calonectris diomedea</i> , <i>Puffinus lherminieri</i> , <i>Puffinus puffinus</i> )		
<b>Biomass</b>	0.000023 t·km <sup>-2</sup>	Abundance estimates based on census at breeding colonies or at-sea counts (Martín et al. 1987, Geraldès 2000, Lorenzo et al. 2003, Rodríguez et al. 2003, Oliveira & Menezes 2004, Granadeiro et al. 2006, Lorenzo & Barone 2007, Catry et al. 2010, SPNM 2015a, SPNM 2015b, Birdlife International 2018)
<b>P/B</b>	0.09 year <sup>-1</sup>	Spatial and temporal distribution (Mougin & Jouanin 1997, Carlsson & Holmström 2004, Nunes et al. 2010, Rodríguez et al. 2010, Catry et al. 2011, Reyes-González et al. 2011, 2012, Rodríguez & Arcos 2012, Ramos et al. 2013, Meirinho et al. 2014, Campioni et al. 2015, Ramos et al. 2015a, Paiva et al. 2016, Alonso et al. 2018). Individual body weight (Robertson & James 1988, Granadeiro 1993, Shoji et al. 2015, Ronconi et al. 2018)
<b>Q/B</b>	42.72 year <sup>-1</sup>	Based on annual survival rates from Harris 1966, Perrins et al. 1973, Ramos et al. 2012
<b>EE</b>	Estimated by the model	Calculated with consumption equation from Daunt et al. (2008) and prey energy densities from Glaser et al. (2015) and Spitz et al. (2010)
<b>Diet</b>		Brown et al. 1981, Thompson 1987, Monteiro et al. 1996, Paiva et al. 2010, Alonso et al. 2014, Ramos et al. 2015a, Alonso et al. 2018
<b>8. Small petrels</b> ( <i>Bulweria bulwerii</i> , <i>Hydrobates castro</i> , <i>Pelagodroma marina</i> , <i>Pterodroma deserta</i> , <i>Pterodroma madeira</i> )		
<b>Biomass</b>	0.000023 t·km <sup>-2</sup>	Abundance estimates based on census at breeding colonies or at-sea counts (Hernández et al. 1990, Zino & Biscoito 1994, Campos & Granadeiro 1999, Biscoito & Zino 2002, Barone & Delgado 2002, Rodríguez et al. 2003, Oliveira & Menezes 2004, Ramos & Trujillo 2004, Barone & Trujillo 2007, Luzardo et al. 2008, Catry et al. 2010, Menezes et al. 2010, Lorenzo & Rodríguez 2011, Menezes et al. 2011, Catry et al. 2015, SPNM 2015a)
		Spatial and temporal distribution (Ramírez et al. 2013, Rodríguez et al. 2013, Dias et al. 2015, 2016)



		Individual body weight (Zino & Zino 1986, Robertson & James 1988, Nunes 2000)
<b>P/B</b>	0.19 year <sup>-1</sup>	Based on annual survival rates from Warham 1996, Oro et al. 2005, Ramos et al. 2015b, Ramos et al. 2016
<b>Q/B</b>	91.24 year <sup>-1</sup>	Calculated with consumption equation from Daunt et al. 2008
<b>EE</b>	Estimated by the model	
<b>Diet</b>		Spear et al. 2007, Waap 2015, Waap et al. 2017
<hr/>		
<b>9. Coastal birds</b> ( <i>Larus michahellis</i> , <i>Sterna hirundo</i> )		
<b>Biomass</b>	0.000023 t·km <sup>-2</sup>	Abundance estimates based on census at breeding colonies or at-sea counts (Oliveira & Menezes 2004 Catry et al. 2010, Matias & Catry 2010, SPNM 2015b)
		Individual body weight (Matias & Catry 2010)
<b>P/B</b>	0.05 year <sup>-1</sup>	Based on annual survival rates from Morais et al. 1998
<b>Q/B</b>	69.73 year <sup>-1</sup>	Calculated with consumption equation from Daunt et al. 2008
<b>EE</b>	Estimated by the model	
<b>Diet</b>		Monteiro et al. 1996, Granadeiro et al. 2002, Catry et al. 2010, Matias & Catry 2010, Pedro et al. 2013, López et al. 2016, Romero et al. 2019
<hr/>		
<b>10. Turtles</b> ( <i>Caretta caretta</i> )		
<b>Biomass</b>	0.000516 t·km <sup>-2</sup>	Density estimates based on radiotracking and transects in the closeby Canary archipelago (OAG 2013)
		Individual body weight (Delgado et al. 2010)
<b>P/B</b>	0.15 year <sup>-1</sup>	Taken from Guénette & Morato 2001
<b>Q/B</b>	3.50 year <sup>-1</sup>	Taken from Guénette & Morato 2001
<b>EE</b>	Estimated by the model	
<b>Diet</b>		Brongersma 1968, Brongersma 1972, Van Nierop & Den Hartog 1984, Frick et al. 2009
<hr/>		
<b>11. Pelagic sharks</b> ( <i>Alopias superciliosus</i> , <i>Carcharhinus galapagensis</i> , <i>Isurus oxyrinchus</i> , <i>Lamna nasus</i> , <i>Prionace glauca</i> , <i>Sphyrna zygaena</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	0.27 year <sup>-1</sup>	Taken from Morato et al. 2016
<b>Q/B</b>	1.44 year <sup>-1</sup>	Average value calculated for the group. K and L <sub>∞</sub> values for each species taken from Froese & Pauly et al. 2018
<b>EE</b>	0.40	
<b>Diet</b>		Clarke 1996, Preti et al. 2008, 2012, Mendonça 2009, Porsmoguer et al. 2013, 2014, 2015, 2017, Klarian et al. 2018
<hr/>		
<b>12. Deep sea sharks</b> ( <i>Centrophorus granulosus</i> , <i>Centrophorus lusitanicus</i> , <i>Centrophorus squamosus</i> , <i>Centrophorus uyato/machiquensis</i> , <i>Centroscymnus coelolepis</i> , <i>Centroscymnus crepidater</i> , <i>Centroscymnus owstonii</i> , <i>Chimaera opalescens</i> , <i>Dalatias licha</i> , <i>Deania calcea</i> , <i>Deania hystricosa</i> , <i>Deania profundorum</i> , <i>Etmopterus princeps</i> , <i>Etmopterus pusillus</i> , <i>Etmopterus spinax</i> , <i>Galeus melastomus</i> , <i>Harriotta haeckeli</i> , <i>Harriotta realeighana</i> , <i>Heptanchias perlo</i> , <i>Hexanchus griseus</i> , <i>Hydrolagus affinis</i> , <i>Mustelus mustelus</i> , <i>Pseudotriakis microdon</i> , <i>Scymnodon ringens</i> , <i>Somniosus rostratus</i> , <i>Squaliolus laticaudus</i> , <i>Zameus squamulosus</i> )		
<b>Biomass</b>	Estimated by the model	

<b>P/B</b>	0.51 year <sup>-1</sup>	Taken from Morato et al. 2016
<b>Q/B</b>	2.01 year <sup>-1</sup>	Average value calculated for the group. K and L <sub>∞</sub> values for each species taken from Froese & Pauly et al. 2018
<b>EE</b>	0.60	
<b>Diet</b>		Mauchline & Gordon 1983a, Cortés et al. 1999, Neiva et al. 2006, Dunn et al. 2010, Pethybridge et al. 2011, Xavier et al. 2012, Navarro et al. 2014, Barría et al. 2015, 2018
<b>13. Rays and skates</b> ( <i>Bathytoshia lata</i> , <i>Dasyatis pastinaca</i> , <i>Gymnura altavela</i> , <i>Mobula birostris</i> , <i>Myliobatis aquila</i> , <i>Pteroplatytrygon violacea</i> , <i>Raja brachyura</i> , <i>Raja maderensis</i> , <i>Rostroraja alba</i> , <i>Taeniura grabata</i> , <i>Torpedo marmorata</i> , <i>Torpedo nobiliana</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	0.22 year <sup>-1</sup>	Taken from Morato et al. 2016
<b>Q/B</b>	2.29 year <sup>-1</sup>	Average value calculated for the group. K and L <sub>∞</sub> values for each species taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	
<b>Diet</b>		Morato et al. 2003, Saglam & Bascinar 2008, Follesa et al. 2010, Kadri et al. 2014
<b>14. Bigeye tuna</b> ( <i>Thunnus obesus</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	1.88 year <sup>-1</sup>	Taken from Couce-Montero et al. 2015
<b>Q/B</b>	6.00 year <sup>-1</sup>	Value calculated for the species. K and L <sub>∞</sub> values taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	
<b>Diet</b>		Matthews et al. 1977, Kim et al. 1997, Moteki et al. 2001, Bertrand et al. 2002, Young et al. 2010, Vaske-Júnior et al. 2012, Horn et al. 2013, Ménard et al. 2013, Romero (unpublished data)
<b>15. Skipjack tuna</b> ( <i>Katsuwonus pelamis</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	1.43 year <sup>-1</sup>	Taken from Couce-Montero et al. 2015
<b>Q/B</b>	10.40 year <sup>-1</sup>	Value calculated for the species. K and L <sub>∞</sub> values taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	
<b>Diet</b>		Thomas 1962, Nakamura 1965, Batts 1972, Ankenbrandt 1985, Roger 1994, Ramos 1995, Ménard et al. 2003, Fofandi et al. 2012, Mendizabal 2013, Alatorre-Ramirez et al. 2017, Romero (unpublished data)
<b>16. Other tunas</b> ( <i>Auxis rochei</i> , <i>Thunnus alalunga</i> , <i>Thunnus albacares</i> , <i>Thunnus thynnus</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	0.96 year <sup>-1</sup>	Taken from Couce-Montero et al. 2015
<b>Q/B</b>	8.07 year <sup>-1</sup>	Average value calculated for the group. K and L <sub>∞</sub> values for each species taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption

<b>Diet</b>	Matthews et al. 1977, Allain 2004, Potier et al. 2004, 2007, Satoh et al. 2004, Pusineri et al. 2005, Mostarda et al. 2007, Consoli et al. 2008, Karakulak et al. 2009, Young et al. 2010, Goñi et al. 2011, Battaglia et al. 2013, Horn et al. 2013, Olson et al. 2014	
-------------	---	--

**17. Large pelagic fish** (*Acanthocybium solandri*, *Coryphaena equiselis*, *Coryphaena hippurus*, *Kajidia albida*, *Makaira nigricans*, *Tetrapterus georgii*, *Tetrapterus pfluegeri*, *Xiphias gladius*)

<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	0.73 year <sup>-1</sup>	Taken from Morato et al. 2016
<b>Q/B</b>	5.27 year <sup>-1</sup>	Average value calculated for the group. W <sub>∞</sub> taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>	Brock 1984, Guerra et al. 1993, Hernández-García V 1995, Massutí et al. 1998, Moteki et al. 2001, Olson & Magaña 2002, Allain 2003, Satoh et al. 2004, Vaske-Junior et al. 2004, Shimose et al. 2006, Pinheiro et al. 2010, Runderhausen et al. 2010, Veiga et al. 2011, Silva et al. 2014, Tripp-Valdez et al. 2015, Brewton et al. 2016, Rosa-Luís et al. 2017	

**18. Medium pelagic fish** (*Caranx crysos*, *Caranx rhonchus*, *Decapterus macarellus*, *Decapterus punctatus*, *Pomatomus saltatrix*, *Pseudocaranx dentex*, *Sarda sarda*, *Seriola dumerili*, *Seriola fasciata*, *Seriola rivoliana*, *Sphyrna viridensis*, *Trachinotus ovatus*)

<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	0.87 year <sup>-1</sup>	Taken from Morato et al. 2016
<b>Q/B</b>	3.93 year <sup>-1</sup>	Average value calculated for the group. W <sub>∞</sub> taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>	Barreiros et al. 2002, 2003, Campo et al. 2006	

**19. Chub mackerel** (*Scomber colias*)

<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	1.98 year <sup>-1</sup>	Taken from Couce-Montero et al. 2015
<b>Q/B</b>	5.60 year <sup>-1</sup>	Value calculated for the species. W <sub>∞</sub> taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>	Habashi & Wojciechowski 1973, Weiss 1974, Castro 1993, Castro & Santana 1994, Castro 1995, Whabi et al. 2015, Gushchin & Corten 2017, Vieira (unpublished data)	

**20. Blue jack mackerel** (*Trachurus picturatus*)

<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	1.76 year <sup>-1</sup>	Taken from Vasconcelos et al. 2018
<b>Q/B</b>	6.50 year <sup>-1</sup>	Value calculated for the species. W <sub>∞</sub> taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>	Olaso et al. 1999, Cabral & Murta 2002, Gushchin & Corten 2017, Hirsch & Christiansen 2010, Vieira (unpublished data)	

---

**21. Small pelagic fish** (*Atherina presbyter*, *Boops boops*, *Macrorhamphosus scolopax* (juveniles), *Sardina pilchardus*, *Sardinella maderensis*, *Scomberesox saurus*, *Trachurus trachurus*)

<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	1.98 year <sup>-1</sup>	Taken from Morato et al. 2016
<b>Q/B</b>	16.03 year <sup>-1</sup>	Average value calculated for the group. K and L <sub>∞</sub> values for each species taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>	Moreno & Castro 1995, Fock et al. 2002, Matthiessen et al. 2003, Christiansen et al. 2009, Hirsch & Christhiansen 2010, Guschin & Corten 2017	

---

**22. Epipelagic fish** (*Balistes* sp. (juveniles), *Cheilopogon exsiliens*, *Cheilopogon pinnatibarbus*, *Coryphaena* sp. (juveniles), *Exocoetus volitans*, *Naucratus ductor*, *Seriola* sp. (juveniles), *Lagocephalus lagocephalus*, *Ranzania laevis*, *Sphoeroides pachygaster*)

<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	0.83 year <sup>-1</sup>	Taken from Morato et al. 2016
<b>Q/B</b>	5.35 year <sup>-1</sup>	Average value calculated for the group. K and L <sub>∞</sub> values for each species taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>	Deudero 2001, Deudero & Morales-Nin 2001, Cassaza 2008, Febyanti & Syahailatua 2008	

---

**23. Large reef-associated fish** (*Enchelycore anatina*, *Epinephelus marginatus*, *Gymnothorax unicolor*, *Muraena augusti*, *Mycteroperca fusca*)

<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	0.44 year <sup>-1</sup>	Taken from Morato et al. 2016
<b>Q/B</b>	3.42 year <sup>-1</sup>	Average value calculated for the group. K and L <sub>∞</sub> values for each species taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>	Azevedo et al. 1995, Barreiros & Santos 1998, Reñones et al. 2002, Soares et al. 2003, Sallami et al. 2014, Zaidi et al. 2017	

---

**24. Medium reef-associated fish** (*Balistes capriscus*, *Bodianus scrofa*, *Canthidermis sufflamen*, *Chelon labrosus*, *Chelon ramada*, *Diplodus cervinus*, *Diplodus sargus*, *Diplodus vulgaris*, *Kyphosus sectatrix*, *Labrus bergylta*, *Oblada melanura*, *Sarpa salpa*, *Sparisoma cretense*)

<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	1.26 year <sup>-1</sup>	Taken from Morato et al. 2016
<b>Q/B</b>	12.14 year <sup>-1</sup>	Average value calculated for the group. K and L <sub>∞</sub> values for each species taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>	Whitehead et al. 1984, Azevedo 1995, Porteiro et al. 1996, Figueiredo et al. 2005, Espino et al. 2015, Friedlander et al. 2016, 2017	

---

**25. Small reef-associated fish** (*Abudefduf luridus*, *Apogon imberbes*, *Aulostomus strigosus*, *Canthigaster capistrata*, *Chromis limbata*, *Coris julis*, *Coryphoblennius galerita*, *Gobius paganellus*, *Heteropriacanthus cruentatus*, *Lepadogaster candolii*, *Lepadogaster lepadogaster*, *Mauligobius maderensis*,

<i>Ophioblennius atlanticus, Parablennius parvicornis, Sphoeroides marmoratus, Stephanolepis hispidus, Symphodus trutta, Thalassoma pavo, Tripterygion delaisi)</i>		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	2.49 year <sup>-1</sup>	Taken from Morato et al. 2016
<b>Q/B</b>	17.53 year <sup>-1</sup>	Average value calculated for the group. K and L $\infty$ values for each species taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>	Azevedo 1995, Porteiro et al. 1996, Mancera-Rodríguez & Castro-Hernández 2015, Friedlander et al. 2016, 2017	
<b>26. Medium benthopelagic fish</b> ( <i>Dentex dentex, Dentex gibbosus, Pagellus acarne, Pagellus bogaraveo, Pagellus erythrinus, Pagrus pagrus, Spondyllosoma cantharus</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	1.01 year <sup>-1</sup>	Taken from Pajuelo & Lorenzo 1998, Pajuelo & Lorenzo 2000
<b>Q/B</b>	4.29 year <sup>-1</sup>	Average value calculated for the group. K and L $\infty$ values for each species taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>	Gonçalves & Erzini 1998, Labropoulou et al. 1999, Morato et al. 2001, El-Fergani & El-Mor 2014	
<b>27. Benthic fish</b> ( <i>Arnoglossus imperialis, Arnoglossus rueppelii, Apterichtus caecus, Aulopus filamentosus, Bothus podas, Chelidonichthys cuculus, Chelidonichthys lastoviza, Chelidonichthys lucerna, Chelidonichthys obscura, Echiichthys vipera, Lepidorhombus boscii, Lepidotrigla dieuzeidei, Microchirus azevia, Microchirus ocellatus, Mullus surmuletus, Pegusa lascaris, Synchiropus phaeton, Synodus saurus, Synodus synodus, Symphurus insularis, Trachinus draco, Uranoscopus scaber</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	1.08 year <sup>-1</sup>	Taken from Malta et al. 2018
<b>Q/B</b>	5.87 year <sup>-1</sup>	Average value calculated for the group. K and L $\infty$ values for each species taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>	Azevedo 1995, Labropoulou et al. 1997, Teixeira et al. 2009, 2010, Hirsch et al. 2010	
<b>28. Large demersal fish</b> ( <i>Conger conger, Gymnothorax bacalladoi, Gymnothorax maderensis, Gymnothorax polygonius, Phycis blennoides, Phycis phycis, Polyprion americanus</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	0.57 year <sup>-1</sup>	Taken from Morato et al. 2016
<b>Q/B</b>	2.88 year <sup>-1</sup>	Average value calculated for the group. K and L $\infty$ values for each species from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>	Morato et al. 1999, Young & Winn 2003	
<b>29. Medium demersal fish</b> ( <i>Beryx decadactylus, Beryx splendens, Gephyroberyx darwini, Helicolenus dactylopterus, Pontinus kuhlii, Scorpaena scrofa, Setarches guentherii</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	0.34 year <sup>-1</sup>	Taken from Couce-Montero et al. 2015, Morato et al. 2016

<b>Q/B</b>	3.67 year <sup>-1</sup>	Average value calculated for the group. K and L <sub>∞</sub> values for each species taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>		Morato et al. 1998, Dürr & González 2002, Cabiddu et al. 2010
<b>30. Small demersal fish</b> ( <i>Gaidropsarus granti</i> , <i>Scorpaena canariensis</i> , <i>Scorpaena maderensis</i> , <i>Scorpaena notata</i> , <i>Serranus atricauda</i> , <i>Serranus cabrilla</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	1.23 year <sup>-1</sup>	Taken from Couce-Montero et al. 2015
<b>Q/B</b>	6.30 year <sup>-1</sup>	Average value calculated for the group. K and L <sub>∞</sub> values for each species taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>		Morato et al. 2001, Morte et al. 2001
<b>31. Black-scabbard fish</b> ( <i>Aphanopus carbo</i> , <i>Aphanopus intermedius</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	0.53 year <sup>-1</sup>	Taken from Guénette & Morato 2001
<b>Q/B</b>	2.52 year <sup>-1</sup>	Average value calculated for the group. W <sub>∞</sub> taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>		Maul 1961, Clarke & Lu 1995, Freitas 1998, Farias et al. 2014
<b>32. Large benthopelagic fish</b> ( <i>Alepisaurus ferox</i> , <i>Benthodesmus simonyi</i> , <i>Lepidocybium flavobrunneum</i> , <i>Lepidopus caudatus</i> , <i>Promethichthys prometheus</i> , <i>Ruvettus pretiosus</i> , <i>Serrivomer beanii</i> , <i>Serrivomer lanceolatooides</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	0.57 year <sup>-1</sup>	Average value calculated for the group based on species-specific values from Guénette & Morato 2001, Couce-Montero 2015, Morato et al. 2016
<b>Q/B</b>	2.04 year <sup>-1</sup>	Average value calculated for the group. W <sub>∞</sub> taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>		Matthews 1977, Clarke & Lu 1995, Lorenzo & Pajuelo 1995, Morato et al. 1998, Moteki 2001, Klimpel et al. 2006, Vasilakopoulos et al. 2011, Viana et al. 2012, Choy et al. 2013
<b>33. Medium benthopelagic fish</b> ( <i>Arctozenus risso</i> , <i>Brama brama</i> , <i>Diplospinus multistriatus</i> , <i>Gadella imberbes</i> , <i>Gadella maraldi</i> , <i>Hoplostethus atlanticus</i> , <i>Laemonema robustum</i> , <i>Lepidion guentheri</i> , <i>Lestidiops jayakari</i> , <i>Macroparalepis nigra</i> , <i>Magnisudis atlantica</i> , <i>Mora moro</i> , <i>Physiculus dalwigki</i> , <i>Schedophilus medusophagus</i> , <i>Schedophilus ovalis</i> , <i>Scopelosaurus argenteus</i> , <i>Sudis hialina</i> , <i>Taractichthys longipinnis</i> , <i>Zenopsis conchifera</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	0.30 year <sup>-1</sup>	Taken from Guénette & Morato 2001
<b>Q/B</b>	3.28 year <sup>-1</sup>	Average value calculated for the group. W <sub>∞</sub> taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>		Mauchline & Gordon 1980, Rosecchi et al. 1988, Bulman & Koslow 1992, Pierrot-Bults 1998, Garibaldi et al. 2010, Yang 2011, Battaglia et al. 2014, Anastasopoulou et al. 2018

<b>34. Small benthopelagic fish</b> ( <i>Anthias antias</i> , <i>Antigonia capros</i> , <i>Callanthias ruber</i> , <i>Capros aper</i> , <i>Centracanthus cirrus</i> , <i>Chlorophthalmus agassizi</i> , <i>Diretmoides pauciradiatus</i> , <i>Hoplostethus mediterraneus</i> , <i>Macrorhamphosus scolopax</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	1.20 year <sup>-1</sup>	Taken from Guénette & Morato 2001
<b>Q/B</b>	5.52 year <sup>-1</sup>	Average value calculated for the group. W <sub>∞</sub> taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>	Fock et al. 2002, Pais 2002, Matthiessen et al. 2003, Christiansen et al. 2009, Hirsch & Christhansen 2010	
<b>35. Myctophids</b> ( <i>Benthoosema suborbital</i> , <i>Bolinichthys indicus</i> , <i>Ceratoscopelus maderensis</i> , <i>Ceratoscopelus warmingii</i> , <i>Diaphus adenomus</i> , <i>Diaphus dumerilii</i> , <i>Diaphus holti</i> , <i>Diaphus metopoclampus</i> , <i>Diaphus mollis</i> , <i>Diaphus perspicillatus</i> , <i>Diaphus rafinesquii</i> , <i>Diaphus termophilus</i> , <i>Diogenichthys atlanticus</i> , <i>Electrona risso</i> , <i>Gonichthys cocco</i> , <i>Hygophum benoiti</i> , <i>Hygophum hygomii</i> , <i>Hygophum reinhardtii</i> , <i>Hygophum taaningi</i> , <i>Lampadena chavesi</i> , <i>Lampanyctus alatus</i> , <i>Lampanyctus ater</i> , <i>Lampanyctus festivus</i> , <i>Lampanyctus intricarius</i> , <i>Lampanyctus photonotus</i> , <i>Lampanyctus pusillus</i> , <i>Lepidophanes gaussi</i> , <i>Lobianchia dofleini</i> , <i>Lobianchia gemellarii</i> , <i>Myctophum nitidulum</i> , <i>Myctophum punctatum</i> , <i>Myctophum selenops</i> , <i>Nannobrachiium atrum</i> , <i>Notolychnus valdiviae</i> , <i>Notoscopelus bolini</i> , <i>Notoscopelus resplendens</i> , <i>Symbolophorus veranyi</i> , <i>Taaningichthys bathyphilus</i> , <i>Taaningichthys minimus</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	1.76 year <sup>-1</sup>	Average value calculated for the group. K and L <sub>∞</sub> values for each species from Froese & Pauly et al. 2018
<b>Q/B</b>	13.88 year <sup>-1</sup>	Average value calculated for the group. W <sub>∞</sub> taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>	Clarke 1980, Podrazhanskaya 1993, Bernal et al. 2013	
<b>36. Small mesopelagic fish</b> ( <i>Argyropelecus aculeatus</i> , <i>Argyropelecus hemigymnus</i> , <i>Argyropelecus gigas</i> , <i>Bonapartia pedaliota</i> , <i>Cyclothone braueri</i> , <i>Cyclothone microdon</i> , <i>Cyclothone pallida</i> , <i>Cyclothone pseudopallida</i> , <i>Diplophos taenia</i> , <i>Diretmichthys parini</i> , <i>Diretmus argenteus</i> , <i>Gonostoma atlanticum</i> , <i>Gonostoma denudatum</i> , <i>Ichthyococcus ovatus</i> , <i>Margrethia obtusirostra</i> , <i>Maurololicus muelleri</i> , <i>Neoscopelus macrolepidotus</i> , <i>Neoscopelus microchir</i> , <i>Notoscopelus bolini</i> , <i>Sigmops elongate</i> , <i>Sternoptyx diaphana</i> , <i>Sternoptyx pseudobscura</i> , <i>Valenciennellus tripunctulatus</i> , <i>Vinciguerrria attenuata</i> , <i>Vinciguerrria nimbaria</i> , <i>Vinciguerrria poweriae</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	2.27 year <sup>-1</sup>	Average value calculated for the group. K and L <sub>∞</sub> values for each species from Froese & Pauly et al. 2018
<b>Q/B</b>	14.34 year <sup>-1</sup>	Average value calculated for the group. W <sub>∞</sub> taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption
<b>Diet</b>	Merrett & Roe 1974, Mauchline & Gordon 1983b, Gordon et al. 1985, Hopkins & Baird 1985, Uchikawa et al. 2001a, , Uchikawa et al. 2001b, Carmo et al. 2015	
<b>37. Large bathydemersal fish</b> ( <i>Histiobranchus bathybius</i> , <i>Saccopharynx ampullaceus</i> , <i>Simenchelys parasitica</i> , <i>Synaphobranchus affinis</i> , <i>Synaphobranchus kaupii</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	0.42 year <sup>-1</sup>	Average value calculated for the group. K and L <sub>∞</sub> values for each species from Froese & Pauly et al. 2018
<b>Q/B</b>	2.98 year <sup>-1</sup>	Average value calculated for the group. W <sub>∞</sub> taken from Froese & Pauly et al. 2018
<b>EE</b>	0.95	Assumption

Diet	Merrett & Domansky 1985, Houston & Haedrich 1986, Gordon & Duncan 1987, Saldanha et al. 1995, Gordon & Mauchline 1996, Martin & Christhiansen 1997, Marques 1998	
38. Medium bathydemersal fish1 ( <i>Coryphaenoides rupestris</i> , <i>Epigonus telescopus</i> , <i>Lyconus brachycolus</i> , <i>Malacocephalus laevis</i> , <i>Polymixia nobilis</i> , <i>Trachyrincus scabrus</i> )		
Biomass	Estimated by the model	
P/B	0.30 year <sup>-1</sup>	Taken from Guenette & Morato 2001, Morato et al. 2016
Q/B	2.23 year <sup>-1</sup>	Average value calculated for the group. W <sub>∞</sub> taken from Froese & Pauly et al. 2018
EE	0.95	Assumption
Diet	Mauchline & Gordon 1984a, Macpherson & Ruel 1987, Houston & Haedrich 1986, García-Mederos et al. 2010	
39. Medium bathydemersal fish2 ( <i>Aldrovandia affinis</i> , <i>Alepocephalus bairdii</i> , <i>Alepocephalus rostratus</i> , <i>Antimora rostrata</i> , <i>Bajacalifornia megalops</i> , <i>Bathygadus favosus</i> , <i>Bathygadus melanobranchus</i> , <i>Cataetix laticeps</i> , <i>Coelorinchus labiatus</i> , <i>Conocara salmoneum</i> , <i>Coryphaenoides armatus</i> , <i>Coryphaenoides leptolepis</i> , <i>Coryphaenoides mediterraneus</i> , <i>Coryphaenoides profundicolus</i> , <i>Coryphaenoides rudis</i> , <i>Echinomacrus mollis</i> , <i>Gadomus arcuatus</i> , <i>Halosauropsis macrochir</i> , <i>Halosaurus johnsonianus</i> , <i>Halosaurus ovenii</i> , <i>Hymenocephalus italicus</i> , <i>Hymenogadus gracilis</i> , <i>Lyconus brachycolus</i> , <i>Nezumia aequalis</i> , <i>Nezumia longebarbata</i> , <i>Nezumia sclerorhynchus</i> , <i>Spectrunculus grandis</i> )		
Biomass	Estimated by the model	
P/B	0.35 year <sup>-1</sup>	Taken from Morato et al. 2016
Q/B	2.42 year <sup>-1</sup>	Average value calculated for the group. W <sub>∞</sub> taken from Froese & Pauly et al. 2018
EE	0.95	Assumption
Diet	Sedberry & Musick 1978, Mauchline & Gordon 1983b, Mauchline & Gordon 1984a, Gordon & Duncan 1987, Houston & Haedrich 1986, Carrassón & Matallanas 1998, 2002a, 2002b, Drazen et al. 2008, Sever et al. 2008, Fanelli & Cartes 2010	
40. Medium bathypelagic fish ( <i>Astronesthes gemmifer</i> , <i>Astronesthes micropogon</i> , <i>Avocettina infans</i> , <i>Bathophilus digitatus</i> , <i>Bathophilus vaillanti</i> , <i>Benthalbella infans</i> , <i>Borostomias antarcticus</i> , <i>Chauliodus danae</i> , <i>Chauliodus sloani</i> , <i>Chiasmodon niger</i> , <i>Chiostomias pliopterus</i> , <i>Echiostoma barbatum</i> , <i>Eurypharynx pelecانoides</i> , <i>Eustomias obscurus</i> , <i>Evermannella melanoderma</i> , <i>Idiacanthus fasciola</i> , <i>Leptostomias gladiator</i> , <i>Malacosteus niger</i> , <i>Melanonus zugmayeri</i> , <i>Melanostomias tentaculatus</i> , <i>Photonectes braueri</i> , <i>Photostomias guernei</i> , <i>Rhadinesthes decimus</i> , <i>Stomias boa</i> , <i>Stomias longibarbatus</i> )		
Biomass	Estimated by the model	
P/B	0.56 year <sup>-1</sup>	Average value calculated for the group. K and L <sub>∞</sub> values for each species from Froese & Pauly et al. 2018
Q/B	4.46 year <sup>-1</sup>	Average value calculated for the group. W <sub>∞</sub> taken from Froese & Pauly et al. 2018
EE	0.95	Assumption
Diet	Mauchline & Gordon 1983b, Sutton & Hopkins 1996, Butler et al. 2001, Gaskett et al. 2001, Sutton 2005, Prokofiev & Kukuev 2009, Yang et al. 2011, Battaglia et al. 2018	
41. Small bathypelagic/demersal fish ( <i>Barbantus curvifrons</i> , <i>Bathylagichthys greyae</i> , <i>Bathymicrops regis</i> , <i>Bathypterois longipes</i> , <i>Cetomimus hampeli</i> , <i>Cetostoma regaini</i> , <i>Chaunax pictus</i> , <i>Chaunax suttkusi</i> , <i>Chlorophthalmus agassizi</i> , <i>Dolicholagus longirostris</i> , <i>Epigonus constanciae</i> , <i>Holtbyrnia macrops</i> , <i>Howella atlantica</i> , <i>Laemonema yarrellii</i> , <i>Maulisia maui</i> , <i>Melamphaes typhlops</i> , <i>Melanolagus bericoides</i> , <i>Opisthoproctus soleatus</i> , <i>Platyberyx opalescens</i> , <i>Poromitra capito</i> , <i>Rouleina maderensis</i> , <i>Rinoctes nasutus</i> , <i>Searsia koefoedi</i> , <i>Scopelogadus beanie</i> , <i>Xenodermichthys copei</i> )		
Biomass	Estimated by the model	
P/B	0.70 year <sup>-1</sup>	Average value calculated for the group. K and L <sub>∞</sub> values for each species from Froese & Pauly et al. 2018



Q/B	5.91 year <sup>-1</sup>	Average value calculated for the group. W <sub>∞</sub> taken from Froese & Pauly et al. 2018
EE	0.95	Assumption
Diet		Mauchline & Gordon 1983b, Mauchline & Gordon 1984c, Roe & Badcock 1984, Gartner & Musick 1989, Novotny 2018, Woodstock 2018
<b>42. Small epipelagic cephalopods</b> ( <i>Ommastrephes bartramii</i> , <i>Sthenoteuthis pteropus</i> , <i>Todarodes sagittatus</i> , <i>Onychoteuthis banksii</i> , <i>Cranchia scabra</i> , <i>Argonauta argo</i> , <i>Japetella diaphana</i> , <i>Tremoctopus violaceus</i> )		
Biomass	Estimated by the model	
P/B	8.50 year <sup>-1</sup>	Taken from Pinkerton 2011
Q/B	25.00 year <sup>-1</sup>	Taken from Pinkerton 2011
EE	0.95	Assumption
Diet	Piatkowsky et al. 1998, Watanabe et al. 2008, Parry 2006	
<b>43. Small mesopelagic cephalopods</b> ( <i>Abraliopsis pfefferi</i> , <i>Abraliopsis morisii</i> , <i>Abraliopsis gilchristi</i> , <i>Enoploteuthis anapsis</i> , <i>Histioteuthis arcturi</i> , <i>Histioteuthis meleagroteuthis</i> , <i>Histioteuthis corona</i> , <i>Pyroteuthis margaritifera</i> , <i>Pterygioteuthis giardi</i> , <i>Pterygioteuthis gemmata</i> , <i>Liocranchia reinhardti</i> , <i>Helicocranchia pfefferi</i> , <i>Taonius pavo</i> , <i>Megalocranchia oceánica</i> , <i>Bathothauma lyromma</i> , <i>Leachia atlantica</i> , <i>Idioteuthis hjorti</i> , <i>Mastigoteuthis magna</i> , <i>Mastigoteuthis schmidtii</i> , <i>Heteroteuthis dispar</i> , <i>Brachioteuthis picta</i> , <i>Brachioteuthis riisei</i> , <i>Ctenopteryx sicula</i> , <i>Ctenopteryx canariensis</i> , <i>Onykia robsoni</i> , <i>Thysanoteuthis rhombus</i> , <i>Chiroteuthis veranyii</i> )		
Biomass	Estimated by the model	
P/B	4.45 year <sup>-1</sup>	Taken from Morato & Pitcher et al. 2002
Q/B	16.86 year <sup>-1</sup>	Taken from Morato & Pitcher et al. 2002
EE	0.95	Assumption
Diet	Passarella & Hopkins 1991	
<b>44. Large mesopelagic cephalopods</b> ( <i>Architeuthis</i> sp., <i>Todarodes sagittatus</i> , <i>Lepidoteuthis grimaldii</i> , <i>Taningia danae</i> , <i>Haliphron atlanticus</i> , <i>Moroteuthis robsoni</i> , <i>Cycloteuthis sirventi</i> and <i>Ancistrocheirus lesueurii</i> )		
Biomass	Estimated by the model	
P/B	2.50 year <sup>-1</sup>	Taken from Morato & Pitcher et al. 2002
Q/B	10.00 year <sup>-1</sup>	Taken from Morato & Pitcher et al. 2002
EE	0.95	Assumption
Diet	Bolstad & O'Shea 2004, Watanabe et al. 2008, Parry 2006	
<b>45. Crabs</b> ( <i>Plagusia depressa</i> , <i>Percnon gibbesi</i> , <i>Grapsus grapsus</i> , <i>Calcinus tubularis</i> , <i>Dardanus calidus</i> , <i>Pagurus cuanensis</i> , <i>Palinurus elephas</i> , <i>Galathea squamifera</i> , <i>Scyllarides latus</i> , <i>Scyllarides latus</i> , <i>Maja squinado</i> and <i>Pagurus anachoretus</i> , <i>Chaceon affinis</i> , <i>Paromola cuvieri</i> , <i>Spinolambrus macrochelos</i> )		
Biomass	Estimated by the model	
P/B	1.60 year <sup>-1</sup>	Taken from Morato & Pitcher et al. 2002
Q/B	10.00 year <sup>-1</sup>	Taken from Morato & Pitcher et al. 2002

<b>EE</b>	0.95	Assumption
<b>Diet</b>		Morato et al. 2016
<b>46. Shrimps</b> ( <i>Plesionika edwardsii</i> , <i>P. ensis</i> , <i>P. narval</i> , <i>Heterocarpus ensifer</i> , <i>H. laevigatus</i> , <i>H. grimaldii</i> , <i>Acantheephyra eximia</i> , <i>Benthesicymus bartletti</i> , <i>Lepas anatifera</i> , <i>Lepas anserifera</i> , <i>Lepas pectinata</i> , <i>Alepa pacifica</i> and <i>Dosima fascicularis</i> , <i>Planes minutes</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	1.45 year <sup>-1</sup>	Average value calculated for the group based on species-specific values from Couce-Montero et al. (2015), Morato et al. (2016) and Torres et al. (2013)
<b>Q/B</b>	11.74 year <sup>-1</sup>	Average value calculated for the group based on species-specific values from Couce-Montero et al. (2015), Morato et al. (2016) and Torres et al. (2013)
<b>EE</b>	0.95	Assumption
<b>Diet</b>		Cartes et al. 1993, Labropoulou & Kostikas 1999
<b>47. Echinoderms</b> ( <i>Asteroidea</i> , <i>Ophiuroidea</i> , <i>Crinoidea</i> , <i>Echinoidea</i> and <i>Holothuroidea</i> : inc. <i>Antedon bifida</i> , <i>Arbacia lixula</i> , <i>Astropecten irregulares</i> , <i>Centrostephanus longispinus</i> , <i>Coscinasterias tenuispina</i> , <i>Diadema africana</i> , <i>Echinaster sepositus</i> , <i>Endoxocrinus wyvillethomsoni</i> , <i>Holothuria sanctori</i> , <i>Luidia ciliaris</i> , <i>Ophioderma longicauda</i> , <i>Paracentrotus lividus</i> , <i>Sphaerechinus granularis</i> , <i>Stylocidaris affinis</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	1.72 year <sup>-1</sup>	Couce-Montero et al. 2015
<b>Q/B</b>	6.91 year <sup>-1</sup>	Couce-Montero et al. 2015
<b>EE</b>	0.85	Assumption
<b>Diet</b>		Christensen 1970, Brun 1972, Franz & Worley 1982, Beddingfield & McClintock 1993, de Juan et al. 2007
<b>48. Suprabenthic zooplânkton</b> (inc. <i>copepods</i> , <i>amphipods</i> , <i>isopods</i> , <i>euphasiids</i> and <i>mysids</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	16.00 year <sup>-1</sup>	Average value from Sánchez & Olaso 2004, Torres et al. 2013
<b>Q/B</b>	42.00 year <sup>-1</sup>	Average value from Sánchez & Olaso 2004, Torres et al. 2013
<b>EE</b>	0.95	Assumption
<b>Diet</b>		Based on Sánchez & Olaso 2004, Torres et al. 2013
<b>49. Mollusks</b> ( <i>Trochidae</i> , <i>Littorinidae</i> , <i>Patellidae</i> , <i>Siphonoridae</i> , <i>Haliotidae</i> , <i>Columbellidae</i> , <i>Aplysiidae</i> , <i>Cerithidae</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	0.90 year <sup>-1</sup>	Couce-Montero et al. 2015
<b>Q/B</b>	5.50 year <sup>-1</sup>	Couce-Montero et al. 2015
<b>EE</b>	0.85	Assumption
<b>Diet</b>		Based on Couce-Montero et al. 2015
<b>50. Other benthos</b> ( <i>polychaets</i> , <i>annelids</i> and <i>other benthic invertebrates</i> )		
<b>Biomass</b>	Estimated by the model	

<b>P/B</b>	2.74 year <sup>-1</sup>	Average value from Olaso & Sánchez 2004, Torres et al. 2013, Morato et al. 2016
<b>Q/B</b>	12.25 year <sup>-1</sup>	Average value from Olaso & Sánchez 2004, Torres et al. 2013, Morato et al. 2016
<b>EE</b>	0.95	Assumption
<b>Diet</b>		Based on Guénette & Morato 2001
<b>51. Jellyfish</b> ( <i>jellyfishes and hydrozoans (e.g., Pelagia noctiluca, Physalia physalis)</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	13.87 year <sup>-1</sup>	Malej 1989, Coll et al. 2006, Torres et al. 2013
<b>Q/B</b>	50.48 year <sup>-1</sup>	Malej 1989, Coll et al. 2006, Torres et al. 2013
<b>EE</b>	0.95	Assumption
<b>Diet</b>		Rosa et al. 2013, Tilves et al. 2016
<b>52. Sessile filter feeders</b> ( <i>sponges, hard corals, soft corals (gorgonian corals), cold-water corals (inc. anthipatharian corals), anemons, ascidians and polypoid hydrozoans</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	0.80 year <sup>-1</sup>	Optiz et al. 1993, Morato & Pitcher 2002
<b>Q/B</b>	9.00 year <sup>-1</sup>	Optiz et al. 1993, Morato & Pitcher 2002
<b>EE</b>	0.50	Assumption
<b>Diet</b>		Based on Morato & Pitcher 2002
<b>53. Large-sized zooplânkton</b> ( <i>This group consists of zooplankton &gt; 5mm, namely crustaceans such as decapod's larvae and some euphasiids, but also chaetognaths, ctenophors, pteropods, fish larvae and free-swimming tunicates, such as appendicularians or salps</i> )		
<b>Biomass</b>	Estimated by the model	
<b>P/B</b>	8.70 year <sup>-1</sup>	Morato et al. 2009
<b>Q/B</b>	29.00 year <sup>-1</sup>	Morato et al. 2009
<b>EE</b>	0.95	Assumption
<b>Diet</b>		Based on Morato et al. 2009
<b>54. Small-sized zooplânkton</b> ( <i>This group consists of zooplankton &lt; 5mm, namely crustaceans such as copepods, ostracods, amphipods and mysids, but also Radiolaria, bivalve larvae and small tunicates, chaetognaths and ctenophors</i> )		
<b>Biomass</b>	10.900000 t·km <sup>-2</sup>	
<b>P/B</b>	11.21 year <sup>-1</sup>	Morato et al. 2009
<b>Q/B</b>	43.29 year <sup>-1</sup>	Morato et al. 2009
<b>EE</b>	Estimated by the model	Assumption
<b>Diet</b>		Based on Guénette & Morato 2001
<b>55. Macroalgae</b> ( <i>inc. Asparagopsis armata, Asparagopsis taxiformis, Caulerpa webbiana, Chaetomorpha sp., Codium adhaerens, Corallina elongata, Cottoniella filamentosa, Cryptopleura ramosa, Cystoseira marina, Dasycladus vermicularis, Dictyota fasciola, Dictyota neglecta, Halopteris scoparia, Jania rubens,</i>		

<i>Laminaria ochroleuca, Laurencia viridis, Lobophora variegata, Lophocladia trichoclados, Padina pavonica, Pterocladia capillacea, Pseudotetraspora marina, Sargassum filipendula, Stypopodium zonale, Symploca hydroides, Ulva lactula, Ulva rigida, Valonia utricularis, Zonaria tournefortii)</i>		
Biomass	Estimated by the model	
P/B	27.95 year <sup>-1</sup>	Couce-Montero et al. 2015
Q/B	-	
EE	0.50	Assumption
Diet	-	
<b>56. Phytoplânkton</b> (inc. diatoms, dinoflagellates and coccolithophores)		
Biomass	9.320000 t·km <sup>-2</sup>	
P/B	121.82 year <sup>-1</sup>	Calculated dividing the annual primary production by the annual mean biomass
Q/B	-	
EE	Estimated by the model	Assumption
Diet	-	
Other parameters	Integrated Chl-a concentration (0-150m of depth) of 0.932 g C m <sup>-2</sup> (Mendonça et al. 2012) Primary production of 113.5391 g C m <sup>-2</sup> ·year <sup>-1</sup> (Kleisner & Hoornaert 2015)	
<b>57. Detritus</b>		
Biomass	21.809400 t·km <sup>-2</sup>	
P/B	-	
Q/B	-	
EE	Estimated by the model	
Diet	-	
Other parameters	Average euphotic zone of 118m (Christiansen et al. 2010) Primary production of 113.5391 g C m <sup>-2</sup> ·year <sup>-1</sup> (Kleisner & Hoornaert 2015)	

Table S2 – Diet composition of each functional group of the food web of the oceanic region of the archipelago of Madeira.

Prey \ Predator	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24
1 Baleen whales																								
2 Toothed whales				0.05																				
3 Sperm whale																								
4 Killer whales																								
5 Dolphins				0.05							0.10													
6 Monk seal																								
7 Shearwaters																								
8 Small petrels									0.01															
9 Coastal birds																								
10 Turtles				0.05		0.03					0.03													
11 Pelagic sharks																								
12 Deep-sea sharks												0.10												
13 Rays											0.08													
14 Bigeye tuna				0.10																				
15 Skipjack tuna				0.10													0.05							
16 Other tunas																								
17 Pelagic fish L																								
18 Pelagic fish M																	0.05							
19 Chub mackerel	0.10			0.05	0.15		0.25	0.05		0.05				0.20	0.25	0.15	0.15	0.25						
20 Blue Jack mackerel	0.10			0.05	0.15		0.10	0.05		0.10				0.20	0.15	0.15	0.10	0.25						
21 Pelagic fish S	0.20			0.05	0.10		0.20	0.19		0.05			0.05	0.10	0.15	0.10	0.10	0.10	0.35	0.25		0.25		
22 Epipelagic fish S/M				0.10	0.05		0.05								0.10	0.05	0.15	0.15						
23 Reef fish L						0.05																		
24 Reef fish M						0.10							0.05											
25 Reef fish S						0.05		0.05	0.05				0.05				0.15							
26 Benthopelagic fish M1					0.05	0.10						0.05	0.05				0.05						0.05	
27 Benthic fish S/M					0.03	0.10							0.15				0.05							0.20
28 Demersal fish L					0.03	0.10																		
29 Demersal fish M						0.05																		
30 Demersal fish S					0.05	0.05							0.05											

Table S2 – Diet composition of each functional group of the food web of the oceanic region of the archipelago of Madeira (continued horizontally from page 178)

[illegible]

Table S2 – Diet composition of each functional group of the food web of the oceanic region of the archipelago of Madeira (continued vertically from 178)

Prey \ Predator	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24
31 Black scabbard fish												0.05												
32 Benthopelagic fish L2																								
33 Benthopelagic fish M2											0.05	0.05		0.10			0.10							
34 Benthopelagic fish S2					0.05							0.15	0.20	0.05	0.05	0.05								
35 Myctophids	0.10	0.10			0.15		0.05	0.25	0.10	0.05	0.05	0.10		0.10	0.05	0.05	0.10		0.05	0.05				
36 Mesopelagic fish S	0.05	0.15			0.10		0.05	0.20							0.05	0.05								
37 Bathydemersal fish L																								
38 Bathydemersal fish M1												0.05												
39 Bathydemersal fish M2											0.05	0.05												
40 Bathypelagic fish M		0.05										0.10				0.05								
41 Bathydemersal fish S								0.05																
42 Epipelagic cephalopods S	0.05			0.05	0.05		0.20	0.05			0.05				0.05	0.15	0.20		0.05	0.05				
43 Mesopelagic cephalopods S/M		0.60	0.35	0.35	0.05	0.05	0.10	0.15		0.05	0.15	0.25		0.15		0.05								
44 Mesopelagic cephalopods L		0.10	0.65								0.10	0.05												
45 Crabs						0.18			0.25		0.05		0.20										0.10	0.15
46 Shrimps									0.10	0.10	0.10		0.05	0.05	0.05	0.05								
47 Echinoderms																							0.15	
48 Suprabenthic zooplankton																								
49 Mollusks						0.15			0.10															
50 Other benthos																							0.10	0.05
51 Jellyfish										0.65														
52 Sessile filter feeders																								
53 Zooplankton L	0.10							0.20	0.10	0.10			0.10	0.05	0.05	0.10		0.05	0.15	0.05	0.15			
54 Zooplankton S	0.30							0.10					0.05		0.05			0.38	0.45	0.80	0.50		0.20	
55 Macroalgae																		0.03		0.05	0.05		0.40	
56 Phytoplankton																		0.05		0.05				
57 Detritus																		0.05	0.05	0.05	0.05		0.05	

Table S2 – Diet composition of each functional group of the food web of the oceanic region of the archipelago of Madeira (continued vertically from page 179)

	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
31				0.05				0.05																						
32							0.10	0.10					0.05							0.05										
33				0.20	0.10	0.05	0.20	0.15						0.10				0.10		0.05										
34		0.10			0.15		0.10	0.20	0.15				0.10	0.05		0.45		0.30	0.15	0.25										
35					0.05		0.05	0.15	0.20				0.10			0.20		0.15	0.10	0.15										
36							0.10																							
37				0.05									0.05							0.05										
38							0.10						0.05	0.05		0.05				0.10										
39							0.05	0.05	0.15				0.10			0.10				0.05										
40							0.05						0.05	0.05	0.05	0.10		0.05		0.10										
41								0.05										0.10		0.05										
42		0.05			0.05		0.15	0.05					0.20				0.05	0.10	0.10	0.15										
43							0.10																							
44	0.05		0.20	0.15	0.05	0.40								0.10	0.05															
45			0.05		0.30	0.10			0.05		0.05	0.05	0.15	0.10	0.10	0.10	0.05	0.10	0.05											
46	0.05	0.05	0.05												0.05							0.05								
47		0.05	0.15							0.05	0.05		0.05	0.30	0.25		0.40		0.05		0.10	0.10	0.10	0.05	0.20	0.25				
48		0.05	0.15	0.05	0.05									0.15	0.15					0.15	0.05	0.15			0.05	0.10				
49	0.25	0.10	0.30	0.05		0.05				0.10		0.10		0.05	0.20		0.15			0.10	0.10	0.20				0.10				
50								0.25							0.05															
51																0.05				0.05		0.05								
52		0.10				0.10		0.05	0.05	0.15	0.05	0.10	0.05		0.05		0.10	0.05	0.10			0.15			0.05	0.05	0.30		0.10	
53	0.30	0.15			0.05	0.05			0.15	0.70	0.75	0.70	0.05	0.05	0.05		0.10	0.05	0.45		0.05	0.25			0.05		0.70	0.20	0.15	
54	0.25																				0.05		0.30			0.10	0.05		0.10	0.15
55											0.05	0.05														0.10			0.10	0.70
56	0.10	0.05									0.05						0.10				0.50	0.30	0.20	0.95	0.45	0.45		0.70	0.55	0.15



Table S3 - Fisheries data for the Madeira System (2000-2017). S – Small, M – Medium, L – Large

Group name	Purse seine	Scabbard fishery	Tuna pole and line	Bottom longline and handline	Squids fishery	Recreational	Pelagic longline (mainland)	Total	References
Pelagic sharks	0.000000	0.000000	0.000000	0.000000	0.000000	0.000000	0.000025	0.000025	Martins & Ferreira 1995, Severino 2004, Pajuelo et al. 2010
Deep-sea sharks	0.000000	0.000225	0.000000	0.000000	0.000000	0.000000	0.000000	0.000225	<a href="https://estatistica.madeira.gov.pt/">https://estatistica.madeira.gov.pt/</a>
Bigeye tuna	0.000000	0.000000	0.002697	0.000000	0.000000	0.000013	0.000000	0.002710	Gouveia et al. 2017
Skipjack tuna	0.000000	0.000021	0.003489	0.000000	0.000000	0.000011	0.000000	0.003520	Gouveia et al. 2017
Other tunas	0.000000	0.000000	0.000279	0.000000	0.000000	0.000004	0.000000	0.000284	Gouveia et al. 2017
Pelagic L	0.000000	0.000011	0.000000	0.000000	0.000000	0.000083	0.000020	0.000115	<a href="https://www.ine.pt">https://www.ine.pt</a>
Pelagic M	0.000000	0.000000	0.000000	0.000000	0.000000	0.000041	0.000000	0.000041	<a href="https://www.ine.pt">https://www.ine.pt</a>
Chub mackerel	0.000946	0.000000	0.000000	0.000000	0.000000	0.000000	0.000000	0.000946	<a href="https://estatistica.madeira.gov.pt/">https://estatistica.madeira.gov.pt/</a>
Blue Jack mackerel	0.001437	0.000000	0.000000	0.000000	0.000000	0.000000	0.000000	0.001437	<a href="https://estatistica.madeira.gov.pt/">https://estatistica.madeira.gov.pt/</a>
Pelagic S	0.000051	0.000000	0.000000	0.000000	0.000000	0.000000	0.000000	0.000051	<a href="https://estatistica.madeira.gov.pt/">https://estatistica.madeira.gov.pt/</a>
Reef-associated M	0.000000	0.000000	0.000000	0.000004	0.000000	0.000000	0.000000	0.000004	<a href="https://estatistica.madeira.gov.pt/">https://estatistica.madeira.gov.pt/</a>
Benthopelagic M1	0.000000	0.000000	0.000000	0.000060	0.000000	0.000000	0.000000	0.000060	<a href="https://estatistica.madeira.gov.pt/">https://estatistica.madeira.gov.pt/</a>
Demersal L	0.000000	0.000000	0.000000	0.000078	0.000000	0.000000	0.000000	0.000078	<a href="https://estatistica.madeira.gov.pt/">https://estatistica.madeira.gov.pt/</a>
Demersal M	0.000000	0.000000	0.000000	0.000004	0.000000	0.000000	0.000000	0.000004	<a href="https://estatistica.madeira.gov.pt/">https://estatistica.madeira.gov.pt/</a>
Demersal S	0.000000	0.000000	0.000000	0.000016	0.000000	0.000000	0.000000	0.000016	<a href="https://estatistica.madeira.gov.pt/">https://estatistica.madeira.gov.pt/</a>
Black scabbard fish	0.000000	0.008244	0.000000	0.000000	0.000000	0.000000	0.000000	0.008244	<a href="https://www.ine.pt">https://www.ine.pt</a>
Benthopelagic L	0.000000	0.000023	0.000000	0.000000	0.000000	0.000000	0.000000	0.000023	<a href="https://estatistica.madeira.gov.pt/">https://estatistica.madeira.gov.pt/</a>
Benthopelagic M2	0.000000	0.000007	0.000000	0.000000	0.000000	0.000000	0.000000	0.000007	<a href="https://estatistica.madeira.gov.pt/">https://estatistica.madeira.gov.pt/</a>
Epipelagic cephalopods S	0.000000	0.000000	0.000000	0.000000	0.000002	0.000000	0.000000	0.000002	<a href="https://estatistica.madeira.gov.pt/">https://estatistica.madeira.gov.pt/</a>
Sum	0.002434	0.008531	0.006465	0.000163	0.000002	0.000152	0.000046	0.017793	

Table S4 – Average biomass contribution of each functional group as prey for other species (in %), and number of predators per each functional group in the Madeira system. Keystoneness index according to (Libralato et al. 2006). S – Small, M – Medium, L – Large

Prey	Nr. Predators	Prey (%)	Keystoneness index
Baleen whales	0	0.00	-0.19
Toothed whales	1	0.09	-1.21
Sperm whale	0	0.00	-0.07
Killer whales	0	0.00	0.09
Dolphins	2	0.28	0.18
Monk seal	0	0.00	-0.02
Shearwaters	0	0.00	-1.50
Small petrels	1	0.02	-2.00
Coastal birds	0	0.00	-0.004
Turtles	3	0.19	-1.03
Pelagic sharks	0	0.00	0.03
Deep-sea sharks	1	0.19	-0.61
Rays	1	0.14	-1.09
Bigeye tuna	1	0.19	-0.50
Skipjack tuna	2	0.28	-0.36
Other tunas	0	0.00	-1.45
Pelagic L	0	0.00	-0.26
Pelagic M	1	0.09	-1.01
Chub mackerel	11	3.06	-0.42
Blue Jack mackerel	13	2.78	-0.48
Pelagic S	17	4.80	-0.27
Epipelagic S/M	8	1.39	-0.66
Reef-associated L	1	0.09	-1.32
Reef-associated M	4	0.93	-0.66
Reef-associated S	10	1.76	-0.51
Benthopelagic M1	7	0.74	-0.76
Benthic S/M	8	1.25	-0.55
Demersal L	2	0.23	0.09
Demersal M	2	0.28	-0.07
Demersal S	6	0.65	-0.31
Black scabbard fish	1	0.09	-0.18
Benthopelagic L	2	0.19	-1.15
Benthopelagic M2	8	1.11	-0.30
Benthopelagic S	14	2.78	-0.49
Myctophids	26	6.20	-0.18
Mesopelagic S	16	3.33	-0.37
Bathydemersal L	1	0.19	-1.03
Bathydemersal M1	4	0.37	-0.55
Bathydemersal M2	7	0.83	-0.05

Bathypelagic M	9	1.30	-0.13
Bathydemersal S	8	0.93	-0.29
Epipelagic cephalopods S	14	2.13	-0.50
Mesopelagic cephalopods S/M	21	5.93	-0.27
Mesopelagic cephalopods L	5	1.85	-0.09
Crabs	13	3.56	-0.24
Shrimps	20	3.24	-0.004
Echinoderms	6	0.74	-0.36
Suprabenthic zooplankton	15	3.98	-0.33
Molluscs	13	2.50	-0.48
Other benthos	16	3.70	-0.34
Jellyfish	3	1.76	-0.23
Sessile filter feeders	3	0.28	-0.98
Zooplankton L	29	5.19	-0.48
Zooplankton S	29	14.49	-0.06
Macroalgae/seagrass	11	2.82	-0.25
Phytoplankton	8	2.22	-0.18
Detritus	18	8.89	

---




Figure S1 - The Mixed Trophic Impact (MTI) of the Madeira system. Blue and red squares represent the positive and negative impacts, respectively, between functional groups.

## References

- Aguilar A (2009) Fin whale: *Balaenoptera physalus*. In: Encyclopedia of Marine Mammals, Second Edition, pp 433-437.
- Allain V (2003) Diet of Mahi-mahi, Wahoo and Lancetfish in the Western and Central Pacific. SCTB16 Working Paper, 16th Meeting of the Standing Committee on Tuna and Billfish, Mooloolaba, Queensland, Australia, 9-16 July 2003, Oceanic Fisheries Programme, Secretariat of the Pacific Community, Noumea, New Caledonia.

- Alatorre-Ramirez VG, Galvan-Magaria F, Torres-Rojas YE, Olson RJ (2017) Trophic segregation of mixed schools of yellow fin tuna (*Thunnus albacares*) and skipjack tuna (*Katsuwonus pelamis*) caught in the eastern tropical Pacific Ocean. *Fishery Bulletin* 115: 252-269.
- Allain V (2004) Diet of yellowfin tuna in different areas of the western and central Pacific Ocean. 17th Meeting of the Standing Committee on Tuna and Billfish, SCTB17, Majuro, Marshall Islands, 9-18 August 2004. BIO-1, 1-20
- Alonso MK, Pedraza SN, Schiavini A, Goodall RNP, Crespo EA (1999) Stomach contents of false killer whales (*Pseudorca crassidens*) stranded on the coasts of the Strait of Magellan, Tierra del Fuego. *Marine Mammal Science* 15: 712-724.
- Alonso H, Granadeiro JP, Waap S, Xavier J, Symondson WO, Ramos JA, Catry P (2014) An holistic ecological analysis of the diet of Cory's shearwaters using prey morphological characters and DNA barcoding. *Molecular Ecology* 23: 3719-3733.
- Alonso H, Granadeiro JP, Dias MP, Catry T, Catry P (2018) Fine-scale tracking and diet information of a marine predator reveals the origin and contrasting spatial distribution of prey. *Progress in Oceanography* 162: 1-12.
- Alves F, Dinis A, Nicolau C, Ribeiro C, Kaufmann M, Fortuna C, Freitas L (2014) Survival and abundance of short-finned pilot whales in the archipelago of Madeira, NE Atlantic, *Marine Mammal Science* 31: 106–121.
- Alves F, Ferreira R, Fernandes M, Halicka Z, Dias L, Dinis A (2018) Analysis of occurrence patterns and biological factors of cetaceans based on long-term and fine-scale data from platforms of opportunity: Madeira Island as a case study. *Marine Ecology* e12499.
- Alves F, Freitas L, Dinis A (2006) Occurrence and diving behaviour of false killer whale off Madeira archipelago (NE Atlantic). In: 20th Conference of the European Cetacean Society B (Vol. 2).
- Anastasopoulou A, Mytilineou C, Smith CJ, Papadopoulou KN (2018) Crustacean prey in the diet of fishes from deep waters of the Eastern Ionian Sea. *Journal of the Marine Biological Association of the United Kingdom* 1-9.
- Ankenbrandt L (1985) Food habits of bait-caught skipjack tuna, *Katsuwonus pelamis*, from the southwestern Atlantic Ocean. *Fishery Bulletin* 83: 379-393.
- Azevedo JMN (1995) Food web of the Azorean shallow water ichthyological communities: a guild approach. *Boletim do Museu Municipal do Funchal. Suplemento 4 A*: 29-53.

- Azevedo J, Rodrigues JB, Mendizabel M, Arruda LM (1995) Study of a sample of dusky groupers, *Epinephelus marginatus* (Lowe, 1834), caught in a tide pool at Lajes do Pico, Azores. Boletim do Museu Municipal do Funchal 4: 55-64.
- Aznar FJ, Míguez-Lozano R, Ruiz B, de Castro AB, Raga JA, Blanco C (2017) Long-term changes (1990-2012) in the diet of striped dolphins *Stenella coeruleoalba* from the western Mediterranean. Marine Ecology Progress Series 568: 231-247.
- Babiddu S, Pedoni C, Pesci P, Porcu C (2010) Trophic habits of red scorpionfish *Scorpaena scrofa* (Osteichtheys, Scopaeniformes) in the Central Western Mediterranean. Rapp. Comm. int. Mer. Medit. 39: 464.
- Barone & Delgado (2002) Datos de interés sobre las aves nidificantes de la isla de Porto Santo (archipiélago de Madeira). Rev. Acad. Canar. Cienc. XIV 219-225.
- Barone & Trujillo (2007) Pardela de Bulwer In: Lorenzo JA (ed.) (2007) Atlas de las aves nidificantes en el archipiélago canario (1993-2003). Dirección General de Conservación de la Naturaleza-SEO/BirdLife, Madrid.
- Barreiros JP, Santos RS (1998) Notes on the food habits and predatory behaviour of the dusky grouper, *Epinephelus marginatus* (Lowe, 1834) (Pisces: Serranidae) in the Azores. Arquipelago 16A: 29-35.
- Barreiros JP, Santos RS, Borba AESD (2002) Food habits, schooling and predatory behaviour of the Yellowmouth Barracuda, *Sphyraena viridensis* Cuvier, 1829 (Perciformes: Sphyraenidae) in the Azores. Cybium–International Journal of Ichthyology 26: 83-88.
- Barreiros JP, Morato T, Santos RS, Borba AESD (2003) Interannual changes in the diet of the almaco jack *Seriola rivoliana* (Perciformes: Carangidae) from the Azores. Cybium–International Journal of Ichthyology 27: 37-40.
- Barría C, Coll M, Navarro J (2015) Unravelling the ecological role and trophic relationships of uncommon and threatened elasmobranchs in the western Mediterranean Sea. Marine Ecology Progress Series 539: 225-240.
- Barría C, Navarro J, Coll M (2018) Trophic habits of an abundant shark in the northwestern Mediterranean Sea using an isotopic non-lethal approach. Estuarine, Coastal and Shelf Science 207: 383-390.
- Battaglia P, Andaloro F, Consoli P, Esposito V, Malara D, Musolino S, Pedà C, Romeo T (2013) Feeding habits of the Atlantic bluefin tuna, *Thunnus thynnus* (L.1758), in the Central Mediterranean Sea (Strait of Messina). Helgol. Mar. Res. 67: 97-107.

- Battaglia P, Musolino S, Esposito V, Ammendolia G, Consoli P, Andaloro F, Romeo T (2014) Feeding habits of juvenile fishes belonging to three medusivorous species (Centrolophidae and Nomeidae) from the Strait of Messina (central Mediterranean Sea). *Marine Biology Research* 10: 927-933.
- Battaglia P, Ammendolia G, Esposito V, Romeo T, Andaloro F (2018) Few But Relatively Large Prey: Trophic Ecology of *Chauliodus sloani* (Pisces: Stomiidae) in Deep Waters of the Central Mediterranean Sea. *Journal of Ichthyology* 58: 8-16.
- Batts BS (1972) Food habits of the skipjack tuna, *Katsuwonus pelamis*, in North Carolina waters. *Chesapeake Science* 13: 193-200.
- Bearzi G, Reeves RR, Remonato E, Pierantonio N, Airoidi S (2011) Risso's dolphin *Grampus griseus* in the Mediterranean Sea. *Mammalian Biology* 76: 385-400.
- Beatson E (2007) The diet of pygmy sperm whales, *Kogia breviceps*, stranded in New Zealand: implications for conservation. *Reviews in Fish Biology and Fisheries* 17: 295-303.
- Beddingfield SD, McClintock JB (1993) Feeding behaviour of the sea star *Astropecten articulatus* (Echinodermata: Asteroidea): an evaluation of energy-efficient foraging in a soft-bottom predator. *Marine Biology* 115: 669-676.
- Bernal A, Valls M, Olivar MP, Sweeting CJ, Fernández de Puellas ML, Molí Ferrer B (2013) Diet and trophic levels of myctophids, gonostomatids and hatchetfish in the Western Mediterranean. Symposium on Integrating New Advances in Mediterranean Oceanography and Marine Biology, 26-29 November 2013, Institut de Ciències del Mar (CSIC), Barcelona, Catalunya, Spain.
- BirdLife International (2018) Important Bird Areas factsheet: Desertas. Downloaded from <http://www.birdlife.org> on 17/04/2018.
- Biscoito M, Zino F (2002) Aves do arquipélago da Madeira. Direção Regional do Ambiente.
- Blanco C, Salomón O, Raga JA (2001) Diet of the bottlenose dolphin (*Tursiops truncatus*) in the western Mediterranean Sea. *Journal of the Marine Biological Association of the United Kingdom* 81: 1053-1058.
- Blanco C, Raduán MÁ, Raga JA (2006) Diet of Risso's dolphin (*Grampus griseus*) in the western Mediterranean Sea. *Scientia Marina* 70: 407-411.
- Bolstad KS, O'Shea S (2004) Gut contents of a giant squid *Architeuthis dux* (Cephalopoda: Oegopsida) from New Zealand waters, *New Zealand Journal of Zoology*, 31:1, 15-21.

- Brewton RA, Ajemian MJ, Young PC, Stunz GW (2016) Feeding Ecology of Dolphinfish in the Western Gulf of Mexico. *Transactions of the American Fisheries Society* 145: 839-853.
- Brock RW (1984) A contribution to the trophic biology of the blue marlin (*Makaira nigricans* Lacepede, 1802) in Hawaii. *Pacif Sci* 38: 141-149.
- Brongersma LD (1968) Notes upon some turtles from the Canary Islands and from Madeira. Koninkl. Nederl. Akademie van Wetenschappen Amsterdam, Proceedings C 71(2), 128-136.
- Brongersma LD (1972) European Atlantic turtles. *Zoologische Verhandelingen* no. 121. E.J. Brill. Leiden. 318 pages.
- Brophy JT, Murphy S, Rogan E (2009) The diet and feeding ecology of the short-beaked common dolphin (*Delphinus delphis*) in the northeast Atlantic. *International Whaling Commission Scientific Committee paper SC/61/SM14*, 18.
- Brown RGB, Barker SP, Gaskin DE, Sandeman MR (1981) The foods of great and sooty shearwaters *Puffinus gravis* and *P. griseus* in eastern Canadian waters. *Ibis* 123: 19-30.
- Brun E (1972) Food and feeding habits of *Luidia ciliaris* Echinodermata: Asteroidea. *J. mar. biol. Ass. U.K.* 52: 225-236.
- Bulman CM, Koslow JA (1992) Diet and food consumption of a deep-sea fish, orange roughy *Hoplostethus atlanticus* (Pisces: Trachichthyidae), off southeastern Australia. *Marine ecology progress series*. Oldendorf 82: 115-129.
- Butler M, Bollens SM, Burkhalter B, Madin LP, Horgan E (2001) Mesopelagic fishes of the Arabian Sea: distribution, abundance and diet of *Chauliodus pammelas*, *Chauliodus sloani*, *Stomias affinis*, and *Stomias nebulosus*. *Deep Sea Research Part II: Topical Studies in Oceanography* 48: 1369-1383.
- Cabral HN, Murta AG (2002) The diet of blue whiting, hake, horse mackerel and mackerel off Portugal. *Journal of Applied Ichthyology* 18: 14-23.
- Campioni L, Granadeiro JP, Catry P (2015) Niche segregation between immature and adult seabirds: does progressive maturation play a role? *Behavioral Ecology* 27: 426-433.
- Campo D, Mostarda E, Castriota L, Scarabello MP, Andaloro F (2006) Feeding habits of the Atlantic bonito, *Sarda sarda* (Bloch, 1793) in the southern Tyrrhenian sea. *Fisheries Research* 81: 169-175.



- Campos AR, Granadeiro JP (1999) Breeding biology of the White-faced Storm-petrel on Selvagem Grande Island, north-east Atlantic. *Waterbirds* 22: 199-206.
- Carlsson B & Holmström N (2004) Trip Report Madeira. 1-9 September.
- Carmo V, Sutton T, Menezes G, Falkenhaus T, Bergstad OA (2015) Feeding ecology of the Stomiiformes (Pisces) of the northern Mid-Atlantic Ridge. 1. The Sternoptychidae and Phosichthyidae. *Progress in Oceanography* 130: 172-187.
- Carrassón M, Matallanas J, Casadevall M (1997) Feeding strategies of deep-water morids on the western Mediterranean slope. *Deep Sea Research Part I: Oceanographic Research Papers* 44: 1685-1699.
- Carrassón M, Matallanas J (1998) Feeding habits of *Alepocephalus rostratus* (Pisces: Alepocephalidae) in the western Mediterranean Sea. *Journal of the Marine Biological Association of the United Kingdom* 78: 1295-1306.
- Carrassón M, Matallanas J (2002a) Diets of deep-sea macrourid fishes in the western Mediterranean. *Marine Ecology Progress Series* 234: 215-228.
- Carrassón M, Matallanas J (2002b) Feeding habits of *Cataetx alleni* (Pisces: Bythitidae) in the deep western Mediterranean. *Scientia Marina* 66: 417-421.
- Cartes JE (1993) Diets of deep-water pandalid shrimps on the Western Mediterranean slope. *Marine Ecology Progress Series* 96: 49-61.
- Cassazza TL (2008) Community structure and diets of fishes associated with pelagic Sargassum and open-water habitats off North Carolina (Doctoral dissertation, University of North Carolina Wilmington).
- Castro JJ (1993) Feeding ecology of chub mackerel *Scomber japonicus* in the Canary Islands area. *South African Journal of Marine Science* 13: 323-328.
- Castro JJ, Santana del Pino A (1994) Feeding preferences of *Scomber japonicus* in the Canary Islands area. *Scientia Marina (Barcelona)* (3-4).
- Castro J (1995) Mysids and euphysiids in the diet of *Scomber japonicus* Houttuyn, 1782 off the Canary Islands. *Boletín Instituto Español de Oceanografía* 11: 77-86.
- Catry P, Dias MP, Catry T, Pedro P, Tenreiro P, Menezes D (2015) Bulwer's petrels breeding numbers on the Desertas Islands (Madeira): improved estimates indicate the NE Atlantic population to be much larger than previously thought. *Airo* 23: 10-14.
- Catry P, Dias MP, Phillips RA, Granadeiro JP (2011) Different means to the same end: long-distance migrant seabirds from two colonies differ in behaviour, despite common wintering grounds. *PLoS One* 6: e26079.

- Catry P, Geraldles P, Pio JP, Almeida A (2010) Aves marinhas da Selvagem Pequena e do Ilhéu de Fora: censos e notas, com destaque para a dieta da Gaivota-de-patas-amarelas. *Airo* 20: 29-35.
- Christiansen B, Martin B, Hirsch S (2009) The benthopelagic fish fauna on the summit of Seine Seamount, NE Atlantic: Composition, population structure and diets. *Deep Sea Research Part II: Topical Studies in Oceanography* 56: 2705-2712.
- Christiansen B, Albers A, Brand T, Chivers A, Christiansen H, Christiansen S, Denda A, Diniz T, George K-H, Irion I, Janßen T, Kaufmann M, Kullmann B, Lamont P, Molodtsova T, Montgomery J, Peine F, Schneeorst A, Schuster A, Springer B, Stahl H, Stefanowitsch B, Turner G, Turnewitsch R, Vieira R, Vogel S, Wehrmann H (2010) Biodiversity and Biological Production at a Shallow NE Atlantic Seamount (Ampère Seamount), METEOR-Berichte, Cruise No. M83/2, 64 pp., DFG-Senatskommission für Ozeanographie.
- Christensen AM (1970) Feeding biology of the sea star *Astropecten irregularis* Pennant. *Ophelia* 8: 1-134.
- Choy CA, Portner E, Iwane M, Drazen JC (2013) Diets of five important predatory mesopelagic fishes of the central North Pacific. *Marine Ecology Progress Series* 492: 169-184.
- Clarke TA (1980) Diets of 14 species of vertically migrating mesopelagic fishes in Hawaiian waters. *Fish Bull* 78: 619-640
- Clarke MR, Lu CC (1995) Cephalopoda of Madeiran waters. *Bolletim do museu municipal do Funchal* 4: 181-200.
- Clarke MR, MacLeod N (1974) Cephalopod remains from a sperm whale caught off Vigo, Spain. *Journal of the Marine Biological Association of the United Kingdom* 54: 959-968.
- Clarke MR, Martins HR, Pascoe P (1993) The diet of sperm whales (*Physeter macrocephalus* Linnaeus 1758) off the Azores. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 339: 67-82.
- Clarke MR, Clarke DC, Martins HR, Silva HM (1996) The diet of the blue shark (*Prionace glauca* L.) in Azorean waters. *ARQUIPÉLAGO - Ciências Biológicas e Marinhas - Life and Marine Sciences* 14: 41-56.
- Clua É, Grosvalet F (2001) Mixed-species feeding aggregation of dolphins, large tunas and seabirds in the Azores. *Aquatic Living Resources* 14: 11-18.

- Coll M, Palomera I, Tudela S, Sardà F (2006) Trophic flows, ecosystem structure and fishing impacts in the South Catalan Sea, Northwestern Mediterranean. *Journal of Marine Systems* 59: 63-96.
- Consoli P, Romeo T, Battaglia P, Castriota L, Esposito V, Andaloro F (2008) Feeding habits of the albacore tuna *Thunnus alalunga* (Perciformes, Scombridae) from central Mediterranean Sea. *Marine Biology* 155: 113.
- Cortés E (1999) Standardized diet compositions and trophic levels of sharks. *ICES Journal of Marine Science* 56: 707-717.
- Couce-Montero ML (2015) Diagnosis de la pesquería artesanal en la isla de gran canaria (Doctoral dissertation, Universidad de Las Palmas de Gran Canaria).
- Couce-Montero L, Christensen V, Castro JJ (2015) Effects of small-scale and recreational fisheries on the Gran Canaria ecosystem. *Ecological modelling* 312: 61-76.
- Daunt F, Wanless S, Greentree S P R, Jensen H, Hamer K C, Harris M P (2008) The impact of the sandeel fishery closure on seabird food consumption, distribution, and productivity in the northwestern North Sea. *Can. J. Fish. Aquat. Sci.* 65: 362-381.
- de Juan S, Cartes JE, Demestre M (2007) Effects of commercial trawling activities in the diet of the flat fish *Citharus linguatula* (Osteichthyes: Pleuronectiformes) and the starfish *Astropecten irregularis* (Echinodermata: Asteroidea). *Journal of Experimental Marine Biology and Ecology* 349: 152-169.
- Delgado C, Canário AV, Dellinger T (2010) Sex ratios of loggerhead sea turtles *Caretta caretta* during the juvenile pelagic stage. *Marine Biology* 157: 979-990.
- Delgado de Molina AR (2017) Datos estadísticos de la pesquería de túnidos de las Islas Canarias durante el periodo 1975 a 2015. *Col. Vol. Sci. Pap. ICCAT* 73: 737- 748.
- Dellinger T (2008) Tartarugas marinhas. In: Loureiro A, Ferrand de Almeida N, Carretero MA, Paulo OS (eds) *Atlas dos Anfíbios e Répteis de Portugal*. Instituto da Conservação da Natureza e Biodiversidade, Lisboa, pp. 193-209.
- Dellinger T, Encarnação H (2000) Accidental capture of sea turtles by fishing fleet based at Madeira Island, Portugal. In: Kalb H, Wibbels T (eds) *Proceedings of the nineteenth Annual Symposium on Sea Turtle Conservation and Biology*. NOAA Technical Memorandum NOAA NMFS-SEFSC-443, pp. 218–218.
- den Hartog JC (1980) Notes on the food of sea turtles: *Eretmochelys imbricata* (Linnaeus) and *Dermochelys coriacea* (Linnaeus). *Netherlands Journal of Zoology* 30: 595-610.
- Deudero S (2001) Interspecific trophic relationships among pelagic fish species underneath FADs. *Journal of Fish Biology* 58: 53-67.

- Deudero S, Morales-Nin B (2001) Prey selectivity in planktivorous juvenile fishes associated with floating objects in the western Mediterranean. *Aquaculture Research* 32: 481-490.
- Dias MP, Alho M, Granadeiro JP, Catry P (2015) Wanderer of the deepest seas: migratory behaviour and distribution of the highly pelagic Bulwer's petrel. *Journal of Ornithology* 156: 955-962.
- Dias MP, Romero J, Granadeiro JP, Catry T, Pollet IL, Catry P (2016) Distribution and at-sea activity of a nocturnal seabird, the Bulwer's petrel *Bulweria bulwerii*, during the incubation period. *Deep Sea Research Part I: Oceanographic Research Papers* 113: 49-56.
- Dinis A, Cascão I, Alves F, Freitas L, Querouil S (2008) Comparing the epipelagic diet between short-beaked common dolphin and Atlantic spotted dolphin in two Atlantic oceanic archipelagos. 22nd Conference of the European Cetacean Society. Egmond aan Zee, The Netherlands.
- Drazen JC, Popp BN, Choy CA, Clemente T, Forest LD, Smith Jr KL (2008) Bypassing the abyssal benthic food web: Macrourid diet in the eastern North Pacific inferred from stomach content and stable isotopes analyses. *Limnology and Oceanography* 53: 2644-2654.
- Dunn MR, Szabo A, McVeagh MS, Smith PJ (2010) The diet of deepwater sharks and the benefits of using DNA identification of prey. *Deep Sea Research Part I: Oceanographic Research Papers* 57: 923-930.
- Dürr J, González JA (2002) Feeding habits of *Beryx splendens* and *Beryx decadactylus* (Berycidae) off the Canary Islands. *Fisheries Research* 54: 363-374.
- El-Fergani ES, El-Mor M (2014) Feeding habits of the Common dentex, *Dentex dentex* (Linnaeus, 1758) (Teleostei: Sparidae) from Benghazi coast, eastern Libya. *Int. J. Bioassays* 3(11): 3517-3522.
- Escáñez A, Guerra A, González A, Tobeña M, Landeria JM, Aguilar N (2011) Trophic resources of blainville's beaked whales (*Mesoplodon densirostris*) and cuvier's beaked whales (*Ziphius cavirostris*) in El Hierro, Canary Island. In: Gauffier P & Verborgh P (eds) Abstract book. 25th Conference of the European Cetacean Society. Long term datasets on marine mammals: learning from the past to manage the future. 21st-23rd March 2011, Cádiz, Spain.
- Espino F, González JA, Haroun R, Tuya F (2015) Abundance and biomass of the parrotfish *Sparisoma cretense* in seagrass meadows: temporal and spatial differences

- between seagrass interiors and seagrass adjacent to reefs. *Environmental biology of fishes* 98: 121-133.
- Fais A, Lewis TP, Zitterbart DP, Álvarez O, Tejedor A, Aguilar Soto N (2016) Abundance and Distribution of Sperm Whales in the Canary Islands: Can Sperm Whales in the Archipelago Sustain the Current Level of Ship-Strike Mortalities? *PLoS ONE* 11(3): e0150660
- Fanelli E, Cartes JE (2010) Temporal variations in the feeding habits and trophic levels of three deep-sea demersal fishes from the western Mediterranean Sea, based on stomach contents and stable isotope analyses. *Marine Ecology Progress Series* 402: 213-232.
- Farias I, Figueiredo I, Janeiro AI, Bandarra NM, Batista I, Morales-Nin B (2014) Reproductive and feeding spatial dynamics of the black scabbardfish, *Aphanopus carbo* Lowe, 1839, in NE Atlantic inferred from fatty acid and stable isotope analyses. *Deep Sea Research Part I: Oceanographic Research Papers* 89: 84-93.
- Febyanti F, Syahailatua A (2008) Food habit of flying fishes, *Hirundichthys oxycephalus* and *Cheilopogon cyanopterus* in Makassar Strait, *J Lit. Perikan. Ind.* 14: 115-122.
- Fernández R, Santos MB, Carrillo M, Tejedor M, Pierce GJ (2009) Stomach contents of cetaceans stranded in the Canary Islands 1996–2006. *Journal of the Marine Biological Association of the United Kingdom* 89: 873-883.
- Ferreira T (2001) Abundância relativa de tartaruga-comum *Caretta caretta* (Linnaeus, 1758) na ZEE da Madeira. Relatório de Estágio da Licenciatura em Biologia Aplicada aos Recursos Animais, Faculdade de Ciências da Universidade de Lisboa, Lisboa.
- Fletcher N, Batjakas IE, Pierce GJ (2013) Diet of the Atlantic bonito *Sarda sarda* (Bloch, 1793) in the Northeast Aegean Sea. *Journal of Applied Ichthyology* 29: 1030-1035.
- Figueiredo M, Morato T, Barreiros JP, Afonso P, Santos RS (2005) Feeding ecology of the white seabream, *Diplodus sargus*, and the ballan wrasse, *Labrus bergylta*, in the Azores. *Fisheries Research* 75: 107-119.
- Fock HO, Matthiessen B, Zidowitz H, von Westernhagen H (2002) Diel and habitat-dependent resource utilisation by deep-sea fishes at the Great Meteor seamount: niche overlap and support for the sound scattering layer interception hypothesis. *Marine Ecology Progress Series* 244: 219-233.

- Fofandi MD, Vegd JA, Fofandi NM (2012) Diet composition and feeding strategy of bait caught skipjack tuna (*Katsuwonus pelamis*) caught along Saurashtra Coast. Environmental Research Journal 6: 121-123.
- Follesa MC, Mulas A, Cabiddu S, Porcu C, Deiana AM, Cau A (2010) Diet and feeding habits of two skate species, *Raja brachyura* and *Raja miraletus* (Chondrichthyes, Rajidae) in Sardinian waters (central-western Mediterranean). Italian Journal of Zoology 77: 53-60.
- Franz DR, Worley EK (1982) Seasonal variability of prey in the stomachs of *Astropecten americanus* (Echinodermata: Asteroidea) from off Southern New England, USA. Estuarine, Coastal and Shelf Science 14: 355-368.
- Freitas ICL (1998) Contribuição para o conhecimento da ecologia alimentar do peixe-espada preto, *Aphanopus carbo* Lowe, 1839 (Pisces: Trichiuridae), no Arquipélago da Madeira. Relatório de Estágio do Curso de Biologia, Universidade da Madeira, Funchal, 40 pp. (in Portuguese).
- Freitas L, Alves F, Ribeiro C, Dinis A, Nicolau C, Carvalho A (2014) Estudo técnico-científico de suporte à proposta de criação de áreas de operação para a actividade de whalewatching e respectiva capacidade de carga. Relatório técnico do Projecto CETACEOSMADEIRA II (LIFE07 NAT/P/000646), Museu da Baleia da Madeira, 87p.
- Freitas L, Dinis A, Alves F, Nóbrega F (2004) Cetaceans in the Madeira Archipelago (In Portuguese). Museu da Baleia, Machico, Madeira.
- Freitas L, Dinis A, Nicolau C, Ribeiro C, Alves F (2012) New records of cetacean species for Madeira Archipelago with an updated checklist. Boletim do Museu Municipal do Funchal 62: 25-43.
- Frick MG, Williams KL, Bolten AB, Bjorndal KA, Martins HR (2009) Foraging ecology of oceanic-stage loggerhead turtles *Caretta caretta*. Endangered Species Research 9: 91-97.
- Friedlander AM, Ballesteros E, Clemente S, Estep A, Gonçalves EJ, Rose P, Shepard M, Thompson C, Meeuwig JJ, Sala E (2016) Marine biodiversity and ecosystem health of Ilhas Selvagens, Portugal. Scientific Report to the Government of Portugal.
- Friedlander AM, Ballesteros E, Clemente S, Gonçalves EJ, Estep A, Rose P, Sala E (2017) Contrasts in the marine ecosystem of two Macaronesian islands: A comparison between the remote Selvagens Reserve and Madeira Island. PloS one 12: e0187935.

- Froese R, Pauly D (Eds) (2018) FishBase. World Wide Web electronic publication. [www.fishbase.org](http://www.fishbase.org), version (06/2018)
- García-Mederos AM, Tuset VM, Santana JI, Gonzalez JA (2010) Reproduction, growth and feeding habits of stout beardfish *Polymixia nobilis* (Polymixiidae) off the Canary Islands (NE Atlantic). *Journal of Applied Ichthyology* 26: 872-880.
- Garibaldi F, Orsi Relini L, Relini G (2010) Medusivorous fishes of the Mediterranean Sea 2. The specialist, *Schedophilus medusoiphagus* Cocco, 1839. *Rapp. Comm. Int. Mer Médit.* 39: 525
- Gartner Jr JV, Musick JA (1989) Feeding habits of the deep-sea fish, *Scopelogadus beanii* (Pisces: Melamphaide), in the western North Atlantic. *Deep Sea Research Part A. Oceanographic Research Papers* 36: 1457-1469.
- Gaskett AC, Bulman C, He X, Goldsworthy SD (2001) Diet composition and guild structure of mesopelagic and bathypelagic fishes near Macquarie Island, Australia. *NZ J Mar Freshw Res* 35: 469-476.
- Geraldes P (2000) Censos de Procelariformes na Ilha da Madeira. Época de nidificação 2000. Relatório final no âmbito do projecto "Novos Atlas das Aves Nidificantes em Portugal". SPNM e ICN.
- Giménez J, Marçalo A, Ramírez F, Verborgh P, Gauffier P, Esteban R, Nicolau L, González-Ortegón E, Baldó F, Vilas C, Vingada J, Forero MG, Stephanis R (2017) Diet of bottlenose dolphins (*Tursiops truncatus*) from the Gulf of Cadiz: Insights from stomach content and stable isotope analyses. *PloS one* 12: e0184673.
- Giménez J, Marçalo A, García-Polo M, García-Barón I, Castillo JJ, Fernández-Maldonado C, Saavedra C, Santos MB, Stephanis R (2018) Feeding ecology of Mediterranean common dolphins: The importance of mesopelagic fish in the diet of an endangered subpopulation. *Marine Mammal Science* 34: 136-154.
- Glaser SM, Waechter KE, Bransome NC (2015) Through the stomach of a predator: regional patterns of forage in the diet of albacore tuna in the California Current System and metrics needed for ecosystem-based management. *Journal of Marine Systems* 146: 38-49.
- Gonçalves JMS, Erzini K (1998) Feeding habits of the Two-banded sea bream (*Diplodus vulgaris*) and the Black sea bream (*Spondyllosoma cantharus*) (Sparidae) from the southwest coast of Portugal. *Cybio* 22(3): 245-254.

- Goñi N, Logan J, Arrizabalaga H, Jarry M, Lutcavage M (2011) Variability of albacore (*Thunnus alalunga*) diet in the Northeast Atlantic and Mediterranean Sea. *Marine biology* 158: 1057-1073.
- Gordon JDM, Nishida S, Nemoto T (1985) The diet of mesopelagic fish from the Pacific coast of Hokkaido, Japan. *Journal of the Oceanographical Society of Japan* 41: 89-97.
- Gordon JDM, Duncan JAR (1987) Deep-sea bottom-living fishes at two repeat stations at 2200 and 2900 m in the Rockall Trough, northeastern Atlantic Ocean. *Marine Biology* 96: 309-325.
- Gordon JDM, Mauchline J (1996) The distribution and diet of the dominant, slope-dwelling eel, *Synaphobranchus kaupii*, of the Rockall Trough. *Journal of the Marine Biological Association of the United Kingdom* 76: 493-503.
- Granadeiro JP (1993) Variation in measurements of Cory's Shearwater between populations and sexing by discriminant analysis. *Ringed & Migration* 14: 103-112.
- Granadeiro JP, Monteiro LR, Silva MC, Furness RW (2002) Diet of common terns in the Azores, Northeast Atlantic. *Waterbirds* 25: 149-155.
- Granadeiro JP, Dias MP, Rebelo R, Santos CD, Catry P (2006) Numbers and population trends of Cory's shearwater *Calonectris diomedea* at Selvagem Grande, Northeast Atlantic. *Waterbirds* 29: 56-60.
- Guénette S, Morato T (2001) The Azores archipelago in 1997. In: Guénette S, Christensen V, Pauly D (ed) *Fisheries impacts on North Atlantic ecosystems: models and analyses*. Fisheries Centre Research Reports Vol 9, University of British Columbia, p 241-270.
- Guerra A, Simon F, Gonzalez AF (1993) Cephalopods in the diet of the Swordfish, *Xiphias gladius*, from the Northeastern Atlantic Ocean. *Recent Advances in Fisheries Biology*, 159-164.
- Guschin AV, Corten A (2017) Feeding of pelagic fish in waters of Mauritania: 3.- Atlantic Chub mackerel *Scomber colias*, Atlantic horse mackerel *Trachurus trachurus*, Cunene horse mackerel *Trachurus trecae*. *Journal of Ichthyology* 57: 410-423.
- Habashi B, Wojciechowski J (1973) Observations on the biology of *Scomber japonicus* off northwest Africa. *Journal du Conseil International pour l'exploration de la mer (CIEM/ICES)* 20, 9pp.
- Harris MP (1966) Breeding biology of the Manx Shearwater *Puffinus puffinus*. *Ibis* 108: 17-33.



- Hernández E, Martín A, Nogales M, Quilis V, Delgado G, Trujillo O (1990) Distribution and status of Bulwer's Petrel (*Bulweria bulwerii* Jardine & Selby, 1828) in the Canary Islands. Bol. Mus. Mun. Funchal 42: 5–14.
- Hernández-García V, Martín V (1994) Stomach contents of two short-finned pilot whale (*Globicephala macrorhynchus* Gray, 1846) (Cetacea, Delphinidae) off the Canary Islands: a preliminary note. International Council for the Exploration of the Sea (ICES), 16.
- Hernández-García V (1995) The diet of the swordfish *Xiphias gladius* Linnaeus, 1758, in the central east Atlantic, with emphasis on the role of cephalopods. Fishery Bulletin 93: 403-411.
- Hernández-García V (2002) Contents of the digestive tract of a false killer whale (*Pseudorca crassidens*) stranded in Gran Canaria (Canary Islands, central east Atlantic). Bulletin of Marine Science 71: 367-369.
- Hirsch S, Christiansen B (2010) The trophic blockage hypothesis is not supported by the diets of fishes on Seine Seamount. Marine Ecology 31: 107-120.
- Hopkins TL, Baird RC (1985) Feeding ecology of four hatchetfishes (Sternoptychidae) in the eastern Gulf of Mexico. Bulletin of Marine Science 36: 260-277.
- Horn PL, Ballara SL, Sutton PJH, Griggs LH (2013) Evaluation of the diets of highly migratory species in New Zealand waters. New Zealand aquatic environment and biodiversity report 116, 140 pp.
- Horwood J (2009) Sei whale: *Balaenoptera borealis*. In Encyclopedia of Marine Mammals, Second Edition, pp 1001-1003.
- Houston KA, Haedrich RL (1986) Food habits and intestinal parasites of deep demersal fishes from the upper continental slope east of Newfoundland, northwest Atlantic Ocean. Marine Biology 92: 563-574.
- INCOGEO (2010) Informe preliminar de Campaña “INCOECO 0310”. Proyecto Life-INDEMARES. Instituto Español de Oceanografía, Centro Oceanográfico de Canarias. Memoria científico-técnica: 11 pp. + Anexos.
- Kadri H, Marouani S, Bradai MN, Bouaïn A (2014) Diet and feeding strategy of thornback ray, *Raja clavata* (Chondrichthyes: Rajidae) from the Gulf of Gabes (Tunisia—Central Mediterranean Sea). Journal of the Marine Biological Association of the United Kingdom 94: 1509-1516.

- Karakulak FS, Salman A, Oray IK (2009) Diet composition of bluefin tuna (*Thunnus thynnus* L. 1758) in the Eastern Mediterranean Sea, Turkey. *Journal of Applied Ichthyology* 25: 757-761.
- Karamanlidis AA, Dendrinos P, De Larrinoa PF, Gücü AC, Johnson WM, Kiraç CO, Pires R (2016) The Mediterranean monk seal *Monachus monachus*: status, biology, threats, and conservation priorities. *Mammal Review* 46: 92-105.
- Karamanlidis AA, Kallianiotis A, Psaradellis M, Adamantopoulou S (2011) Stomach contents of a subadult Mediterranean monk seal (*Monachus monachus*) from the Aegean Sea. *Aquatic Mammals* 37: 280.
- Kawamura A (1980) A review of food of balaenopterid whales. *Sci. Rep. Whales Res. Inst* 32: 155-197.
- Kato H, Perrin WF (2009) Bryde's whales: *Balaenoptera edeni/brydei*. In *Encyclopedia of Marine Mammals* (Second Edition).
- Klarian SA, Canales-Cerro C, Barría P, Zárate P, Concha F, Hernández S, Meléndez R (2018) New insights on the trophic ecology of blue (*Prionace glauca*) and shortfin mako sharks (*Isurus oxyrinchus*) from the oceanic eastern South Pacific. *Marine Biology Research* 14: 173-182.
- Kleisner K and Hoornaert C (2015) Primary Production. In: *Sea Around Us area parameters and definitions*. Website methods, [www.seaaroundus.org/sea-around-us-area-parameters-and-definitions/](http://www.seaaroundus.org/sea-around-us-area-parameters-and-definitions/)
- Klimpel S, Rückert S, Piatkowski U, Palm HW, Hanel R (2006) Diet and metazoan parasites of silver scabbard fish *Lepidopus caudatus* from the Great Meteor Seamount (North Atlantic). *Marine Ecology Progress Series* 315: 249-257.
- Konishi K, Tamura T, Isoda T, Okamoto R, Hakamada T, Kiwada H, Matsuoka K (2009) Feeding strategies and prey consumption of three baleen whale species within the Kuroshio-Current extension. *J North Atl Fish Sci* 42: 27-40.
- Labropoulou M, Kostikas I (1999) Patterns of resource use in deep-water decapods. *Marine Ecology Progress Series* 184: 171-182.
- Labropoulou M, Machias A, Tsimenides N, Eleftheriou A (1997) Feeding habits and ontogenetic diet shift on the striped mullet, *Mullus surmuletus* Linnaeus, 1758. *Fisheries Research* 31: 257-267.
- Labropoulou M, Papadopoulou-Smith K-N (1999) Foraging behaviour patterns of four sympatric demersal fishes. *Estuarine, Coastal and Shelf Science* 49: 99-108.
- Libralato S, Christensen V, Pauly D (2006) A method for identifying keystone species in

- food web models. *Ecol Modell* 195:153–171.
- Lorenzo JA, Alonso JRA, Tosco RB (2003) Atlas de las aves nidificantes en la isla de Tenerife.
- Lorenzo JA, Barone R (2007) Pardela cenicienta, *Calonectris borealis*. In: Lorenzo JA (ed.) Atlas de las aves nidificantes en el archipiélago canario (1997-2003). Dirección General de Conservación de la Naturaleza. Sociedad Española de Ornitología. Madrid.
- Lorenzo JM, Pajuelo JG 1995 Population biology of the roudi escolar *Promethichthys prometheus* (Gempylidae) off the Canary Islands. *Scientia Marina* 59: 235-240.
- Lorenzo JM, Pajuelo JG 1997 Biology of a deep benthopelagic fish, roudi escolar, *Promethichthys prometheus* (Gempylidae) off the Canary Islands. *Fish Bull* 1: 92-9.
- Lorenzo JA, Rodríguez B (2011) Estatus y distribución de los petreles y paños nidificantes en las islas Canarias. *Boletín del GIAM* 34: 31-33.
- López H, Pérez AJ, Rumeu B, Nogales M (2016) Trophic strategies of Yellow-legged Gull *Larus michahellis* on oceanic islands surrounded by deep waters. *Bird study* 63: 337-345.
- Luzardo J, López-Darias M, Suárez V, Calabuig P, García E, & Martín C (2008) First breeding population of Bulwer's petrel *Bulweria bulwerii* recorded on Grán Canaria (Canary islands)—population size and morphometric data. *Marine Ornithology* 36: 159-162.
- MacLeod CD, Santos MB, Pierce GJ (2003) Review of data on diets of beaked whales: evidence of niche separation and geographic segregation. *Journal of the Marine Biological Association of the United Kingdom* 83: 651-665.
- Macpherson E, Roel BA (1987) Trophic relationships in the demersal fish community off Namibia. *South African Journal of Marine Science* 5: 585-596.
- Malej A (1989) Behaviour and trophic ecology of the jellyfish *Pelagia noctiluca* (Forsskål, 1775). *Journal of Experimental Marine Biology and Ecology* 126: 259-270.
- Mancera-Rodríguez NJ, Castro-Hernández JJ (2015) Feeding ecology of the planehead filefish *Stephanolepis hispidus* (Pisces: Monacanthidae), in the Canary Islands area. *Revista de Biología Marina y Oceanografía* 50: 221-234.
- Martín A (1987) Atlas de las aves nidificantes en la isla de Tenerife. 1987. Instituto de Estudios Canarios. Monografía nº32.

- Martin B, Christiansen B (1997) Diets and standing stocks of benthopelagic fishes at two bathymetrically different midoceanic localities in the northeast Atlantic. *Deep Sea Research Part I: Oceanographic Research Papers* 44: 541-558.
- Martins HR (1985) A pygmy sperm whale, *Kogia breviceps* (Blainville, 1838) (Cetacea: Odontoceti) stranded on Faial Island, Azores, with notes on cephalopod beaks in stomach. *Ciencias Biológicas* 6: 63-69.
- Martins R, Ferreira C (1995) Line Fishing for Black Scabbardfish (*Aphanopus carbo* Lowe, 1839) and Other Deep-Water Species in the Eastern Mid Atlantic to the North of Madeira. In: Hopper A.G. (eds) *Deep-Water Fisheries of the North Atlantic Oceanic Slope*. NATO ASI Series (Series E: Applied Sciences), vol 296. Springer, Dordrecht.
- Marques A (1998) A note on the diet *Synaphobranchus kaupi* (Pisces: Synaphobranchidae) from the Porcupine Seabight, north-east Atlantic. *Journal of the Marine Biological Association of the United Kingdom* 78: 1385-1388.
- Massutí E, Deudero S, Sánchez P, Morales-Nin B (1998) Diet and feeding of dolphin (*Coryphaena hippurus*) in western Mediterranean waters. *Bulletin of Marine Science* 63: 329-341.
- Matthews FD, Damkaer DM, Knapp LW, Colette BB (1977) Food of the western north Atlantic tunas (*Thunnus*) and lancetfishes (*Alepisaurus*). *NOAA Tech. Rep. NMFS SSRF* 706: 1-19.
- Matthiessen B, Fock HO, Westernhagen H (2003) Evidence for two sympatric species of snipefishes *Macroramphosus* spp. (Syngnathiformes, Centriscidae) on Great Meteor Seamount. *Helgoland Marine Research* 57: 63-72.
- Matias R, Catry P (2010) The diet of Atlantic Yellow-legged Gulls (*Larus michahellis atlantis*) at an oceanic seabird colony: estimating predatory impact upon breeding petrels. *European Journal of Wildlife Research* 56: 861-869.
- Mauchline J, Gordon JDM (1980) The food and feeding of the deep-sea morid fish *Lepidion eques* (Gunther, 1887) in the Rockall Trough. *Journal of the Marine Biological Association of the United Kingdom* 60: 1053-1059.
- Mauchline J, Gordon JDM (1983a) Diets of the sharks and chimaeroids of the Rockall Trough, northeastern Atlantic Ocean. *Marine Biology* 75: 269-278.
- Mauchline J, Gordon JDM (1983b) Diets of clupeoid, stomiatoid and salmonoid fish of the Rockall Trough, northeastern Atlantic Ocean. *Marine Biology* 77: 67-78.

- Mauchline J, Gordon JDM (1984a) Diets and bathymetric distributions of the macrourid fish of the Rockall Trough, northeastern Atlantic Ocean. *Marine Biology* 81: 107-121.
- Mauchline J, Gordon JDM (1984b) Feeding and bathymetric distribution of the gadoid and morid fish of the Rockall Trough. *Journal of the Marine Biological Association of the United Kingdom* 64: 657-665.
- Mauchline J, Gordon JDM (1984c) Occurrence and feeding of berycomorphid and percomorphid teleost fish in the Rockall Trough. *ICES Journal of Marine Science* 41: 239-247.
- Maul GE (1961) The Ceratioid Fishes in the Collection of the Museu Municipal do Funchal (Melanocetidae, Himantolophidae, Oneirodidae, Linophrynidae). *Boletim do Museu Municipal do Funchal*. N°XIV, Art. 50: 87-159.
- Meirinho A, Barros N, Oliveira N, Catry P, Lecoq M, Paiva V, Geraldés P, Granadeiro JP, Ramírez I, Andrade J (2014) Atlas das Aves Marinhas de Portugal. Sociedade Portuguesa para o Estudo das Aves. Lisboa.
- Ménard F, Marchal E (2003) Foraging behaviour of tuna feeding on small schooling *Vinciguerria nimbaria* in the surface layer of the equatorial Atlantic Ocean. *Aquatic Living Resources* 16: 231-238.
- Mendizabal MG (2013) The reproductive biology, condition and feeding ecology of the skipjack, *Katsuwonus pelamis*, in the Western Indian Ocean. Universidad del País Vasco.
- Mondonça A, Arístegui J, Vilas JC, Montero MF, Ojeda A, Espino M, Martins A (2002) Is there a seamount effect on microbial community structure and biomass? The case study of Seine and Sedlo seamounts (Northeast Atlantic). *PLoS ONE* 7(1): e29526.
- Menezes D, Oliveira P, Ramírez I (2010) Pterodromas do arquipélago da Madeira. Duas espécies em recuperação. Serviço do Parque Natural da Madeira, Funchal.
- Menezes D, Oliveira P, Ramírez I (2011) Medidas Urgentes para a Recuperação da Freira do Bugio *Pterodroma feae* e do seu Habitat. Relatório Final. Serviço do Parque Natural da Madeira/ Sociedade Portuguesa para o Estudo das Aves.
- Mendonça A (2009) Diet of the blue shark, *Prionace glauca*, in the Northeast Atlantic. Master's Thesis, University of Porto, Portugal.
- Merrett NR, Domanski PA (1985) Observations on the ecology of deep-sea bottom-living fishes collected off northwest Africa: II. The Moroccan slope (27–34 N), with special reference to *Synaphobranchus kaupii*. *Biological Oceanography* 3: 349-399.

- Merrett N, Roe HSJ (1974) Patterns and selectivity in the feeding of certain mesopelagic fishes. *Marine Biology* 28: 115-126.
- Mintzer VJ, Gannon DP, Barros NB, Read AJ (2008) Stomach contents of mass-stranded short-finned pilot whales (*Globicephala macrorhynchus*) from North Carolina. *Marine Mammal Science* 24: 290-302.
- Monteiro LR, Ramos JA, Furness RW, Del Nevo AJ (1996) Movements, morphology, breeding, molt, diet and feeding of seabirds in the Azores. *Colonial Waterbirds* 82-97.
- Morais L, Santos C, Vicente L (1998) Population increase of Yellow-legged gulls *Larus cachinnans* breeding on Berlenga island (Portugal), 1974-1994. *Sula* 12: 27-38
- Morato T, Solà E, Grós MP, Menezes G, Pinho MR (1998) Trophic relationships and feeding habits of demersal fishes from the Azores: importance to multispecies assessment. *ICES CM 1998/O: 7*. 21 pp.
- Morato T, Solà E, Grós MP, Menezes G (1999) Diets of forkbeard (*Phycis phycis*) and Conger eel (*Conger conger*) off the Azores during spring of 1996 and 1997. *Archipelago. Life and Marine Sciences* 17A: 51-64.
- Morato T, Pitcher T (2002) Challenges and problems in modelling seamount ecosystems and their fisheries. *ICES 2002/M: 8*, 28pp.
- Morato T, Solà E, Grós MP, Menezes G (2003) Diets of thornback ray (*Raja clavata*) and tope shark (*Galeorhinus galeus*) in the bottom longline fishery of the Azores, northeastern Atlantic. *Fishery Bulletin* 101: 590-602.
- Morato T, Bulman C, Pitcher, TJ (2009) Modelled effects of primary and secondary production enhancement by seamounts on local fish stocks. *Deep Sea Research Part II: Topical Studies in Oceanography* 56: 2713-2719.
- Morato T, Lemey E, Menezes G, Pham CK, Brito J, Soszynski A, Pitcher TJ, Heymans JJ (2016) Food-web and ecosystem structure of the open-ocean and deep-sea environments of the Azores, NE Atlantic. *Frontiers in Marine Science* 3: 245.
- Moreno T, Castro J (1995) Community structure of the juvenile of coastal pelagic fish species in the Canary Islands Waters. *Scientia Marina* 59: 405-413.
- Morissette L, Kaschner K, Gerber LR (2010) Ecosystem models clarify the trophic role of whales off Northwest Africa. *Marine Ecology Progress Series* 404: 289-302.
- Morte S, Redon MJ, Sanz-Brau A (2001) Diet of *Scorpaena porcus* and *Scorpaena notata* (Pisces: Scorpaenidae) in the western Mediterranean. *Cah. Biol. Mar.* 42: 333-344.

- Mostarda E, Campo D, Castriota L, Esposito V, Scarabello MP, Andaloro F (2007) Feeding habits of the bullet tuna *Auxis rochei* in the southern Tyrrhenian Sea. *Journal of the Marine Biological Association of the United Kingdom* 87: 1007-1012.
- Moteki M, Arai M, Tsuchiya K, Okamoto H (2001) Composition of piscine prey in the diet of large pelagic fish in the eastern tropical Pacific Ocean. *Fisheries science* 67: 1063-1074.
- Mougin JL, Jouanin C (1997) Prospection alimentaire du Puffin cendré *Calonectris diomedea borealis* de Selvagem Grande (30 09' N, 15 52' W) pendant l'incubation, par télémétrie satellitaire. *Comptes Rendus de l'Académie des Sciences-Series III-Sciences de la Vie* 320: 825-831.
- Nakamura EL (1965) Food and feeding habits of skipjack tuna (*Katsuwonus pelamis*) from the Marquesas and Tuamotu Islands. *Transactions of the American Fisheries Society* 94: 236-242.
- Navarro J, López L, Coll M, Barría C, Sáez-Liante R (2014) Short-and long-term importance of small sharks in the diet of the rare deep-sea shark *Dalatias licha*. *Marine Biology* 161: 1697-1707.
- Neiva J, Coelho R, Erzini K (2006) Feeding habits of the velvet belly lanternshark *Etmopterus spinax* (Chondrichthyes: Etmopteridae) off the Algarve, southern Portugal. *Journal of the Marine Biological Association of the United Kingdom* 86: 835-841.
- Nicolau C, Alves F, Ferreira R, Henriques F, Carvalho A, Cunha I, Freitas L (2014) Surveillance of the conservation status of cetaceans species in Madeira offshore waters (Deliverable A.8\_I). Relatório técnico do Projecto CETÁCEOSMADEIRA II (LIFE07 NAT/P/000646). Museu da Baleia da Madeira.
- Novotny M (2018) The Assemblage Structure and Trophic Ecology of a Deep-Pelagic Fish Family (Platyroctidae) in the Gulf of Mexico. Master's thesis. Nova Southeastern University.
- Nunes M (2000) New data on the Bulwer's Petrel (*Bulweria bulwerii*) breeding biology in the Desertas Islands (Madeira archipelago). *Life and Marine Sciences Suppl.* 2 (Part A): 167-173.
- Nunes J, Nunes M, Fagundes AI, Valkenburg T (2010) Contributo para a conservação do Fura-bucho-do-atlântico *Puffinus puffinus*, uma espécie ameaçada na ilha da Madeira. *Airo* 20: 12-21.

- OAG (2013) Estado de conservación de la tortuga boba (*Caretta caretta*) en las islas Canarias. Observatorio Ambiental Granadilla Technical Paper. OAG, Santa Cruz de Tenerife, 154pp
- Olaso I, Cendrero O, Abaúnza P (1999) The diet of the horse mackerel, *Trachurus trachurus* (Linnaeus 1758), in the Cantabrian Sea (north of Spain). J. Appl. Ichthyol. 15: 193-198.
- Oliveira P & Menezes D (2004) Aves do Arquipélago da Madeira. Serviço do Parque Natural da Madeira/Arquipélago Verde produtos promocionais, Funchal, Madeira.
- Olson RJ, Galván-Magaña F (2002) Food habits and consumption rates of common dolphinfish (*Coryphaena hippurus*) in the eastern Pacific Ocean. Fishery Bulletin 100: 279-298.
- Olson RJ, Duffy LM, Kuhnert PM, Galvan-Magana F, Bocanegra-Castillo N, Alatorre-Ramirez V (2014) Decadal diet shift in yellowfin tuna *Thunnus albacares* suggests broad-scale food web changes in the eastern tropical Pacific Ocean. Marine Ecology Progress Series 497: 157-178.
- Opitz S (1993) A quantitative model of the trophic interactions in a Caribbean coral reef ecosystem. Pp: 259-267 in V. Christensen and D. Pauly (eds.) Trophic models of aquatic ecosystems. ICLARM Conf. 26, 390 p.
- Oro D, de León A, Minguez E, Furness RW (2005) Estimating predation on breeding European storm-petrels (*Hydrobates pelagicus*) by yellow-legged gulls (*Larus michahellis*). Journal of Zoology 265: 421-429.
- Öztürk B, Salman, A, Öztürk A A, Tonay A (2007) Cephalopod remains in the diet of the Striped dolphins (*Stenella coeruleoalba*) and Risso's dolphins (*Grampus griseus*) in the eastern Mediterranean Sea. Vie et milieu – Life and Environment 57(1/2): 53-59.
- Pais C (2002) Diet of a deep-sea fish, *Hoplostethus mediterraneus*, from the south coast of Portugal. Journal of the Marine Biological Association of the United Kingdom 82: 351-352.
- Paiva VH, Xavier J, Geraldés P, Ramirez I, Garthe S, Ramos JA (2010) Foraging ecology of Cory's shearwaters in different oceanic environments of the North Atlantic. Marine Ecology Progress Series 410: 257-268.
- Paiva VH, Fagundes AI, Romão V, Gouveia C, Ramos JA (2016) Population-scale foraging segregation in an apex predator of the North Atlantic. PloS one 11: e0151340.



- Pajuelo JG, Lorenzo JM (1998) Population biology of the common pandora *Pagellus erythrinus* (Pisces: Sparidae) off the Canary Islands. Fisheries Research 36: 75-86.
- Pajuelo JG, JM Lorenzo (2000) Reproduction, age, growth and mortality of axillary seabream, *Pagellus acarne* (Sparidae), from the Canarian Archipelago. J. Appl. Ichthyol. 16: 41-47.
- Pajuela JG, González JA, Santan JI (2010) Bycatch and incidental catch of the black scabbardfish (*Aphanopus* spp.) fishery off the Canary Islands. Fisheries Research 106: 448-453.
- Pauly D, Bartz MLS, Palomares MLD (1993) Improved construction, parametrization and interpretation of steady-state ecosystem models, pp 1-13. In: Christensen V, Pauly D (Eds) Trophic models of aquatic ecosystems. ICLARM Conference Proceedings 26, 390p.
- Parry M (2006) Feeding behaviour of two ommastrephid squids *Ommastrephes bartramii* and *Sthenoteuthis oualaniensis* off Hawaii. Marine Ecology Progress Series 318: 229-235.
- Passarella KC, Hopkins TL (1991) Species composition and food habitats of the micronektonic cephalopod assemblage in the eastern Gulf of Mexico. Bulletin of Marine Science 49(1-2): 638-659.
- Pedro PI, Ramos JA, Neves VC, Paiva VH (2013) Past and present trophic position and decadal changes in diet of Yellow-legged Gull in the Azores Archipelago, NE Atlantic. European Journal of Wildlife Research 59: 833-845.
- Perrin WF, Robertson KM, Walker WA (2008) Diet of the Striped Dolphin, *Stenella coeruleoalba*, in the Eastern Tropical Pacific Ocean.
- Perrin WF (2009) Atlantic Spotted Dolphin: *Stenella frontalis*. In Encyclopedia of Marine Mammals (Second Edition) (pp. 54-56).
- Perrins CM, Harris MP, Britton CK (1973) Survival of Manx shearwaters *Puffinus puffinus*. Ibis 115: 535-548.
- Pethybridge H, Daley RK, Nichols PD (2011) Diet of demersal sharks and chimaeras inferred by fatty acid profiles and stomach content analysis. Journal of Experimental Marine Biology and Ecology 409: 290-299.
- Piatkowski U, Hernández-García V, Clarke MR (1998) On the biology of the European Flying squid *Todarodes sagittatus* (Lamarck, 1798) (Cephalopoda, Ommastrephidae) in the Central Eastern Atlantic.

- Pierce GJ, Hernandez-Milian G, Santos MB, Dendrinos P, Psaradellis M, Tounta E, Androukaki E, Edridge A (2011) Diet of the monk seal (*Monachus monachus*) in Greek waters. *Aquatic Mammals* 37: 284.
- Pierrot-Bults AC, Van der Spoel S (1998) Pelagic Biogeography ICoPB II. In Proceedings of the 2nd International Conference. IOC Workshop Report (Vol. 142, pp. 1-387).
- Pinheiro PB, Júnior TV, Hazin FHV, Travassos P, Tolotti MT, Barbosa TM (2010) Diet of the white marlin (*Tetrapturus albidus*) from the southwestern equatorial Atlantic Ocean. *Collect. Vol. Sci. Pap. ICCAT* 65: 1843-1850.
- Pinkerton MH (2011) A balanced trophic model of the Chatham Rise, New Zealand, pp 60.
- Pires R, Neves HC, Karamanlidis AA (2008) The critically endangered Mediterranean monk seal *Monachus monachus* in the archipelago of Madeira: priorities for conservation. *Oryx* 42: 278-285.
- Pires R (2011) Lobos-marinhos do arquipélago da Madeira. Edições do Serviço do Parque Natural da Madeira, Funchal, Madeira.
- Piroddi C, Coll M, Steenbeek J, Moy DM, Christensen V (2015) Modelling the Mediterranean marine ecosystem as a whole: addressing the challenge of complexity. *Marine Ecology Progress Series* 533: 47-65
- Podrazhanskaya SG (1993) Feeding habits of Mesopelagic species of fish and estimation of plankton graze in the northwest Atlantic. *NAFO Scientific Council Studies* 19: 79-85.
- Porsmoguer SB, Banaru D, Béarez P, Dekeyser I, Merchán Fornelino M, Míguez Lozano R, Boudouresque CF (2013) Diet of the shortfin Mako *Isurus oxyrinchus* and the blue shark *Prionace glauca* in the northeastern Atlantic Ocean. *Rapp Comm Int Mer Médit*, 40: 499.
- Porsmoguer SB, Bănar D, Bearez P, Dekeyser I, Fornelino MM, Boudouresque CF (2014) Unexpected headless and tailless fish in the stomach content of shortfin mako *Isurus oxyrinchus*. *PloS one* 9: e88488.
- Porsmoguer SB, Banaru D, Boudouresque CF, Dekeyser I, Viricel A, Merchán M (2015) DNA evidence of the consumption of short-beaked common dolphin *Delphinus delphis* by the shortfin mako shark *Isurus oxyrinchus*. *Marine Ecology Progress Series* 532: 177-183.

- Porsmoguer SB, Banaru D, Boudouresque CF, Dekeyser I, Béarez P, Míguez Lozano R (2017) Compared diet of two pelagic shark species in the Northeastern Atlantic Ocean. *Vie et Milieu-Life and Environment* 67: 21-25.
- Porteiro FM, Barreiros JP, Santos RS (1996) Wrasses (Teleostei: Labridae) of the Azores. *ARQUIPÉLAGO. Ciências Biológicas e Marinhas= Life and Marine Sciences* 14: 23-40.
- Potier M, Marsac F, Lucas V, Sabatié R, Hallier JP, Ménard F (2004) Feeding partitioning among tuna taken in surface and mid-water layers: the case of yellowfin (*Thunnus albacares*) and bigeye (*T. obesus*) in the western tropical Indian Ocean. *Western Indian Ocean Journal of Marine Science* 3: 51-62.
- Potier M, Marsac F, Cherel Y, Lucas V, Sabatié R, Maury O, Ménard F (2007) Forage fauna in the diet of three large pelagic fishes (lancetfish, swordfish and yellowfin tuna) in the western equatorial Indian Ocean. *Fisheries Research* 83: 60-72.
- Preti A, Kohin S, Dewar H, Ramon D (2008) Feeding habits of the Bigeye Thresher shark (*Alopias supercilliosus*) sampled from the California-based drift gillnet fishery. *CalCOFI rep.* 49, 10 p.
- Preti A, Soykan CU, Dewar H, Wells RD, Spear N, Kohin S (2012) Comparative feeding ecology of shortfin mako, blue and thresher sharks in the California Current. *Environmental Biology of Fishes* 95: 127-146.
- Prokofiev AM, Kukuev EI (2009) Systematics and distribution of black swallows of the genus *Chiasmodon* (Perciformes: Chiasmodontidae). *Journal of Ichthyology* 49: 899.
- Pusineri C, Vasseur Y, Hassani S, Meynier L, Spitz J, Ridoux V (2005) Food and feeding ecology of juvenile albacore, *Thunnus alalunga*, off the Bay of Biscay: a case study. *ICES Journal of Marine Science* 62: 116-122.
- Pusineri C, Magnin V, Meynier L, Spitz J, Hassani S, Ridoux V (2007) Food and feeding ecology of the common dolphin (*Delphinus delphis*) in the oceanic Northeast Atlantic and comparison with its diet in neritic areas. *Marine Mammal Science* 23: 30-47.
- Ramírez I, Paiva VH, Menezes D, Silva I, Phillips RA, Ramos JA, Garthe S (2013) Year-round distribution and habitat preferences of the Bugio petrel. *Marine Ecology Progress Series* 476: 269-284.
- Ramos AG, Lorenzo, Pajuelo (1995) Food habits of bait-caught skipjack tuna *Katsuwonus pelamis* off the Canary Islands. *Scientia Marina* 59: 65-69.

- Ramos R, Granadeiro JP, Nevoux M, Mougin JL, Dias MP, Catry P (2012) Combined spatio-temporal impacts of climate and longline fisheries on the survival of a trans-equatorial marine migrant. *PloS one* 7: e40822.
- Ramos R, Granadeiro JP, Rodríguez B, Navarro J, Paiva VH, Bécares J, Reyes-González JM, Fagundes I, Ruiz A, Arcos P, González-Solís J, Catry P (2013) Metapopulation feeding grounds of Cory's shearwater in the subtropical Atlantic Ocean: implications for the definition of Marine Protected Areas based on tracking studies. *Diversity and Distributions* 19: 1284-1298.
- Ramos JA, Fagundes AI, Xavier JC, Fidalgo V, Ceia FR, Medeiros R, Paiva VH (2015a) A switch in the Atlantic Oscillation correlates with inter-annual changes in foraging location and food habits of Macaronesian shearwaters (*Puffinus baroli*) nesting on two islands of the sub-tropical Atlantic Ocean. *Deep Sea Research Part I: Oceanographic Research Papers* 104: 60-71.
- Ramos R, Sanz V, Militão T, Bried J, Neves VC, Biscoito M, Phillips RA, Zino F, González-Solís J (2015b) Leapfrog migration and habitat preferences of a small oceanic seabird, Bulwer's petrel (*Bulweria bulwerii*). *Journal of Biogeography* 42: 1651-1664.
- Ramos R, Ramírez I, Paiva VH, Militão T, Biscoito M, Menezes D, Phillips RA, Zino F, González-Solís J (2016) Global spatial ecology of three closely-related gadfly petrels. *Scientific reports* 6: 23447.
- Ramos JJ, Trujillo D (2004) Petrel de Bulwer *Bulweria bulwerii*. *Libro Rojo de las aves de España*. Madrid, Spain: Dirección General para la Biodiversidad-SEO/BirdLife, 37-39.
- Renones O, Polunin NVC, Goni R (2002) Size related dietary shifts of *Epinephelus marginatus* in a western Mediterranean littoral ecosystem: an isotope and stomach content analysis. *Journal of fish biology* 61: 122-137.
- Reyes-González, Rodríguez, Arcos, Bécares (2011) Campaña de marcaje de SEO/BirdLife-UB: Pardela cenicienta – GPS. Campañas oceanográficas Proyecto LIFE+ INDEMARES. Veneguera, Canarias.
- Reyes-González, Rodríguez, Arcos (2012) Campaña de marcaje de SEO/BirdLife: Pardela cenicienta – GPS. Campañas oceanográficas Proyecto LIFE+ INDEMARES. Veneguera, Canarias.
- Ringelstein J, Pusineri C, Hassani S, Meynier L, Nicolas R, Ridoux V (2006) Food and feeding ecology of the striped dolphin, *Stenella coeruleoalba*, in the oceanic waters

- of the north-east Atlantic. Journal of the Marine Biological Association of the United Kingdom 86: 909-918.
- Robertson HA, James PC (1988) Morphology and egg measurements of seabirds breeding on Great Salvage Island, North Atlantic. Bull BOC 108: 79-87.
- Rodríguez B, de León L, Martín A, Alonso J, Nogales M (2003) Status and distribution of breeding seabirds in the northern islets of Lanzarote, Canary Islands. Atlantic seabirds 5: 41-56.
- Rodríguez, Arcos (2010) Censo de aves marinas en el Banco de La Concepción: Informe de SEO/BirdLife para la campaña Banco de la Concepción - INCOGEO (IEO).
- Rodríguez, Arcos, Bécares (2010) Campaña de marcaje de SEO/BirdLife: pardela cenicienta – GPS. Campañas oceanográficas Proyecto LIFE+ INDEMARES. Alegranza, Canarias.
- Rodríguez, Arcos (2012) Campaña de marcaje de SEO/BirdLife: Pardela cenicienta – GPS. Campañas oceanográficas Proyecto LIFE+ INDEMARES. Parque Nacional de Timanfaya, Lanzarote.
- Rodríguez B, Bécares J, Martínez JM, Rodríguez A, Ruiz A, Arcos JM (2013) Satellite tracking of Bulwer's Petrels *Bulweria bulwerii* in the Canary Islands. Bird Study 60 270-274.
- Roe HSJ, Badcock J (1984) The diel migrations and distributions within a mesopelagic community in the North East Atlantic. 5. Vertical migrations and feeding of fish. Progress in Oceanography 13: 389-424.
- Roger C (1994) Relationships among yellowfin and skipjack tuna, their prey-fish and plankton in the tropical western Indian Ocean. Fisheries Oceanography 3: 133-141.
- Romero J, Catry P, Menezes D, Coelho N, Silva JP, Granadeiro JP (2019) A gull that scarcely ventures on the ocean: Yellow-legged gull *Larus michahellis atlantis* on the oceanic island of Madeira. Ardeola 66(1): 101-112.
- Ronconi RA, Schoombie S, Westgate AJ, Wong SN, Koopman HN, Ryan PG (2018) Effects of age, sex, colony and breeding phase on marine space use by Great Shearwaters *Ardenna gravis* in the South Atlantic. Marine Biology 165: 58.
- Rosa S, Pansera M, Granata A, Guglielmo L (2013) Interannual variability, growth, reproduction and feeding of *Pelagia noctiluca* (Cnidaria: Scyphozoa) in the Straits of Messina (Central Mediterranean Sea): Linkages with temperature and diet. Journal of Marine Systems 111: 97-107.

- Rosas-Luis R, Navarro J, Loor-Andrade P, Forero MG (2017) Feeding ecology and trophic relationships of pelagic sharks and billfishes coexisting in the central eastern Pacific Ocean. *Marine Ecology Progress Series* 573: 191-201.
- Rosecchi E, Tracey DM, Webber WR (1988) Diet of orange roughy, *Hoplostethus atlanticus* (Pisces: Trachichthyidae) on the Challenger Plateau, New Zealand. *Marine Biology* 99: 293-306.
- Runderhausen PJ, Buckel JA, Edwards J, Gannon DP, Butler CM, Averett TW (2010) Feeding ecology of blue marlins, dolphinfish, yellowfin tuna, and wahoos from the North Atlantic Ocean and comparisons with other oceans. *Transactions of the American Fisheries Society* 139: 1335-1359.
- Saglam H, Bascinar NS (2008) Feeding ecology of thornback ray (*Raja clavata* Linnaeus, 1758) on the Turkish coast of the south-eastern Black Sea. *Marine Biology Research* 4: 451-457.
- Saldanha L, Almeida AJ, Andrade F, Guerreiro J (1995) Observations on the diet of some slope dwelling fishes of southern Portugal. *Internationale Revue der gesamten Hydrobiologie und Hydrographie* 80: 217-234.
- Sallami B, Ben Salem M, Reynaud C, Capapé C (2014) Diet of Mediterranean moray, *Muraena helena* (Actinopterygii, Anguilliformes, Muraenidae), from the north-eastern Tunisian coast (central Mediterranean). *Acta Ichthyologica et Piscatoria* 44: 273-283.
- Sambolino A, Alves F, Correia AM, Ferreira R, Carvalho P, Silva G, Dinis A (2017) Monitoring cetaceans in the Madeira Archipelago from a ferry along a fixed transect
- Sánchez F, Olaso I (2004) Effects of fisheries on the Cantabrian Sea shelf ecosystem. *Ecological Modelling* 172: 151-174.
- Santos J, Araújo H, Ferreira M, Henriques A, Miodonski J, Monteiro S, Oliveira I, Rodrigues P, Duro G, Oliveira F, Pinto N, Sequeira M, Eira C, Vingada J (2012) Chapter I: Baseline estimates of abundance and distribution of target species. Annex to the Midterm Report of project LIFE MarPro NAT/PT/00038.
- Santos MB, Martin V, Arbelo M, Fernández A, Pierce GJ (2007) Insights into the diet of beaked whales from the atypical mass stranding in the Canary Islands in September 2002. *Journal of the Marine Biological Association of the United Kingdom* 87: 243-251.
- Santos MB, Pierce GJ, Herman J, Lopez A, Guerra A, Mente E, Clarke MR (2001) Feeding ecology of Cuvier's beaked whale (*Ziphius cavirostris*): a review with new

- information on the diet of this species. *Journal of the Marine Biological Association of the United Kingdom* 81: 687-694.
- Santos MB, Pierce GJ, Lopez A, Reid RJ, Ridoux V, Mente E (2006) Pygmy sperm whales *Kogia breviceps* in the Northeast Atlantic: New information on stomach contents and strandings. *Marine Mammal Science* 22: 600-616.
- Santos AR, Trueman C, Connolly P, Rogan E (2013) Trophic ecology of black scabbardfish, *Aphanopus carbo* in the NE Atlantic-assessment through stomach content and stable isotope analyses. *Deep Sea Research Part I: Oceanographic Research Papers* 77: 1-10.
- Satoh K, Yokawa K, Saito H, Matsunaga H, Okamoto H, Uozumi Y (2004) Preliminary stomach contents analysis of pelagic fish collected by Shoyo-Maru 2002 research cruise in the Atlantic Ocean. *Collect. Vol. Sci. Pap. ICCAT* 56: 1096-1114.
- Sedberry GR, Musick JA (1978) Feeding strategies of some demersal fishes of the continental slope and rise off the mid-Atlantic coast of the USA. *Marine Biology* 44: 357-375.
- Sekiguchi K, Klages NTW, Best PB (1992) Comparative analysis of the diets of smaller odontocete cetaceans along the coast of southern Africa. *South African Journal of Marine Science* 12: 843-861.
- Sever TM, Filiz H, Bayhan B, Taskavak E, Bilge G (2008) Food habits of the hollowsnout grenadier, *Caelorinchus caelorhincus* (Risso, 1810), in the Aegean Sea, Turkey. *Belgian Journal of Zoology* 138: 81-84.
- Shimose T, Shono H, Yokawa K, Saito H, Tachihara K (2006) Food and feeding habits of blue marlin, *Makaira nigricans*, around Yonaguni Island, southwestern Japan. *Bulletin of Marine Science* 79: 761-775.
- Shoji A, Aris-Brosou S, Fayet A, Padget O, Perrins C, Guilford T (2015) Dual foraging and pair coordination during chick provisioning by Manx shearwaters: empirical evidence supported by a simple model. *Journal of Experimental Biology* 218: 2116-2123.
- Silva FVA, Pinheiro PB, Bezerra NPA, Vaske-Junior T, Filho ALV, Hazin FH (2014) Hábito alimentar do Agulhão-Verde, *Tetrapturus pfluegeri* (Robins & de Sylva, 1963), (Teleostei-Istiophoridae), capturado no Atlântico Oeste Tropical. *Tropical Oceanography* 42: 128-36.
- Soares MS, Sousa L, Barreiros JP (2003) Feeding habits of the lizardfish *Synodus saurus* (Linnaeus, 1758). *Aqua - Journal of Ichthyology and Aquatic Biology* 7: 29-38.

- Spear LB, Ainley DG, Walker WA (2007) Foraging dynamics of seabirds in the eastern tropical Pacific Ocean. *Studies in Avian Biology* 35: 1–99.
- Spitz J, Mourocq E, Schoen V, Ridoux V (2010) Proximate composition and energy content of forage species from the Bay of Biscay: high-or low-quality food? *ICES Journal of Marine Science* 67: 909-915.
- SPNM (2015a) Planos de Ação das 14 espécies alvo do projeto LIFE ILHÉUS DO PORTO SANTO.
- SPNM (2015b) LIFE RECOVER NATURA - Midterm Report Covering the project activities from 01/10/2013 to 30/11/2015
- Sutton TT, Hopkins TL (1996) Trophic ecology of the stomiid (Pisces: Stomiidae) fish assemblage of the eastern Gulf of Mexico: strategies, selectivity and impact of a top mesopelagic predator group. *Marine Biology* 127: 179-192.
- Sutton TT (2005) Trophic ecology of the deep-sea fish *Malacosteus niger* (Pisces: Stomiidae): An enigmatic feeding ecology to facilitate a unique visual system? *Deep Sea Research Part I: Oceanographic Research Papers* 52: 2065-2076.
- Teixeira CM, Pinheiro A, Cabral HN (2009) Feeding ecology, growth and sexual cycle of the sand sole, *Solea lascaris*, along the Portuguese coast. *J. Mar. Biol. Ass. U.K.* 89(3): 621-627.
- Teixeira CM, Batista MI, Cabral HN (2010) Diet, growth and reproduction of four flatfishes on the Portuguese coast. *Scientia Marina* 74(2): 223-233.
- Tershy BR, Acevedo-Gutiérrez A, Breese D, Strong C (1993) Diet and feeding behavior of fin and Bryde's whales in the central Gulf of California, Mexico. *Revista de Investigación Científica* 1: 31-38.
- Thomas PT (1964) Food of *Katsuwonus pelamis* (Linnaeus) and *Neothunnus macropterus* (Temminck and Schlegel) from Minicoy waters during the season 1960-61.
- Thompson KR (1987) The ecology of the Manx shearwater *Puffinus puffinus* on Rhum, West Scotland (Doctoral dissertation, University of Glasgow).
- Tilves U, Purcell JE, Fuentes VL, Torrents A, Pascual M, Raya V, Sabatés A (2016) Natural diet and predation impacts of *Pelagia noctiluca* on fish eggs and larvae in the NW Mediterranean. *Journal of Plankton Research* 38: 1243-1254.
- Tonay AM, Danyer E, Dede A, Öztürk B, Öztürk AA (2016) The stomach content of a Mediterranean Monk Seal (*Monachus monachus*): finding of Green Turtle (*Chelonia mydas*) remains. *Zoology in the Middle East* 62: 212-216.



- Torres MÁ, Coll M, Heymans JJ, Christensen V, Sobrino I (2013) Food-web structure of and fishing impacts on the Gulf of Cadiz ecosystem (South-western Spain). *Ecological modelling* 265: 26-44.
- Tripp-Valdez A, Galvan-Magaña F, Ortega-Garcia S (2015) Food sources of common dolphinfish (*Coryphaena hippurus*) based on stomach content and stable isotopes analyses. *Journal of the Marine Biological Association of the United Kingdom* 95: 579-591.
- Trites AW, Pauly D (1998) Estimating mean body masses of marine mammals from maximum body lengths. *Canadian Journal of Zoology* 76: 886-896.
- Uchikawa K, Kitagawa D, Sakurai Y (2001a) Notes on Feeding Habits of the Mesopelagic Fish *Maurolicus japonicus* off the Pacific Coast of Northern Japan. *Bull. Fac. Fish. Hokkaido Univ.* 52: 151-156.
- Uchikawa K, Yamamura O, Sakurai Y (2001b) Feeding habits of the mesopelagic fish *Gonostoma gracile* in the northwestern North Pacific. *Journal of oceanography* 57: 509-517.
- Vasconcelos J, Sousa R, Riera R, Delgado J, Faria G, Gordo LS (2018) Stock assessment of the blue jack mackerel, *Trachurus picturatus*, in the North-eastern Atlantic. *Fish. Manag. Ecol* 25: 233-239.
- Vasilakopoulos P, Pavlidis M, Tserpes G (2011) On the diet and reproduction of the oilfish *Ruvettus pretiosus* (Perciformes: Gempylidae) in the eastern Mediterranean. *Journal of the Marine Biological Association of the United Kingdom* 91: 873-881.
- van Nierop MM, den Hartog JC (1984) A study on the gut contents of five juvenile loggerhead turtles, *Caretta caretta* (Linnaeus) (Reptilia, Cheloniidae), from the south-eastern part of the North Atlantic Ocean, with emphasis on Coelenterate identification. *Zoologische Mededelingen* 59: 35-54.
- Vaske-Júnior T, Vooren CM, Lessa RP (2004) Feeding habits of four species of Istiophoridae (Pisces: Perciformes) from northeastern Brazil. *Environmental Biology of Fishes* 70: 293-304.
- Veiga P, Xavier JC, Assis CA, Erzini K (2011) Diet of the blue marlin, *Makaira nigricans*, off the south coast of Portugal. *Marine Biology Research* 7: 820-825.
- Viana DDL, Tolotti MT, Porto M, Araújo RJVD, Vaske-Júnior T, Hazin FHV (2012) Diet of the oilfish *Ruvettus pretiosus* (Perciformes: gempylidae) in the Saint Peter and saint Paul Archipelago, Brazil. *Brazilian Journal of Oceanography* 60: 181-188.

- Waap S (2015) Trophic relationships among pelagic predators of the deep seas of the Madeira islands. PhD thesis, Cardiff University
- Waap S, Symondson WOC, Granadeiro JP, Alonso H, Serra-Gonçalves C, Dias MP, Catry P (2017) The diet of a nocturnal pelagic predator, the Bulwer's petrel, across the lunar cycle. *Scientific Reports* 7: 1384.
- Wahbi F, Tojo N, Ramzi A, Somoue L, Manchi K, Errhif A (2015) Seasonal and size-dependent variability in diet of *Scomber colias* (Gmelin, 1789) of the Atlantic Coast of the Northwest Africa. *International Journal* 3: 485-497.
- Warham J (1996) The behaviour, population biology and physiology of the petrels. Academic Press, London.
- Watanabe K, Kubodera T, Ichii T, Sakai M, Moku M, Seitou M (2008) Diet and sexual maturation of the neon flying squid *Ommastrephes bartramii* during autumn and spring in the Kuroshio-Oyashio transition region. *J. mar. biol. Ass. U.K.* 88(2): 381-389.
- Weiss R (1974) Feeding behaviour and formation of fish concentrations in the chub mackerel (*Scomber colias*) in the Northwest African fishing grounds. *ICES Doc CM* 1974/J:15, 19pp.
- West K, Walker W, Baird R, White W, Levine G, Brown E, Schofield D (2009) Diet of pygmy sperm whales (*Kogia breviceps*) in the Hawaiian Archipelago. *Marine Mammal Science* 25: 931-943.
- West KL, Walker WA, Baird RW, Mead JG, Collins PW (2017) Diet of Cuvier's beaked whales *Ziphius cavirostris* from the North Pacific and a comparison with their diet world-wide. *Marine Ecology Progress Series* 574: 227-242.
- Whitehead PJP, Bauchot M-L, Hureau J-C, Nielsen J, Tortonese E (1984) Fishes of the north-eastern Atlantic and Mediterranean. United Nations Educational, Scientific, and Cultural Organization, Paris.
- Woodstock M (2018) Trophic Ecology and Parasitism of a Mesopelagic Fish Assemblage. Master's thesis. Nova Southeastern University.
- Xavier JC, Vieira C, Assis C, Cherel Y, Hill S, Costa E, Borges TC, Coelho R (2012) Feeding ecology of the deep-sea lanternshark *Etmopterus pusillus* (Elasmobranchii: Etmopteridae) in the northeast Atlantic. *Scientia Marina* 76: 301-310.
- Yang MS (2011) Diet of Nineteen Mesopelagic Fishes in the Gulf of Alaska. Alaska Fisheries Science Center, US Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service.

- Young JW, Lansdell MJ, Campbell RA, Cooper SP, Juanes F, Guest MA (2010) Feeding ecology and niche segregation in oceanic top predators off eastern Australia. *Marine Biology* 157: 2347-2368.
- Young RF, Winn HE (2003) Activity patterns, diet, and shelter site for two species of Moray Eels, *Gymnothorax moringa* and *Gymnothorax vicinus*, in Belize. *Copeia* 2003(1): 44-55.
- Zaidi R, Derbal F, Kara MH (2017) Temporal and ontogenic variations of diet of the goldblotch grouper *Epinephelus costae* (Serranidae) in the eastern coast of Algeria. *Journal of the Marine Biological Association of the United Kingdom* 97: 259-267.
- Zino F, Biscoito M (1994) Breeding seabirds in the Madeira archipelago. Seabirds on islands: threats, case studies and action plans. BirdLife International, Cambridge, United Kingdom, 172-185.
- Zino PA, Zino FJ (1986) Contribution to the study of the petrels of the genus *Pterodroma* in the archipelago of Madeira. *Bol. Mus. Mun. Funchal* 38: 141-165



## CHAPTER 7

---



Credits: Joana Romero

## General Discussion

The studies in this thesis have contributed to the existing knowledge on oceanic regions by focusing on the diet and trophic interactions between some iconic species and groups found in the waters of the archipelago of Madeira. In Chapter 2, we described the diet of the two most abundant small pelagic species, the Atlantic chub mackerel *Scomber colias* and the Blue jack mackerel *Trachurus picturatus*, which also allowed us to have a first insight on the most common and abundant planktonic species in the region. In Chapters 3 and 4, we studied how top predators, namely tunas (3) and gulls (4), use the available marine resources around Madeira. In Chapter 5, we explored the interannual variability of the oceanic epipelagic fish and squid community through the study of the diet and foraging behaviour of the Cory's shearwater. Finally, in Chapter 6, we built a mass-balance model of the oceanic region of Madeira to study the structure and dynamics of its food web.

### Diet, Prey abundance and Community assessment

Diet studies are important for the understanding of the ecology of species. Not only it reveals the prey composition, but also reflects the type of habitats used by the consumer and its foraging behaviour. The study of the diet of an organism also allows to explore interactions with other organisms or even with humans (Brodeur et al. 2008, Wallace et al. 2009), and to describe the prey community (e.g. Velarde et al. 1994) and its fluctuations (e.g. Velarde et al. 2015).

Despite the importance of tunas to the fisheries industry, our knowledge on their ecology is still very limited, especially in the Atlantic, where the feeding behaviour, vertical movements and migratory routes are some of the less studied topics. Tunas are difficult to observe or sample, and therefore more difficult to study, which is why collaborations with fishermen are important. Fishermen are key elements in the study of species targeted by fisheries, as *connoisseurs* of good sampling sites, and of how to find and capture those species more efficiently, which allows to achieve sufficient sample sizes. Still, some techniques used by fishermen might raise challenges in the study of the diet of tunas, or other species. In Madeira, and in a few other regions like the Azores and Canary Islands (Zeller et al. 2001), tunas are captured using live bait. The use of bait, together with showers, mimic the feeding frenzy phenomenon, which attracts tunas and facilitates their capture. However, during these events, tunas are likely to eat bait, which can bias the analysis of stomach contents. The analysis of the concentration of mercury

in the tissues of Bigeye and Skipjack tunas, and of two other predators, indicated that the Bigeye tuna did not have a mesopelagic diet, unlike in the north Pacific or equatorial Atlantic ocean (e.g. Ménard & Marchal 2003, Ohshimo et al. 2018). Instead, its diet was mostly composed by Atlantic chub mackerel and Blue jack mackerel (Chapter 3), depending to a large extent on epipelagic waters.

As mentioned before, diet studies are useful tools to assess prey communities. As such, top predators like seabirds and tunas, may constitute valuable sampling agents with associated low costs. They further provide near real-time information on the relative abundance and availability of commercial, non-commercial and unsurveyed species, extending sampling possibilities to less accessible organisms (Ainley et al. 1986). The conclusions on the diet of tunas (Chapter 3), allied to data collected on the diet and foraging areas of Cory's shearwaters (Chapter 5) and biomass estimations used to build the mass-balanced food web model (Chapter 6), allowed to conclude that among the intermediate trophic levels of Madeira, the most abundant groups in the pelagic ecosystem were myctophids, mackerels and flying squids. These were also the main groups sustaining other upper trophic levels and top predators, like marlins (e.g. Veiga et al. 2011), seabirds (e.g. Waap et al. 2017), and cetaceans (e.g. Fernández et al. 2009). Mesopelagic fish are among the most abundant fish families in these waters (Chapter 6), mainly represented by myctophids, like lantern fish of the genus *Diaphus* sp. and *Hygophum* sp. (Chapter 3). Blue jack and Atlantic chub mackerels are the most abundant epipelagic species in the region of Madeira (Chapter 3 and 5), together with Pilot fish (Chapter 5). The collection of regurgitations and GPS data of Cory's shearwaters over several years (Chapter 5), further revealed an increase in the population of Longspine snipefish *Macroramphosus scolopax* around the archipelago of Madeira in 2017/2018, suggesting a shift in the ecosystem of this oceanic region. Among cephalopods, the Neon and the European flying squids (Ommastrephidae) and Hooked squids (Onychoteuthidae), seem to be the most commonly distributed, according to the diet of tunas (Chapter 3) and Cory's shearwaters (Chapter 5).

The Atlantic chub mackerel and the Blue jack mackerel are the third most important sources of income for fishermen in Madeira, right after tunas and Black scabbardfish (*Aphanopus carbo* and *A. intermedius*; Hermida & Delgado 2016), and are also among the most important prey for top predators in the pelagic ecosystem. Describing their main prey is key to understand what sustains their populations. Both species revealed to be planktonic and piscivorous, feeding on dinoflagellates (*Lingulodinium polyedra* and

*Scrippsiella* sp.), but also on diatoms of the family Bacillariophyceae and Pennales, and coccolithophores. They also fed on zooplanktonic species, like calanoid and cyclopoid copepods (*Candacia* sp., *Pleuromamma* sp., and *Oncea* sp.), and on very small Atlantic saury, clupeids, Longspine snipefish, myctophids and chub mackerels (Chapter 2). These results revealed the importance of long-term studies in marine environments. A previous study by Costa et al. (2013) briefly described the diet of Blue jack mackerel as being composed by fish and euphausiids. Although it is not possible to compare the fish species consumed, as Costa et al. (2013) did not identify them further, the planktonic species found in the mackerel's diet appear to be result of opportunism. In the face of shifts in the planktonic community, Blue jack mackerel seems capable to adapt its diet according to prey availability. The generalist and opportunistic behaviour of the Atlantic chub mackerel also suggests that shifts in its prey abundance and composition should not affect the population of Madeira significantly, which might indicate some resilience by these two species.

We also investigated the diet and foraging behaviour of the Yellow-legged gull in the archipelago of Madeira to understand how this top predator uses its marine environment (Chapter 4). Our study revealed that the Yellow-legged gull has an overall terrestrial behaviour with strong associations to anthropogenic sites and activities, like fisheries, but with low interaction with the marine environment. When at sea, around half of the trips were associated with purse-seine vessels. The low use of the marine environment by gulls in this region seems to be mostly related to the high availability of human refuse but also with a small fleet of purse-seiners and other near shore fisheries.

### **Food web structure and Fisheries**

The oligotrophic oceanic region of the archipelago of Madeira has its overall biomass concentrated on the lower trophic levels, i.e. primary producers, zooplanktonic species, molluscs, suspension filter-feeders, and detritivores (Chapter 6), which are also the base of the food web and through which most of energy and matter flow. The food web was characterized by a more linear-like food chain in opposition to a more web-like food chain, with a big proportion of specialist organisms, like dolphins, shearwaters, and large pelagic fish. These species fed mainly on small pelagic and mesopelagic fish and cephalopods. Despite the low mean trophic level of the system, the mean trophic level of fisheries was 4.2, targeting mainly top predators, like tunas and Black scabbardfish, which



were also considered key components of the ecosystem, and with the most impact (Chapter 6).

In Madeira, top predators occur in large numbers and show high diversity. There, one can find a total of eight species of seabirds (Meirinho et al. 2014), 29 species of cetaceans (seven whales and 22 dolphins) (Freitas et al. 2012), five species of tunas (Gouveia et al. 2019), 27 species of sharks (both pelagic and deep-sea sharks; Biscoito et al. 2018), and many other predator fish, including the Blue and White marlins and the Black scabbardfish. According to our study in Chapter 6, top predators like sharks, dolphins and large demersal fish, were identified as important keystone species, meaning that they have a significant impact in the ecosystem, despite their low abundance, shaping the dynamics and functioning of food webs (Paine 1995). Our model suggested that the ecosystem should remain in equilibrium if fisheries are to continue exploiting the marine resources of Madeira at the same rate. Even so, it has been showed that the selective removal of certain species and size ranges will eventually deplete stocks and change the structure of the ecosystem (Zhou et al. 2010). For the scabbardfish, this can imply the disappearance of its population, as Madeira and the Canary Islands are its spawning grounds (Farias et al. 2013). For tunas, which are migratory species, it will ultimately depend mostly on the status of the population in the spawning and nursing areas. Since 1950, the total biomass of Bigeye tuna in the Atlantic has decreased 73%, of which 53% was in the last 30 years (ICCAT 2018). Although total landings have decreased since 1994, when catches reached a maximum of 134,933 tonnes, since 2006 catches have been slowly increasing again (ICCAT 2018). This is worrisome as ICCAT (International Commission for the Conservation of Atlantic Tunas) has defined this stock as overexploited. Similar decreasing and increasing trends of biomass and fisheries, respectively, have also been observed for Skipjack tunas in eastern Atlantic (ICCAT 2014). If the exploitation rate in the feeding areas remain unchanged, the populations of these three species will certainly continue to decrease and will have significant impacts on the structure of the food web, most probably impacting the epipelagic and deep-sea ecosystems, and consequently also impacting the economy of the archipelago of Madeira.

Fisheries do not only affect their targeted species, but can indirectly affect other species, like gulls. We have seen that gulls have a majorly terrestrial behaviour in Madeira. However, if for instance, the purse-seine fleet would increase in response to the depletion of stocks of higher trophic levels, or if regulatory measures would be applied to fisheries and/or to the main foraging area of gulls (landfills), Yellow-legged gulls could

be expected to increase their interaction with purse-seine vessels. There is a likelihood that gulls would also increase predation of small petrels. If these behaviour shifts were to be observed, gulls would consequently use more the marine environment and have a more substantial role in the marine food web of Madeira than presently.

Fisheries and climate change are known to cause alterations on life-traits of organisms, such as the reduction of their average length (Heino & Godø 2002, van Rijn et al. 2017), which can cause size-based trophic mismatches and perturbations on the prey-predator relationships. For instance, if the defence mechanism of a prey would be its size, then its reduction would increase the predation rate and, consequently, decrease its abundance. These are some of the consequences that can alter the types and strengths of flows of energy and matter in ecosystems, and impact the ecological networks and the ecosystem structure and functioning (Parmesan & Matthews 2006). Continued efforts to try to reduce pressure on fisheries and mitigate the consequences of climate change are needed to avoid irreversible changes in the structure of the ecosystem which cannot be compensated by its adaptation capacity.

## **Future research**

This thesis has revealed new information on the diet and foraging behaviour of some iconic species from the archipelago of Madeira and has allowed us to have a better understanding of the trophic web of this ecosystem. But as expected, it also raises new questions to be answered. The study of the diet of the Bigeye tuna has revealed that in the archipelago of Madeira, this species is mainly feeding on epipelagic species, while it feeds on mesopelagic species elsewhere. Why would a species that tends to spend most of its time at higher depths, according to other studies (Brill et al. 2005, Junior et al. 2012), would feed on epipelagic species in the oceanic region of Madeira? This feeding behaviour should be further investigated focusing on the vertical movements of the tuna, but also on oceanographic phenomena which can affect their behaviour, and on the behaviour of their prey in Madeira. Long-term studies on the diet of Bigeye tuna are also important to better understand its feeding ecology.

In Chapter 5, we used the diet and foraging movements of the Cory's shearwater to show that very sudden and significant variations in this community can occur, as was the case of the appearance of the Longspine snipefish in the Selvagens region, which very quickly assumed a top position as prey for these seabirds in the region. Large scale and annual oceanographic variables did not explain the snipefish fluctuations. Further studies

on Longspine snipefish population dynamics and breeding biology in regions where its presence is more common, such as studies of oceanographic phenomena in the oceanic region of Madeira, may help shed more light on the drivers of these fluctuations. As done in chapter 5, it would be interesting to use other top predators, like the Bulwer's petrel or the Black scabbardfish, to describe the prey communities of deeper layers.

The Ecosim with Ecopath software, used to model the food web of Madeira, showed that there is still a lot to know about this ecosystem. Several species, mainly deep-sea species, have not been properly catalogued or sampled, which means that information on their biomass, diet, or reproduction biology are still unknown. Biomass was the most frequently missing parameter in the model, which shows the need for studies estimating population sizes.

## References

- Ainley DG, Fraser WR, Sullivan CW, Torres JJ, Hopkins TL, Smith WO (1986) Antarctic mesopelagic micronekton: evidence from seabirds that pack ice affects community structure. *Science* 232:847–849
- Biscoito M, Ribeiro C, Freitas M (2018) Annotated checklist of the fishes of the archipelago of Madeira (NE Atlantic): I-Chondrichthyes. *Zootaxa* 4429:459–494
- Brill RW, Bigelow KA, Musyl MK, Fritsches KA, Warrant EJ (2005) Bigeye tuna (*Thunnus obesus*) behavior and physiology and their relevance to stock assessments and fishery biology. *Col Vol Sci Pap ICCAT* 57:142–161
- Brodeur RD, Suchman CL, Reese DC, Miller TW, Daly EA (2008) Spatial overlap and trophic interactions between pelagic fish and large jellyfish in the northern California Current. *Mar Biol* 154:649–659
- Costa G, Santamaria MTG, Vasconcelos J, Perera CB, Melo-Moreira E (2013) Endoparasites of *Trachurus picturatus* (Pisces: Carangidae) from the Madeira and Canary Islands: Selecting parasites for use as tags. *Sci Mar* 77:61–68
- Farias I, Morales-Nin B, Lorange P, Figueiredo I (2013) Black scabbardfish, *Aphanopus carbo*, in the northeast Atlantic: Distribution and hypothetical migratory cycle. *Aquat Living Resour* 26:333–342
- Fernández R, Santos MB, Carrillo M, Tejedor M, Pierce GJ (2009) Stomach contents of cetaceans stranded in the Canary Islands 1996–2006. *J Mar Biol Assoc United Kingdom* 89:873–883

- Freitas L, Dinis A, Nicolau C, Ribeiro C, Alves F (2012) New records of cetacean species for Madeira archipelago with an updated checklist. *Bol do Mus Munic do Funchal (História Nat)* 62:25–43
- Gouveia L, Amorim A, Alves A, Hermida M (2019) Updated fishery statistics of tuna species caught off Madeira archipelago. *Collect Vol Sci Pap ICCAT* 75:2081–2097
- Heino M, Godø OR (2002) Fisheries-induced selection pressures in the context of sustainable fisheries. *Bull Mar Sci* 70:639–656
- Hermida M, Delgado J (2016) High trophic level and low diversity: Would Madeira benefit from fishing down? *Mar Policy* 73:130–137
- ICCAT (2014) Report of the 2014 ICCAT East and West Atlantic Skipjack stock assessment meeting. pp 98
- ICCAT (2018) Report of the 2018 ICCAT Bigeye tuna stock assessment meeting. pp 92
- Junior T V., Travassos PE, Hazin FH V., Tolotti MT, Barbosa TM (2012) Forage fauna in the diet of Bigeye tuna (*Thunnus obesus*) in the Western tropical Atlantic Ocean. *Brazilian J Oceanogr* 60:89–97
- Meirinho A, Barros N, Oliveira N, Catry P, Lecoq M, Paiva V, Geraldés P, Granadeiro JP, Ramírez I, Andrade J (2014) Atlas das aves marinhas de Portugal. Sociedade Portuguesa para o Estudo das Aves. Lisboa
- Ménard F, Marchal E (2003) Foraging behaviour of tuna feeding on small schooling *Vinciguerria nimbaria* in the surface layer of the equatorial Atlantic Ocean. *Aquat Living Resour* 16:231–238
- Ohshimo S, Hiraoka Y, Sato T, Nakatsuka S (2018) Feeding habits of bigeye tuna (*Thunnus obesus*) in the North Pacific from 2011 to 2013. *Mar Freshw Res* 69:585–606
- Paine RT (1995) Diversity on Refining the Concept of Keystone. *Conserv Biol* 9:962–964
- Parmesan C, Matthews J (2006) Biological Impacts of Climate Change. In: *Principles of Conservation Biology*. p 333–374
- Rijn I van, Buba Y, DeLong J, Kiflawi M, Belmaker J (2017) Large but uneven reduction in fish size across species in relation to changing sea temperatures. *Glob Chang Biol* 23:3667–3674
- Veiga P, Xavier JC, Assis C a., Erzini K (2011) Diet of the blue marlin, *Makaira nigricans*, off the south coast of Portugal. *Mar Biol Res* 7:820–825

- Velarde E, Ezcurra E, Anderson DW (2015) Seabird diet predicts following-season commercial catch of Gulf of California Pacific Sardine and Northern Anchovy. *J Mar Syst* 146:82–88
- Velarde E, Tordesillas M de la S, Rocio Esquivel LV (1994) Seabirds as indicators of important fish populations in the Gulf of California. *CalCofi Rep* 35:137–143
- Waap S, Symondson WOC, Granadeiro JP, Alonso H, Serra-Gonçalves C, Dias MP, Catry P (2017) The diet of a nocturnal pelagic predator, the Bulwer’s petrel, across the lunar cycle. *Sci Rep* 7
- Wallace BP, Avens L, Braun-McNeill J, McClellan CM (2009) The diet composition of immature loggerheads: Insights on trophic niche, growth rates, and fisheries interactions. *J Exp Mar Bio Ecol* 373:50–57
- Zeller D, Watson R, Pauly D (2001) Fisheries Impacts on North Atlantic Ecosystems: Catch, Effort and National/Regional Data sets.
- Zhou S, Smith ADM, Punt AE, Richardson AJ, Gibbs M, Fulton EA, Pascoe S, Bulman C, Bayliss P, Sainsbury K (2010) Ecosystem-based fisheries management requires a change to the selective fishing philosophy. *Proc Natl Acad Sci U S A* 107:9485–94

